

Isolation of Active Compound from Zingiber Purpureum Roxb. Using Bioassay Guided Fractination Method for WIDR Colon Adenocarcinoma Cell Line

by MonicaEva

Submission date: 06-May-2021 10:34PM (UTC-0400)

Submission ID: 1580142674

File name: Compound_from_Zingiber_Purpureum_Roxb._Using_Bioassay_Guided.pdf (366.58K)

Word count: 3425

Character count: 16690

12

Isolation of Active Compound from *Zingiber Purpureum* Roxb. Using Bioassay Guided Fractination Method for WIDR Colon Adenocarcinoma Cell Line

Rollando Rollando, Maureen Engracia, Laurensia Nina Irawati, Edward Hartono, Eva Monica, Rehmadata Sitepu

16

Program of Pharmacy, Faculty of Science and Technology, Ma Chung University, Malang Villa Puncak Tidar N-01, Malang 65151.

23

*Corresponding Author: Email: ro.llando@machung.ac.id

Abstract

Zingiber purpureum Roxb. was a plant that has the potential to be developed as a chemotherapy agent. Bioassay guided fractination aims to examine the anticancer activity of n-hexane, ethyl acetate, and methanol fractions from ethanolic extract of bangle plant rhizome to colon cancer cell type WiDr and Vero cells. The apoptotic test and modulating cell cycle were observed by flow cytometry method. Isolation process of the most active fraction. The pure isolates obtained were identified by structures with structural elucidation using ultraviolet, 1D-NMR, 2D-NMR and liquid chromatography-mass spectrometer. Obtained 2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one compound of ethyl acetate fraction number 21. With a value of CC_{50} to a WiDr cell of $11.24 \pm 3.44 \mu\text{g/mL}$ and $342.22 \pm 4.98 \mu\text{g/mL}$ to Vero cells.

Keywords: *Zingiber purpureum* Roxb., WiDr, Vero, 3-dihydro-5-hydroxy-7-methoxy-2 phenylchromen-4-one

Introduction

Colorectal cancer ranked fourth in all men and women worldwide, with an incidence of 1,400,000 new cases diagnosed in 2012 [1]. IARC states that cancers of the colon and rectum are the most common incidence of cancer with the number 3 (34,000 per year, 15.9%) and the third cause of death (18,000 per year, 10.8%). Chemotherapy was one of the therapies in the treatment of cancer. Chemotherapy agents generally have low selectivity properties because they are antiproliferative to normal selkanker and cell [2]. In addition, some chemotherapy has a narrow therapeutic index, and may lead to multidrug resistance (MDR) and adverse side effects [3].

One of the problems that often arise in the treatment of cancer is the resistance of chemotherapy drugs (drug resistance). Doxorubicin is a chemotherapy agent of various types of cancer of the anthracycline class that has provoked resistance. In addition to causing resistance, doxorubicin may also cause cardiotoxicity in long-term

use [4]. Increased doses of doxorubicin are not an appropriate way to overcome resistance problems because they can cause normal cells to be exposed by cytotoxic drugs that will trigger toxicity and transform normal cells into cancer cells [5].

One of the medicinal plants in Indonesia that potentially developed as a chemopreventive agent in colon cancer is bangle. The bangle plant has been used by society traditionally to cope with several diseases. Empirically, boiled water rhizome bangle used the community as a headache medicine, constipation drugs, abdominal pain, jaundice, as body warmers, slimming, helps the gas out of the digestive tract, asthma, rheumatism, and antipyretics.

The research on scientific proof of the efficacy of the rhizome of bangle plant is still minimal publication and the efficacy of rhizome of bangle plant still many that have not been explored. So in this study, conducted an active fraction search on the ethanolic extract

of rhizome plant bangle that can kill and suppress the growth of cancer cells, especially in colon cancer cell type WiDr.

Material and Instrument

Methanol, Hexane and ethyl acetate fraction of faloak bark, DMSO 0,1%, cisplatin (Wako), High glucose DMEM (Dulbecco's Modified Eagle Media) (Gibco), (FBS 10% (v/v) (Qualified FBS, Gibco, Invitrogen USA), penicillin-streptomycin 1,5% (v/v) (Gibco, Invitrogen USA and Fungizone 0,5% v/v Gibco). Trypsin-EDTA 0,25% (Gibco, Invitrogen Canada), 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma Aldrich, USA). MTT was prepared with a concentration of 5 mg/mL dissolved in phosphate buffer saline (PBS) 1 x pH 7, 4. 0, 01 N HCl (Merck, Darmstadt, Germany). PBS containing 1 mg/ml (minimum 95% (HPLC), Sigma-Aldrich Co., St. Louise, MO, 63178, USA) 10 mg/ml RNase (obtained from Laboratory of Animal Sciences, NAIST, Japan) and Triton X-100 for GC, E. Merck, 64271, Darmstadt, Germany).

Liquid nitrogen, labconco purifier class II biosafety cabinets (Delta Series, Labconco Corporation, Missouri, USA), cell counter, micropipet (Pipetman® la Gilson, France), digital camera (Sony), centrifuge (Sigma 203, B.Braun Biotech International), digital balance sheets (Mettler Toledo, AG204 Delta Rang®), mixer (Maxi Mix II, Thermolyne type 37600 mixer, Iowa, USA), oven (Mettmert), ELISA reader (Bio-Rad microplate reader Benchmark serial No. 11565, Japan), FACTS calibur flow cytometer. FTIR (FTIR-100 Perkin Elmer), MS (Mariner Biospectrometry System HRESIMS) and NMR spectrometer (Delta 2 400MHz for ¹H-NMR and 100 MHz for ¹³C-NMR). Autoclave (Hirayama HV 25 020585175, Hirayama Manufacturing Co., Japan), inverted microscope (Nikon, Eclipse, TE 2000-U), hemocytometer (Nebauer improved 0.100 mm Tiefe Depth Profondeur 0.0025 mm², Germany), stirrer (Nuova, Thermolyne).

Procedure

Plant Material

Collected plant sample was washed with running tap water and shade dried at room temperature for three weeks and grounded. The plant material (250 g) was successively extracted ethanol with maceration method.

All extractives were filtered through Whatman filter paper (No: 1) and concentrated.

Fractionation with Column Chromatography

Separation of the active compound by column chromatography using silica gel GF₂₅₄ as stationary phase and the 200 mL n-hexane, ethyl acetate and methanol as mobile phase was carried out gradually with the addition of 1 drop of CH₃COOH. Each fraction obtained was evaporated at ± 40 °C to obtain a concentrated fraction and then dried in an oven at ± 50 °C. Each fraction tested to the cytotoxic activity to obtain the value of CC₅₀.

Cytotoxicity Assay

Cells were washed twice with PBS and inoculated with 200 µL of each fraction diluted in DMEM supplemented with 2% FCS (two-fold dilutions, ranging from 2000 µg to 0.975 µg). Cells were allowed to grow for 72 h at 37 °C 5% CO₂. Then, medium was removed, cells were washed in PBS and 20 µL of MTT solution at 5 mg/ml (MTT, Sigma, Saint Louis, MO, USA) were added to each well. Plates were incubated for 4 h at 37 °C 5% CO₂ and the formed formazan crystals were solubilized in DMSO (Sigma, USA). The absorbance of formazan-generated dye was determined using a microplate reader (Thermo Fisher Scientific, USA) at 570 nm. The % of viability was calculated using the following formula:

$$\%V = \frac{\text{Absorbance of treated cells}}{\text{Absorbance of control cells}}$$

The viability of control cells was set to 100%, the CC₅₀ value was derived from the corresponding dose-response curves as the concentration of the oil that reduced cell viability by 50%.

Isolation and Purification under Bioassay-Guided Screening

According to cytotoxicity test the compounds were identified by comparing their UV, infrared, LC-MS and NMR data with those of published data test, only one fraction was the most toxic (ethyl acetate fraction 21). The fraction were separated by Sephadex LH-20 using CH₂Cl₂: MeOH (4:7). Combined fractions, according to TLC analysis, were purified by preparative TLC using a mixture of solvents.

Determination of Cell Cycle and Apoptosis

WiDr cells were grown on glass coverslips in tissue culture dishes (Falcon) and were allowed to attach for 24 hours prior to the addition of any drug. After the cells were incubated with a drug for 24 hours, the coverslips were washed once in PBS and fixed in object glass. Treated cells were stained with acridine orange and ethidium bromide 5 μ L and visualized by fluorescence microscopy.

Result and Discussion

Results of Cytotoxic Test

The cytotoxic activity of the fraction was expressed as CC_{50} . The test was performed by dissolving 44 fractions of fractionation of ethanolic extract with gradient system (n-

hexane, ethyl acetate and methanol), then into it added DMSO solution until obtained the desired concentration. A single cytotoxic effect showed decreased cell viability and morphological changes in colon cancer cells of WiDr. The test results (Fig.1) show that Ethyl Acetate Fraction 21 treatment (Fig. 1b-c) shows the wrinkled cell. This indicates a decrease in living cells when compared to controls (Fig. 1a).

The cells appear round and fragmented to indicate a change in cell morphology, but unknown changes were due to cell deaths due to necrotic or apoptotic processes and proliferative inhibition processes. Meanwhile, the CC_{50} value of the largest was ethyl acetate fraction 21 with concentrations of 11.24 ± 3.44 μ g/mL and doxorubicin was 13.88 ± 0.34 μ g/mL.




Fig.1: Ethyl Acetate Fraction 21 testing was performed using cell controls (a) and Vero cells (d) Effect of Ethyl Acetate Fraction 21 on WiDr colon cancer cells (b-c). Figure (b) at concentrations of 200 μ g/mL and image (c) at a concentration of 300 μ g /mL. The Vero cells were treated with a fraction of concentration of 200 μ g/mL.

Table 1: Cytotoxic activity on WiDr and Vero

Number	Experiment	CC ₅₀ on WiDr Cell	CC ₅₀ on Vero Cell
1	Doxorubicin	13,88±0,34 µg/mL	160,67±0,76 µg/mL
2	Methanol fraction 1	56,54 ±1,67 µg/mL	240,89 ± 0,96 µg/mL
3	Methanol fraction 2	70,88 ±0,92 µg/mL	140,92 ± 1,22 µg/mL
4	Methanol fraction 3	31,22 ±2,22 µg/mL	341,67 ± 2,39 µg/mL
5	Methanol fraction 4	48,88 ±0,88 µg/mL	131,33 ± 1,96 µg/mL
6	Methanol fraction 5	74,38 ±0,55 µg/mL	402,43 ± 1,07 µg/mL
7	Methanol fraction 6	99,68 ±0,87 µg/mL	148,56 ± 3,84 µg/mL
8	Methanol fraction 7	58,33 ±1,99 µg/mL	340,44 ± 1,45 µg/mL
9	Ethyl Acetate Faction 1	41,11 ±2,68 µg/mL	441,44 ± 0,34 µg/mL
10	Ethyl Acetate Faction 2	66,78 ±1,03 µg/mL	561,90 ± 1,52 µg/mL
11	Ethyl Acetate Faction 3	36,99 ±0,86 µg/mL	445,95 ± 1,14 µg/mL
12	Ethyl Acetate Faction 4	98,82 ±3,35 µg/mL	123,98 ± 3,22 µg/mL
13	Ethyl Acetate Faction 5	95,85 ±3,08 µg/mL	248,07 ± 2,58 µg/mL
14	Ethyl Acetate Faction 6	88,55 ±2,88 µg/mL	338,77 ± 4,95 µg/mL
15	Ethyl Acetate Faction 7	13,11 ±2,08 µg/mL	565,44 ± 1,08 µg/mL
16	Ethyl Acetate Faction 8	36,88 ±1,07 µg/mL	341,69 ± 0,11 µg/mL
17	Ethyl Acetate Faction 9	32,54 ±1,91 µg/mL	249,22 ± 0,65 µg/mL
18	Ethyl Acetate Faction 10	16,22 ±0,79 µg/mL	241,53 ± 0,96 µg/mL
19	Ethyl Acetate Faction 11	99,11 ±2,07 µg/mL	299,19 ± 4,98 µg/mL
20	Ethyl Acetate Faction 12	88,19 ±5,88 µg/mL	349,93 ± 0,96 µg/mL
21	Ethyl Acetate Faction 13	14,33 ±4,33 µg/mL	240,11 ± 1,93 µg/mL
22	Ethyl Acetate Faction 14	88,11 ±3,34 µg/mL	360,24 ± 5,11 µg/mL
23	Ethyl Acetate Faction 15	16,92± 2,88 µg/mL	130,11 ± 8,22 µg/mL
24	Ethyl Acetate Faction 16	28,22 ±3,30 µg/mL	223,11 ± 2,30 µg/mL
25	Ethyl Acetate Faction 17	92,22 ±3,93 µg/mL	292,12 ± 2,33 µg/mL
26	Ethyl Acetate Faction 18	31,21 ±3,33 µg/mL	234,44 ± 4,33 µg/mL
27	Ethyl Acetate Faction 19	39,92 ±4,22 µg/mL	234,11 ± 3,43 µg/mL
28	Ethyl Acetate Faction 20	21,22 ±2,22 µg/mL	134,21 ± 3,11 µg/mL
29	Ethyl Acetate Faction 21	11,24 ±3,44 µg/mL	342,22 ± 4,98 µg/mL
30	Ethyl Acetate Faction 22	41,98 ±9,92 µg/mL	232,22 ± 1,92 µg/mL
31	Ethyl Acetate Faction 23	34,22 ±2,33 µg/mL	234,03 ± 3,22 µg/mL
32	Ethyl Acetate Faction 24	92,23 ±2,22 µg/mL	445,11 ± 2,39 µg/mL
33	Ethyl Acetate Faction 25	82,11 ±3,98 µg/mL	354,73 ± 1,39 µg/mL
34	Hexane Fraction 1	77,92 ±2,93 µg/mL	551,82 ± 3,11 µg/mL
35	Hexane Fraction 2	98,28 ±3,34 µg/mL	458,11 ± 1,28 µg/mL
36	Hexane Fraction 3	90,01 ±2,88 µg/mL	388,29 ± 3,29 µg/mL
37	Hexane Fraction 4	34,22 ±1,98 µg/mL	123,21 ± 2,92 µg/mL
38	Hexane Fraction 5	22,11 ±2,62 µg/mL	220,91 ± 0,16 µg/mL
39	Hexane Fraction 6	97,14 ±2,62 µg/mL	240,11 ± 0,22 µg/mL
40	Hexane Fraction 7	36,22 ±4,11 µg/mL	233,11 ± 0,92 µg/mL
41	Hexane Fraction 8	90,11 ±2,66 µg/mL	102,11 ± 0,81 µg/mL
42	Hexane Fraction 9	83,29 ±2,98 µg/mL	358,09 ± 0,12 µg/mL
43	Hexane Fraction 10	73,12 ±2,93 µg/mL	241,72 ± 0,02 µg/mL
44	Hexane Fraction 11	99,54 ±1,67 µg/mL	229,11 ± 0,88 µg/mL
45	Hexane Fraction 12	96,11 ±2,22 µg/mL	221,11 ± 0,11 µg/mL

Structure Elucidation

Compound 1, yellow crystal, m.p. 97-99 °C, UV (MeOH) λ_{maks} nm (log ϵ): 232 (3.92), 290 (4.10), dan 325 sh (3.66) nm. EIMS: m/z (270, M⁺, 100, base peak), 193 (76.1), 166 (80.3), 138 (41.4), 103 (21.2), and 77 (16.4). ¹H NMR (400 MHz in acetone *d*6), δ_{H} ppm: 12.17 (1H, *br, s*, 5-OH), 7.58 (3H, *m*, H-3',4',5'), 7.41 (2H, *m*, H-2',6'), 6.09 (1H, *d*, $J=2.2$ Hz, H-8), 6.07 (1H, *d*, $J=2.2$ Hz, H-6), 5.61 (1H, *dd*, $J=3.8$; 12.3 Hz, H-2), 3.35 (3H, *s*, 7-OCH₃),

3.24 (1H, *dd*, $J=12.8$; 16.4 Hz, H-3_{ax}), and 2.81 (1H, *dd*, $J=3.8$; 16.4 Hz, H-3_{eq}). ¹³C NMR (100 MHz in acetone *d*6), δ_{C} ppm: 196.3 (C-4), 168.6 (C-7), 164.8 (C-5), 163.3 (C-8a), 140.1 (C-1'), 129.6 (C-3',5'), 128.5 (C-4'), 123.9 (C-2',6'), 101.5 (C-4a), 95.09 (C-8), 95.66 (C-6), 79.6 (C-2), 56.1 (7-OCH₃), and 43.9 (C-3). The elucidation results showed that the compound of the ethyl acetate fraction 21 was 2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one.




Fig. 2: 2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one Compound (HMBC)

Observation of Apoptosis and the Cell Cycle

Furthermore, modulation test of cell cycle. The process of DNA synthesis in cancer cells through the same cell cycle as normal cells [6]. One of the main targets in inhibiting cancer cell proliferation is the modulation of cell cycle that can be observed using flowcytometry method (Rollando, 2000).

Flowcytometry is able to detect each phase in the cell cycle based on the number of chromosomes on each phase (G1, S, and G2 / M) [7]. Propidium iodide is used to color each phase because it is capable of interacting with DNA [8]. Observation of cell cycle profile was done at 24 hours [9]. The percentage of cell cycle distribution in detail is shown in Table 2.

Table 2: Percentage of cell cycle distribution

Sample	G1 (%)	S (%)	G2/M (%)	% CV
Control	38,11	12,34	13,34	3,87
Doxorubicin 2,5 μM	42,88	13,11	24,55	2,22
2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one 2,5 μM	34,11	23,11	11,23	5,98




Fig. 3: Distribution of cell cycle percentages. (a) Control (b) Doxorubicin (c) 2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one Compound

The test results showed doxorubicin causing accumulation in phase S. The 2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one

compound causes cell accumulation in phase S when compared with cell control. In the combination treatment the percentage of cell

cycle distribution in the S phase amounted to 23.11% greater than the single doxorubicin of 13.11%. Cell accumulation in the S-phase of combination was increased compared with

cells without treatment (cell control) of 12.34%. Cell accumulation is possible because of cell cycle arrest in that phase.

Table 3: Percentage of death after treatment

	Control	Doxorubicin 2.5 µM	2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one 2.5 µM
Initial apoptosis (%)	2,11	2,33	9,87
End apoptosis (%)	1,63	2,53	1,24
Necrosis (%)	0,86	0,87	1,62
Total	3,94	6,04	13,78

Observation of apoptotic compounds 2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one with flow cytometer. Induction of apoptosis was used to investigate the mechanism of cell death from treatment of the 2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one compound on colon cancer cell WiDr incubated for 24 hours [10].

The method used in this study is Annexin V method that was detected using flowcytometry to see the induction of apoptosis that occurred in the treated cells. Annexin V is a member of the family of phospholipid binding proteins that are on strongly negatively charged cellular membranes. Cell cell death caused by apoptosis or necrosis can be distinguished by the staining of Propidium Iodide (PI) through intercalation with DNA [11]. The percentage of cell death after treatment of 2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one induced by apoptosis or necrosis is shown in Table 3.

Analysis of the percentage of cell death after 2, 3- dihydro- 5- hydroxy-7-methoxy-2-phenylchromen-4-one (Table 3) showed that

control cells showed a 3.94% percentage of cell death. Single-treated cells with doxorubicin showed 6.04% cell death, whereas 2, 3-dihydro-5- hydroxy-7- methoxy-2-phenylchromen-4-one compound resulted in 13.78% cell death. It showed an increase in the percentage of cell death by 7.74% in the 2,3-dihydro- 5-hydroxy- 7-methoxy- 2-phenylchromen-4-one compound compared with doxorubicin.

Conclusion

The results showed that isolate 2 was 3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one having activity cytotoxic in breast cancer cell type T47D with active category with IC₅₀ 11, 24 ±3,44 µg/mL.

Acknowledgements

Gratefully acknowledge for the financial support from Ministry of Research, Technology and Higher Education on Student Creativity Program 2018.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

References

- 1.Yuniati Y, Alfanaar R, Rollando R (2018) Structural conformational study of isoflavon derivatives in soybean using semiempirical methods. Journal of Global Pharma Technology, 10(05):220-225.
- 2.Valeriotte F, Grieshaber CK, Media J, Pietraszkewicz H, Hoffmann J, Pan M (2002) Discovery and development of anticancer agents from plants. J. Exp. Ther. Oncol., 2:228-236.
- 3.Rollando R (2018) Combination of hedyotis corymbosa L. and tinospora crispa ethanolic extract increase cisplatin cytotoxicity on T47D

- 2 breast cancer cells. Asian J Pharm Clin Res, 11(7):171-177.
- 4.Rollando R. Hedyotis corymbosa L. and sterculia quadrifida R. (2018) Br ethanolic extract enhances cisplatin's cytotoxicity on T47D breast cancer cells through cell cycle modulation. The Journal of Pure and Applied Chemistry Research, 7(2):160-172.
- 5.Bravo-Cellar A, Ortiz-Lazareno PC, Lerma-Diaz JM, Dominguez Rodriguez J.r., Jave-Suarez LF, Aguilar-Lemarroy A (2010) Sensitization of cervix cancer cells to adriamycin by

- pentoxyfiline induces an increases in apoptosis and decrease senescence. *Molecular Cancer*, 9:114.
6. Stuart BH (2004) *Infrared Spectroscopy: Fundamentals and Applications.* John Wiley & Sons.
 7. Yuniati Y, Rollando R (2018) Isolation of antibacterial compounds from endophyte fungal of fusarium sp. In phyllanthus niruri linn. Leaves. *J. Pharm. Sci. & Res.* 10(2):260-264.
 8. Rollando R, dan Prilianti KR (200) Fraksi etil asetat kulit batang faloak (*Sterculia quadrifida* R.Br) menginduksi apoptosis dan siklus sel pada sel kanker payudara T47D. *J. Pharm. Sci. Community*;14:1-14.
 9. Masuda T, Ohba S, Kawada M, Osono M, Ikeda D, Esumi H (2006) Antitumor Effect of Kigamicin D on Mouse Tumor Models. *The Journal of Antibiotics*, 59(4):209-211.
 10. Rollando R, Sitepu R, Monica E (2018) Cytotoxic activity of 2- iminoethyl 2-(2-(1-hydroxypentan-2-yl) phenyl) acetate from *sterculia quadrifida* R.Br ethyl acetate fraction. *Journal of Global Pharma Technology*, 10(06):204-212.

Isolation of Active Compound from Zingiber Purpureum Roxb. Using Bioassay Guided Fractination Method for WIDR Colon Adenocarcinoma Cell Line

ORIGINALITY REPORT

16%

SIMILARITY INDEX

7%

INTERNET SOURCES

12%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Barros, Iuri Bezerra de, Marco André Abud Kappel, Priscila Moraes dos Santos, Valdir Florêncio da Veiga Junior, Eliane D'Elia, and Ivan Napoleão Bastos. "The inhibitory action of Bauhinia purpurea extracts on the corrosion of carbon steel in sulfuric acid medium", *Materials Research*, 2016.
Publication 1%
- 2** jpsr.pharmainfo.in
Internet Source 1%
- 3** Luma Majeed Ahmed. "Bulk and Nanocatalysts Applications in Advanced Oxidation Processes", *IntechOpen*, 2020
Publication 1%
- 4** P. B. Schiff, S. B. Horwitz. "Taxol stabilizes microtubules in mouse fibroblast cells.", *Proceedings of the National Academy of Sciences*, 1980
Publication 1%

5

Srinivasan, R., D. Natarajan, M.S. Shivakumar, T. Vinuchakkaravarthy, and D. Velmurugan. "Bioassay guided isolation of mosquito larvicidal compound from acetone leaf extract of *Elaeagnus indica* Servett Bull and its in-silico study", *Industrial Crops and Products*, 2015.

Publication

1 %

6

www.hindawi.com

Internet Source

1 %

7

Hua-Bin Li, Feng Chen, Yue Jiang. "Determination of vitamin B12 in multivitamin tablets and fermentation medium by high-performance liquid chromatography with fluorescence detection", *Journal of Chromatography A*, 2000

Publication

1 %

8

www.jstage.jst.go.jp

Internet Source

1 %

9

Adam Hermawan, Annisa Khumaira, Muthi Ikawati, Herwandhani Putri et al. "Identification of Key Genes of Hesperidin in Inhibition of Breast Cancer Stem Cells by Functional Network Analysis", *Computational Biology and Chemistry*, 2020

Publication

1 %

10	Saoussen Hammami, Habib Jmii, Ridha Mokni, Abdelbaki Khmiri et al. "Essential Oil Composition, Antioxidant, Cytotoxic and Antiviral Activities of Teucrium pseudochamaepitys Growing Spontaneously in Tunisia", <i>Molecules</i> , 2015 Publication	1 %
11	dx.doi.org Internet Source	1 %
12	sintadev.ristekdikti.go.id Internet Source	1 %
13	Ramadan, M.A.. "Minor phenolics from <i>Crinum bulbispermum</i> bulbs", <i>Phytochemistry</i> , 200008 Publication	1 %
14	Sergio Valente, Daniela Trisciuglio, Maria Tardugno, Rosaria Benedetti et al. " - Butylcarbamate-Containing Histone Deacetylase Inhibitors: Apoptosis Induction, Cytodifferentiation, and Antiproliferative Activities in Cancer Cells ", <i>ChemMedChem</i> , 2013 Publication	<1 %
15	ann-clinmicrob.biomedcentral.com Internet Source	<1 %
16	doaj.org Internet Source	<1 %

17

www.oak.go.kr

Internet Source

<1 %

18

Wenyi Gu, Lisa Putral, Nigel McMillan.
"Chapter 12 siRNA and shRNA as Anticancer
Agents in a Cervical Cancer Model", Springer
Science and Business Media LLC, 2008

Publication

<1 %

19

Corona, P.. "Quinoxaline chemistry. Part 11. 3-
Phenyl-2 [phenoxy- and phoxymethyl]-6(7)
or 6,8-substituted quinoxalines and N-[4-(6(7)-
substituted or 6,8-disubstituted-3-
phenylquinoxalin-2-yl)hydroxy or
hydroxymethyl]benzoylglutamates. Synthesis
and evaluation of in vitro anticancer activity
and enzymatic inhibitory activity against
dihydrofolate reductase and thymidylate
synthase", Il Farmaco, 19980730

Publication

<1 %

20

Ilina Krasteva, Stefan Platikanov, Stefan
Nikolov, Maki Kaloga. " Flavonoids from ",
Natural Product Research, 2007

Publication

<1 %

21

Mustafa, K.. "Hydrogen-bonded rotamers of
2',4',6'-trihydroxy-3'-formyldihydrochalcone,
an intermediate in the synthesis of a
dihydrochalcone from *Leptospermum
recurvum*", Tetrahedron, 20030804

Publication

<1 %

22	X. -Y. Guo. "9, 10-Dihydrophenanthrene derivatives from <i>Pholidota yunnanensis</i> and scavenging activity on DPPH free radical", <i>Journal of Asian Natural Products Research</i> , 3/2007 Publication	<1 %
23	jurnal.ugm.ac.id Internet Source	<1 %
24	tessera.spandidos-publications.com Internet Source	<1 %
25	www.jbc.org Internet Source	<1 %
26	www.wjgnet.com Internet Source	<1 %
27	Alexander Baranovsky, Bettina Schmitt, Daniel J. Fowler, Bernd Schneider. "Synthesis of New Biosynthetically Important Diarylheptanoids and Their Oxa- and Fluoro- Analogues by Three Different Strategies", <i>Synthetic Communications</i> , 2003 Publication	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Isolation of Active Compound from Zingiber Purpureum Roxb. Using Bioassay Guided Fractination Method for WIDR Colon Adenocarcinoma Cell Line

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
