

BUKTI KORESPONDENSI

Jurnal Nasional Terakreditasi Peringkat 3 dan 4

Judul Artikel : *Evaluation of Single and Combination Chemotherapy Agents in Patients with Metastatic Breast Cancer*

Jurnal : Jurnal Ilmiah Farmasi Vol. 18 No. 2, 30 Desember 2022

Penulis :

No.	Perihal	Tanggal
1.	Bukti konfirmasi submit artikel	9 November 2021
2.	Pemberitahuan revisi pertama	18 Juni 2022
3.	Submit revisi	19 Juni 2022
4.	<i>Accept Submission</i>	4 Agustus 2022
5.	Proofread	25 Agustus 2022
7.	Final Proofread	12 Desember 2022
8.	Publish Artikel	30 Desember 2022

Bukti Konfirmasi Submit Artikel (9 November 2021)

Jurnal Ilmiah Farmasi

... Back to Submissions

21570 / Aditya et al. / Evaluation of single and combination chemotherapy agents in patients with metastatic breast cancer

Library

Workflow Publication

Submission Review Copyediting Production

Submission Files [Q Search](#)

58157	admin, Breast Cancer Article.docx	November 9, 2021	Article Text
58159	Cover letter	November 9, 2021	Other

[Download All Files](#)

Pre-Review Discussions [Add discussion](#)

Name	From	Last Reply	Replies	Closed
No Items				

Pemberitahuan Revisi Pertama (18 Juni 2022)

Jurnal Ilmiah Farmasi

... Back to Submissions

Notifications ×

[JIF] Editor Decision

2022-06-18 02:39 AM

Godeliva Adriani Hendra:

We have reached a decision regarding your submission to Jurnal Ilmiah Farmasi, "Evaluation of Single and Combination Chemotherapy Agents in Patients with Metastatic Breast Cancer".

Our decision is: Revisions Required

_____ Jurnal Ilmiah Farmasi <http://journal.uui.ac.id/index.php/JIF>

Reviewer's Attachments [Q Search](#)

66840	jif-review-assignment-21570-Other-65287.doc	May 10, 2022
-------	---	--------------

Revisions [Q Search](#) [Upload File](#)

68961	Breast Cancer Article.docx	June 19, 2022	Article Text
-------	----------------------------	---------------	--------------

Submit Revisi (19 Juni 2022)

Jurnal Ilmiah Farmasi

--- Back to Submissions

Notifications

- [\[JIF\] Editor Decision](#) 2022-06-18 02:39 AM
- [\[JIF\] Editor Decision](#) 2022-08-04 03:12 AM

Reviewer's Attachments

[66840](#) [jif-review-assignment-21570-Other-65287.doc](#) May 10, 2022

Revisions

[68961](#) [Breast Cancer Article.docx](#) June 19, 2022 Article Text

Review Discussions

Name	From	Last Reply	Replies	Closed
No Items				

Accept Submission (4 Agustus 2022)

Jurnal Ilmiah Farmasi

--- Back to Submissions

Notifications

[JIF] Editor Decision

2022-08-04 03:12 AM

Godeliva Adriani Hendra:

We have reached a decision regarding your submission to Jurnal Ilmiah Farmasi, "Evaluation of Single and Combination Chemotherapy Agents in Patients with Metastatic Breast Cancer".

Our decision is to: Accept Submission

Mohon dapat melakukan proofread manuskrip dan melampirkan :

1. Naskah yang telah direvisi sesuai proofread
2. Bukti proofread
3. Surat pernyataan proofread manuskrip dari lembaga terpercaya

Naskah final kami tunggu paling lambat 3 september 2022. Terima kasih

Jurnal Ilmiah Farmasi <http://journal.uii.ac.id/index.php/JIF>

Review Discussions

[Add discussion](#)

Proofread (25 Agustus 2022)

Jurnal Ilmiah Farmasi 🔔 👤

← Back to Submissions

21570 / Aditya et al. / Evaluation of single and combination chemotherapy agents in patients with metastatic breast cancer Library

Workflow **Publication**

Submission **Review** Copyediting **Production**

Copyediting Discussions Add discussion

Name	From	Last Reply	Replies	Closed
Copyediting	admin_ori 2022-03-18 09:09 PM	-	0	<input type="checkbox"/>
▶ Artikel, bukti proofread, dan surat pernyataan proofread	godeliva13 2022-08-25 04:41 AM	-	0	<input type="checkbox"/>
▶ Perihal Proofread Artikel	godeliva13 2022-09-21 03:04 AM	-	0	<input type="checkbox"/>
Proofread Author	dianmedisa 2022-12-12 01:10 AM	godeliva13 2022-12-30 04:52 AM	3	<input type="checkbox"/>

[https://journal.uii.ac.id/JIF/\\$\\$\\$call\\$\\$\\$/tab/author-dashboard/author-dashboard-tab/fetch-tab?submissionId=21570&stageId=4](https://journal.uii.ac.id/JIF/$$$call$$$/tab/author-dashboard/author-dashboard-tab/fetch-tab?submissionId=21570&stageId=4)

Final Proofread (30 Desember 2022)

Jurnal Ilmiah Farmasi 🔔 👤

← Back to Submissions

Copyediting Discussions Add discussion

Name	From	Last Reply	Replies	Closed
Copyediting	admin_ori 2022-03-18 09:09 PM	-	0	<input type="checkbox"/>
▶ Artikel, bukti proofread, dan surat pernyataan proofread	godeliva13 2022-08-25 04:41 AM	-	0	<input type="checkbox"/>
▶ Perihal Proofread Artikel	godeliva13 2022-09-21 03:04 AM	-	0	<input type="checkbox"/>
Proofread Author	dianmedisa 2022-12-12 01:10 AM	godeliva13 2022-12-30 04:52 AM	3	<input type="checkbox"/>

Copyedited Search

No Files

Evaluation of Single and Combination Chemotherapy Agents in Patients with Metastatic Breast Cancer

Abstract

Background: There are several ways to treat breast cancer, one of which is administering chemotherapy agents. Chemotherapy agents have activity in inhibiting the cell cycle. That will affect the effectiveness of therapy and the side effects of chemotherapy agents.

Objective: This study aimed to evaluate single and combination chemotherapy agent therapeutic efficacy and side effects

Method: The design of this study used an observational cohort study with retrospective data collection from January to December 2019. Patients obtained from medical records were diagnosed with Ca mammae metastatic stage at Panti Nirmala Hospital, Malang. The effectiveness of chemotherapy agent therapy was seen from the CEA and CA 15-3 (examinations were carried out in the first and third cycles) and analyzed using the Wilcoxon and U-Mann Whitney tests. Side effects of chemotherapy agents were analyzed descriptively.

Results: Analysis of the Wilcoxon test showed differences between the two groups of chemotherapeutic agents in CEA and CA 15-3 ($p < 0.05$). U-Mann Whitney test analysis showed no difference after administration of the two groups of chemotherapy agents at CEA ($p > 0.05$). However, there was a difference in CA 15-3 ($p < 0.05$). Both chemotherapy agents' most common side effects were pain, nausea/vomiting, and alopecia.

Conclusion: Patients who received combination chemotherapy agents had lower CA 15-3 levels than single chemotherapy agents.

Keywords: Ca Mammae Metastatic Stage, CEA, CA 15-3, Side Effects of Chemotherapy Agents

1. Introduction

Metastatic stage breast cancer is one of the highest malignancies among other types of cancer in Indonesia and globally, with an increasing incidence (Kementerian Kesehatan Republik Indonesia, 2019; Yang et al., 2017). As the implementation, metastatic breast cancer patients need to be monitored continuously to increase the survival rate. One way to assess the therapeutic effect is by measuring tumor biomarkers (McDonald, Clark, Tchou, Zhang, & Freedman, 2016; Yang et al., 2017). Reference tumor biomarkers are widely used to measure response to treatment, early recurrence, and predict prognosis. The biomarkers that are widely referenced to assess the effectiveness of therapy are carcinoembryonic antigen (CEA) which is a protein involved in cell adhesion, and cancer antigen (CA) 15-3 or MUC-1, which represents the mucin sequence in cells undergoing malignancy (Hosseini, Razmjoo, Arvandi, & Barat, 2015; Yang et al., 2017). CEA and CA 15-3 are less sensitive for early detection but can predict positive treatment responses (Yang et al., 2017). Serum levels of CEA and CA 15-3 are also widely used to expect a response to therapy in metastatic breast cancer (GENG, LIANG, YE, & ZHAO, 2015).

Commented [DL1]: Kalimat diparafrase untuk mengurangi ambiguitas (peletakan kata both di belakang)--- most common effects itu untuk kedua agen kemoterapi

Commented [DL2]: Dijadikan keterangan, karena pada kalimat sumber justru menjadikannya tidak lengkap.

In addition to the effectiveness of therapy, the side effects of chemotherapy agents also need to be monitored. Monitoring the side effects of chemotherapy agents aims to improve the patient's quality of life. Many reports have stated the side effects induced by chemotherapeutic agents, including spinal cord suppression, neuropathy, gastrointestinal disorders, hair loss, weakness, and skin disorders (Chan & Ismail, 2014). In Indonesia, there have not been many studies that have looked at the differences in biomarkers and the frequency of possible side effects that occur with single chemotherapy agents compared to combinations. This background underlies the need for a study to determine the differences in biomarkers of CEA and CA 15-3 and the side effects of using a single agent compared to a combination that may occur.

Commented [DL3]: Source: many reports have related to side effects induced by chemotherapeutic agents → kalimat tidak memberikan makna yang jelas karena jika have merupakan kata kerja (transitive verb-simple present tense), maka seharusnya diikuti oleh obyek langsung bukannya keterangan (to side effects)

2. Methods

The design of this study used an observational cohort study with retrospective data collection. Data was collected through medical records and laboratory results from January to December 2019. This research has passed ethical test No.E.5.a/209/KEPK-UMM/VIII/2020. The study location was at Panti Nirmala Hospital Malang and carried out in the Medical Record Unit and inpatient ward. The design of this study used an observational cohort study with retrospective data collection. Data were taken through medical records and laboratory results from January to December 2019. This research has passed ethical test No.E.5.a/209/KEPK-UMM/VIII/2020.

Commented [DL4]: Urutan penyampaianya diubah untuk memperjelas bahwa efek samping yang dimaksud adalah yang merupakan akibat dari penggunaan agen tunggal → maka keterangan mengenai agen tunggal diletakkan di awal.

2.1 Sampling and Sampling Techniques

The population of this study were patients diagnosed with breast cancer in the metastatic stage who were hospitalized at Panti Nirmala Hospital and received chemotherapy agents. The study sample was breast cancer patients with metastases who entered the inclusion criteria. The sampling technique used in this study was the total sampling technique, where all population members were sampled. There were 45 samples from the medical record unit, 19 patients received single chemotherapy agents and 26 received combination chemotherapy agents.

Commented [DL5]: Data collection disingkat menjadi data saja, karena sudah mewakili keseluruhan (data adalah bentuk jamak dari datum dalam Bahasa Inggris)

Commented [DL6]: The location of the study disederhanakan menjadi study location (kalimat dalam jurnal cenderung singkat, padat, dan jelas), dan untuk mengurangi pengulangan penggunaan struktur kalimat yang sama pada kalimat berikutnya

Commented [DL7]: Data collection disingkat menjadi data saja, karena sudah mewakili keseluruhan (data adalah bentuk jamak dari datum dalam Bahasa Inggris)

Commented [DL8]: Ada pengulangan kalimat "the location of the study at Panti Nirmala Hospital was carried out in the Medical Record Unit and inpatient ward", maka kalimat tersebut dihilangkan.

Commented [DL9]: The sample of the study disederhanakan menjadi study sample (kalimat dalam jurnal cenderung singkat, padat, dan jelas).

Commented [DL10]: There were 45 samples obtained disederhanakan menjadi there were 45 samples from (agar menjadi lebih ringkas)

2.2 Inclusion and Exclusion Criteria

The inclusion criteria included metastatic breast cancer patients who received single and combination chemotherapy agents; patient data was taken from the patient medical record viewed from biomarkers Carcinoembryonic Antigen (CEA) and Cancer Antigen (CA) 15-3. Exclusion criteria were patients who passed away and patients who did not have complete laboratory data when administering chemotherapy agents.

Commented [DL11]: Disederhanakan menjadi passed away agar kalimat tidak terlalu panjang

2.2 Data Collection

The data collected in this study were in the form of patient characteristics data which included: age, education, employment status, marital status, comorbidities, and history of hormonal contraceptive use. The data on the effectiveness of therapy was seen from the biomarkers of CEA and CA 15-3. Data on side effects were seen from using single and combination chemotherapy agents. The single chemotherapy agents used were Carboplatin, Navelbin, and Zometa/Zolenic. Combination chemotherapy agents consist of two chemotherapeutic agents and three chemotherapeutic agents. Two chemotherapeutic agents include Carboplatin+Docetaxel, Cisplatin+Docetaxel, and Cisplatin+Paclitaxel. The three chemotherapy agents have Carboplatin+Docetaxel+Fluorouracil, Cyclophosphamide+Epirubicin+Fluorouracil.

2.3 Data Analysis

The data of this study were analyzed using the open-source software R. The test that looked at the relationship between chemotherapy agents and the characteristics of the respondents used the Chi-Square test. A test that compares therapeutic efficacy as measured by biomarkers CEA and CA 15-3 uses the U-Mann Whitney test. Side effect data were analyzed descriptively as a percentage of events.

3 Result

Characteristics of metastatic breast cancer patients in the group receiving single or combination chemotherapy agents showed no relationship ($p>0.05$). Patient characteristics data are briefly presented in Table 1.

Table 1. Patient Characteristics Data

Characteristic	Single Chemotherapy* (n=19)	Combination Chemotherapy** (n=26)	p-value
Age (year)			
26-45	3 (15,8)	5 (19,23)	0,917
46-65	14 (73,68)	19 (73,07)	
66-85	2 (10,53)	2 (7,69)	
Mean 52 tahun			

Education			
Elementary	5(26,31)	9 (34,61)	0,844
Junior High School	0	1 (3,85)	
Senior High Scholl	6 (31,58)	7 (26,92)	
Associate Degree	3 (15,79)	3 (11,54)	
Bachelor Degree	2 (10,53)	6 (23,08)	
Master Degree	3 (15,79)	0	
Comorbidities			
Yes	6 (31,58)	6 (23,08)	0,767
No	13 (68,42)	20 (76,92)	
Occupation			
Yes	6 (31,58)	12 (46,15)	0,498
No	13 (68,42)	14 (53,85)	
Married Status			
Married	15 (78,95)	18 (69,23)	0,308
Single	0	3 (11,54)	
Divorce	4 (21,05)	5 (19,23)	
Hormonal History			
Contraception Hormonal	16 (84,21)	17 (65,38)	0,171
Contraception Non-Hormonal	1 (5,26)	7 (26,92)	
No	2 (10,53)	2 (7,69)	
Biomarker CEA			
Normal	1 (5)	1 (3,85)	1
Abnormal	18 (95)		
Biomarker CA 15-3			
Normal	1 (5)	1 (3,85)	1
Abnormal	18 (95)	25 (96,15)	

Information:

*Carboplatin, Navelbin, Zometa/Zolenic

**Combination of two therapy agents: carboplatin+docetaxel, cisplatin+docetaxel, cisplatin+paclitaxel; combination of three therapy agent: carboplatin+docetaxel+fluorouracil, cychlophosphamide+epirubicin+fluorouracil

Patients were dominated by 46-65 years, had no comorbidities, married status, and previously used hormonal family planning. Furthermore, the analysis of differences in CEA and CA15-3 biomarkers between those receiving single and combined chemotherapy agents can be seen in Table 2. There are differences in CA15-3 biomarkers after administration of single and combination chemotherapy agents. However, there was no difference in the CEA biomarker after administering single chemotherapy agents (mean CEA = 9.42 ng/ml) and combination chemotherapy agents (mean CEA = 9.70 ng/ml). CA15-3 levels of chemotherapeutic agents combinations had lower concentrations (mean CA15-3 = 105.60 U/ml) than single chemotherapy agents (mean CA15-3 = 129.48 U/ml).

Table 2. Different Tests of Single and Combination Chemotherapy Agents

Biomarker	W	p-value*
CEA	256	0.422
CA 15-3	325.5	0.036

The side effects data for single chemotherapy agent are represented in Table 3. The most common side effects are nausea/vomiting, alopecia, pain.

Table 3. Descriptive Data Side Effects of Single Chemotherapy Agents

Single Chemotherapy Agent (n=19 patients)	Side Effects	Number of Patients (%)
Carboplatin 400mg inj (n=1)	Nausea/vomiting and alopecia	1 (100)
Navelbin 40mg inj (n=15)	Pain	8 (53.33)
	Extravasation	1 (6.67)
	Nauseous/vomiting	12 (80)
	Short of breath	1 (6.67)
	Alopecia	12 (80)
Zometa/Zolenic 4mg (n=3)	Pain	2 (66.67)
	Nauseous/vomiting	3 (100)
	Alopecia	1 (33.33)

The side effects data for combination chemotherapy agents are represented in Table 4. The most common side effects are nausea/vomiting, alopecia, pain, extravasation, shortness of breath.

Table 4. Descriptive Data Side Effects of Combination Chemotherapy Agents

Combination Chemotherapy Agent (n=26 patients)	Side Effects	Number of Patients (%)
Carboplatin 300mg inj + docetaxel 110mg inj (n=3)	Pain	2 (66.67)
	Nausea/vomiting	3 (100)
	Alopecia	3 (100)
Carboplatin 300mg inj + docetaxel 120mg inj (n=1)	Pain, nausea/vomiting, and Alopecia	1 (100)
Carboplatin 400mg inj + docetaxel 120mg inj (n=1)	Extravasation, pain,nausea/vomiting, and alopecia	1 (100)
Carboplatin 450mg inj + docetaxel 110mg inj (n=2)	Pain, nausea/vomiting, Alopecia, and short of breath	2 (100)
Carboplatin 450mg inj + docetaxel 120mg inj (n=4)	Extravasation	1 (25)
	Pain	3 (75)
Cisplatin 50mg inj + docetaxel 100mg inj (n=4)	Extravasation	1 (36.36)
	Pain	2 (50)
	Nausea/vomiting	4 (100)
	Alopecia	1 (25)

Table 4. Descriptive Data Side Effects of Combination Chemotherapy Agents (Continued)

Combination Chemotherapy Agent (n=26 patients)	Side Effects	Number of Patients (%)
Carboplatin 60 mg inj + docetaxel 120 mg inj (n=3)	Pain	3 (100)
	Nausea/vomiting	3 (100)
	Alopecia	1 (50)

Cisplatin 50 mg inj + paclitaxel 200 mg inj (n=1)	Extravasation and nausea/vomiting	1 (100)
Cisplatin 60 mg inj + paclitaxel 230 mg inj (n=1)	Extravasation, nausea/vomiting and alopecia	1 (100)
Carboplatin 450mg inj, belotaxel 120 mg inj* + curacil 500 mg inj* (n=5)	Extravasation	1 (20)
	Pain	5 (100)
	Nausea/vomiting	5 (100)
	Short of breath	1 (20)
Cyclovid 600 mg inj*, epirubicin 500 mg inj + curacil 500 mg inj* (n=1)	alopecia	4 (80)
	Extravasation, pain, nausea/vomiting, and alopecia	1 (100)

3 Discussion

Administration of single and combined chemotherapeutic agents decreased CA 15-3 biomarkers, but not CEA. That showed a significant response to the effectiveness of therapy after chemotherapy. Parameters of response to chemotherapy agents evaluated through CEA's tumor markers in metastatic breast cancer had CEA levels that increased continuously. That explained that cancer cells did not respond to treatment/relapse after getting treatment. In addition, patients with metastatic breast cancer had high CEA levels. CEA is an insensitive biomarker in breast cancer, so it cannot be used as a screening. This condition caused the need for supporting data in the form of a CA15-3 biomarker. The CA15-3 biomarker aims to determine the prognosis of metastatic breast cancer. Monitoring response to therapy on CA15-3 was seen from increased levels of CA15-3, which was associated with the high severity of breast cancer (Kabel, 2017).

In a retrospective study, serum CA15-3 correlated with the location of the number of metastases in breast cancer patients but not with CEA. CEA has a higher sensitivity in patients with metastases that spread to other organs. So, it can be concluded that in patients with metastatic breast cancer, the sensitivity of the CA15-3 biomarker is intended for serum tumor markers in the breast. In contrast, the CEA biomarker is designed for sensitivity/marker of metastases in breast cancer patients (Yang et al., 2017).

The limitation of the problem in this study was that the sampling of CEA and CA 15-3 biomarker data was only in the first and third cycles.

The most frequent post-chemotherapy side effects using single and combination chemotherapy agents were nausea/vomiting. Nausea/ vomiting was caused by chemotherapy-induced nausea and vomiting (CINV). The risk of developing CINV was higher with the short-term IV infusion route than with long-term or oral preparations. Neurotransmitters that play a role in activating CINV are 5-hydroxytryptamine (5-HT₂, 5-HT₃, and 5-HT₄), dopamine (D₂), histamine (H₁), and acetylcholine (ACh). Receptors of the

Commented [DL12]: Kalimat sumber disederhanakan

Commented [DL13]: Naskah sumber: in the condition of patients with metastatic breast cancer----disederhanakan menjadi in patinets with metastatic breast cancer (disederhanakan)

Commented [DL14]: Naskah sumber: the most incidence kurang sesuai dengan konteks, maka diganti menjadi the most-frequent

neurotransmitter 5-HT and dopamine were abundant in the intestinal mucosa and activated when the neurotransmitter is released, causing nausea and vomiting. Chemotherapy agents given to patients will also quickly bind to the chemoreceptor trigger zone (CTZ) in the brain close to the vomiting center (vomiting center), resulting in nausea and vomiting (Antonarakis & Hain, 2004; Singh, Yoon, & Kuo, 2016).

The most common side effect was alopecia/chemotherapy-induced alopecia (CIA) (Rossi et al., 2017). The incidence of alopecia was associated with the type of chemotherapeutic agent used in this study. The finding showed that 80% of the incidence occurred in the antimicrotubule group (Vinorelbine, Paclitaxel, Docetaxel, Epirubicin), 60% in alkylators (carboplatin and cisplatin), and 10-50% in antimetabolites (fluorouracil), 60-100% on topoisomerase. Alopecia results from the main target of chemotherapeutic agents, the keratinocyte matrix, proliferates during the anagen/hair formation phase. The keratinocyte matrix is sensitive to chemotherapeutic agents and causes rapid apoptosis, which affects the anagen phase, namely hair follicle growth, and causes baldness (Haslam & Smart, 2019). However, this baldness is not permanent; where within six months, the hair growth will occur again (Rossi et al., 2017).

The subsequent most common side effect was a painful condition, often referred to as chemotherapy-induced peripheral neuropathy (CIPN). Peripheral neuropathy induced by platinum-based drugs (carboplatin and cisplatin) causes glial cell activation increasing the inflammatory pro-cytokines that increase nociceptor sensitivity and hyperexcitability of peripheral neurons. The effects, together with ROS (reactive oxygen species), damage the blood-brain barrier and leads to the development of neuroinflammation, damage to mitochondria, and increases ROS, which causes damage to enzymes, proteins, and lipids in neurons as well as dysregulation of calcium homeostasis which induces apoptotic changes in peripheral nerves. Peripheral neuropathy in using antimicrotubule (Vinorelbine, Paclitaxel, Docetaxel, Epirubicin) interferes with axonal transport. It causes degeneration of distal nerve segments, alters ion channel activity and hyperexcitability of peripheral neurons, and modifies the expression and function of Na⁺, K⁺ ion channels. Activation of microglia leads to the release and increase of pro-inflammatory cytokines. This process leads to nociceptor sensitivity and the development of neuroinflammation (Zajackowska et al., 2019).

4 Conclusion

The achievement in this study was that the CA15-3 biomarker in combination chemotherapy agents has lower levels than single chemotherapy agents. However, CEA

Commented [DL15]: Kalimat asal terlalu panjang, maka dibagi 2 agar tidak mengaburkan makna.

Commented [DL16]: Kalimat diparafrase agar lebih mudah dipahami

Commented [DL17]: Kalimat terlalu panjang, maka dijadikan 2 kalimat untuk mempermudah pemahaman.

Commented [DL18]: Ditambahkan conjunction "and"

levels did not give a significant change in single and combined agents. Both chemotherapy agents' most occurring side effects were nausea/vomiting, alopecia, and pain.

Acknowledgment

We would like to thank Panti Nirmala Hospital Malang for assisting the research administration process, Mrs. Suhul Raos, who helped complete the research data, and Mrs. Martanty Aditya, who assisted in preparing this article.

References

- Antonarakis, E. S., & Hain, R. D. W. (2004). Nausea and vomiting associated with cancer chemotherapy: Drug management in theory and in practice. *Archives of Disease in Childhood*, 89(9), 877–880. <https://doi.org/10.1136/adc.2003.037341>
- Chan, H. K., & Ismail, S. (2014). Side effects of chemotherapy among cancer patients in a Malaysian general hospital: Experiences, perceptions and informational needs from clinical pharmacists. *Asian Pacific Journal of Cancer Prevention*, 15(13), 5305–5309. <https://doi.org/10.7314/APJCP.2014.15.13.5305>
- GENG, B., LIANG, M.-M., YE, X.-B., & ZHAO, W.-Y. (2015). Association of CA 15-3 and CEA with clinicopathological parameters in patients with metastatic breast cancer. *Molecular and Clinical Oncology*, 3(1), 232–236. <https://doi.org/10.3892/mco.2014.419>
- Haslam, I. S., & Smart, E. (2019). Chemotherapy-Induced Hair Loss: The Use of Biomarkers for Predicting Alopecic Severity and Treatment Efficacy. *Biomarker Insights*, 14. <https://doi.org/10.1177/1177271919842180>
- Hosseini, S. M., Razmjoo, S., Arvandi, S. H., & Barat, T. (2015). CEA and CA 15-3 serum level in metastatic breast cancer and its correlation with distant metastasis. *Biomedical and Pharmacology Journal*, 8SE, 721–727. <https://doi.org/10.13005/bpj/775>
- Kabel, A. M. (2017). Tumor markers of breast cancer: New prospectives. *Journal of Oncological Sciences*, 3(1), 5–11. <https://doi.org/10.1016/j.jons.2017.01.001>
- Kementerian Kesehatan Republik Indonesia. (2019). Panduan Penatalaksanaan Kanker Payudara (Breast Cancer Treatment Guideline). *Jurnal Kesehatan Masyarakat*, 4(4), 1–50. Retrieved from <http://kanker.kemkes.go.id/guidelines/PPKPayudara.pdf>
- McDonald, E. S., Clark, A. S., Tchou, J., Zhang, P., & Freedman, G. M. (2016). Clinical diagnosis and management of breast cancer. *Journal of Nuclear Medicine*, 57, 9S–16S. <https://doi.org/10.2967/jnumed.115.157834>
- Rossi, A., Fortuna, M., Caro, G., Pranteda, G., Garelli, V., Pompili, U., & Carlesimo, M. (2017). Chemotherapy-induced alopecia management: clinical experience and practical advice. *J Cosmet Dermatol*, 16(4), 537–541. <https://doi.org/10.1111/jocd.12308>
- Singh, P., Yoon, S. S., & Kuo, B. (2016). Nausea: A review of pathophysiology and therapeutics.

Commented [DL19]: Auxiliary verb “had” dihilangkan karena penggunaan past perfect tense tidak sesuai untuk konteks kalimat. Maka diganti dengan menggunakan simple past tense.

Commented [DL20]: Auxiliary verb “had” dihilangkan karena penggunaan past perfect tense tidak sesuai untuk konteks kalimat. Maka diganti dengan menggunakan simple past tense.

Therapeutic Advances in Gastroenterology, 9(1), 98–112.
<https://doi.org/10.1177/1756283X15618131>

Yang, Y., Zhang, H., Zhang, M., Meng, Q., Cai, L., & Zhang, Q. (2017). Elevation of serum CEA and CA15-3 levels during antitumor therapy predicts poor therapeutic response in advanced breast cancer patients. *Oncology Letters*, 14(6), 7549–7556.
<https://doi.org/10.3892/ol.2017.7164>

Zajaczkowska, R., Kocot-Kępska, M., Leppert, W., Wrzosek, A., Mika, J., & Wordliczek, J. (2019). Mechanisms of chemotherapy-induced peripheral neuropathy. *International Journal of Molecular Sciences*, 20(6). <https://doi.org/10.3390/ijms20061451>

LEMBAGA PENGEMBANGAN BAHASA

LANGUAGE CENTER (LC-UMM)

lc.umm.ac.id | lc@umm.ac.id | lembagabahasa.umm@gmail.com

UMM
est. 1964

UNIVERSITAS
MUHAMMADIYAH
MALANG

Malang, August 12, 2022

Number : E.6./413/LC-UMM/VIII/2022
Attachment : -
Subject : Certification of Translation


To Whom It May Concern

Language Center of Universitas Muhammadiyah Malang certifies that the following document:

1. Title : Evaluation of Single and Combination Chemotherapy Agents in Patients with Metastatic Breast Cancer
2. Type : Scientific Journal of Pharmacy
3. Author : 1. Martanty Aditya
2. Godeliva Adriani Hendra
3. Suhul Raos Kumawula Ing Gustya
4. Institution : University of Ma Chung

was translated to the best of our ability and knowledge by the translator team of Language Center Universitas Muhammadiyah Malang.

Acknowledged by
behalf of the Director,



Dessy Listyarini
Dessy Listyarini
Coordinating Translator

Kampus I
Jl. Bandung 1 Malang, Jawa Timur
P: +02 341 551 253 (Hunting)
F: +02 341 480 435

Kampus II
Jl. Bendungan Sutami No. 188 Malang, Jawa Timur
P: +02 341 551 149 (Hunting)
F: +02 341 582 000

Kampus III
Jl. Raya Tlogomas No 240 Malang, Jawa Timur
P: +02 341 404 318 (Hunting)
F: +02 341 400 435
E: webmaster@umm.ac.id

