

Bukti Korespondensi ATC/DDD

Jurnal Nasional Terakreditasi Sinta 3

Judul artikel : Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients

Jurnal : Jurnal Farmasi Galenika Vol. 10, No. 1 (2024)

Penulis : Godeliva Adriani Hendra, Dhanang Prawira Nugraha, Tuswatul Anggi Krisdia

No.	Perihal	Tanggal
1.	Submit artikel	20 Oktober 2023
2.	Lampiran reviewer pertama tahap pertama	20 November 2023
3.	Lampiran reviewer kedua tahap pertama	27 November 2023
4.	Submit revisi artikel tahap pertama	27 Desember 2023
5.	Lampiran reviewer tahap kedua	5 Januari 2024
6.	Keputusan Editor " <i>Revisions Required</i> "	26 Januari 2024
7.	Submit revisi artikel tahap kedua	31 Januari 2024
8.	Lampiran reviewer tahap ketiga	9 Februari 2024
9.	Keputusan Editor " <i>Revisions Required</i> " dan Submit revisi artikel tahap ketiga	15 Februari 2024
10.	Keputusan Editor " <i>Revisions Required</i> " dan Submit revisi artikel tahap keempat	16 Februari 2024
11.	Keputusan Editor " <i>Revisions Required</i> "	20 Februari 2024
12.	Submit revisi artikel tahap kelima	21 Februari 2024
13.	<i>Copyediting</i>	25 Februari 2024
14.	<i>Copyright transfer agreement</i>	2 Maret 2024
15.	<i>Publishing</i>	3 April 2024

1. Submit artikel 20 Oktober 2023

Jurnal Farmasi Galenika (Galenika Journal of Pharma...) Tasks 0 English View Site godeliva13

Submission Library View Metadata

Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients
Godeliva Adriani Hendra, Dhanang Prawira Nugraha, Tuswatul Anggi Krisdia

Submission Review Copyediting Production

Submission Files Search

56770-1	godeliva13, Godeliva Adriani Hendra_Semnas Kefarmasian Untad 2023.docx	October 20, 2023	Article Text
56777-1	zubair, Evaluation_of_Anti_diabetic_Drugs_using_ATC_DDD_an, 13% plagiarism.pdf	October 20, 2023	Article Text
56779-1	zubair, Format review 16624-Article Text-56770-1-2-20231020.docx	October 20, 2023	Article Text

Download All Files

Pre-Review Discussions Add discussion

Name	From	Last Reply	Replies	Closed
No Items				

2. Lampiran reviewer pertama tahap pertama 20 November 2023

Jurnal Farmasi Galenika (Galenika Journal of Pharma...) Tasks 0 English View Site godeliva13

Reviewer's Attachments Search

57083-1	16624-Article Text-56939-1-4-20231020 review 1.docx	November 20, 2023
57216-1	16624-Article Text-56939-1-4-20231020.docx	November 27, 2023

Revisions Search Upload File

57681-1	Article Text, Revisi ATC-DDD Diabetes Mellitus.docx	December 27, 2023	Article Text
---------	---	-------------------	--------------

Review Discussions Add discussion

Name	From	Last Reply	Replies	Closed
No Items				

https://bestjournal.untad.ac.id/index.php/Galenika/\$\$call\$\$\$tab/author-dashboard/author-dashboard-review-round-tab/fetch-review-round-info?submissionId=16624&stageId=3&reviewRoundId=11843

Lampiran Reviewer Pertama Tahap Pertama

20 November 2023

Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients

ABSTRACT

Background: Diabetes Mellitus (DM), which results from insulin resistance, is closely linked to long-term damage to pancreatic beta cells, organ dysfunction, and organ failure, particularly in the eyes, kidneys, nerves, heart, and blood vessels. The development of the ATC/DDD system is necessary to increase global drug knowledge, ensure equitable drug availability, and promote responsible drug use. **Objectives:** This study aimed to identify data on patient characteristics and oral anti-diabetic drugs and insulin using the ATC/DDD and DU90% methods. **Material and Methods:** This study is a cross-sectional evaluation of oral anti-diabetic medications and insulin using the ATC/DDD method and DU90% at the Bala Keselamatan Bokor Turen Hospital from January to December 2022. The inclusion criteria were individuals diagnosed with type 1 or type 2 diabetes mellitus, with or without comorbidities, who were treated with oral anti-diabetic medications and a combination of insulin and oral medications. In the present study, the incomplete medical record data served as the exclusion criteria. The sample for this investigation consisted of 238 patients selected using the complete sampling technique. **Results:** The majority of patients who used oral anti-diabetic medications and insulin were between the ages of 46 and 65 (71.34%), were female (73.2%), had standard body mass indices (BMIs between 18.5 and 25), and had diagnoses of DM + HT (20.73%) and Type II DM (18.29%). Glulisin was the most commonly used anti-diabetic drug, accounting for 589 DDD/100 days of hospitalization, and DU90% was 21.39%. **Conclusion:** Glulisin is the most frequently prescribed anti-diabetic medication at Bokor Turen Hospital.

Keywords: Antidiabetic Oral and Insulin; Diabetes Mellitus; ATC/DDD; DU 90%

ABSTRAK

Latar Belakang: Resistensi insulin yang berkembang menjadi Diabetes Melitus (DM) erat hubungannya dengan terjadinya kerusakan sel beta pankreas dalam jangka panjang, disfungsi organ, dan kegagalan organ terutama pada mata, ginjal, saraf, jantung, dan pembuluh darah. Pengembangan sistem ATC/DDD diperlukan untuk meningkatkan pengetahuan tentang penggunaan obat di seluruh dunia, memastikan ketersediaan obat secara merata, dan mendorong penggunaan obat yang bijak. Tujuan penelitian ini adalah mengidentifikasi data karakteristik pasien serta obat anti-diabetes oral dan insulin menggunakan metode ATC/DDD dan DU90%. Bahan dan Metode: Rancangan penelitian ini menggunakan *cross-sectional study* yang mengevaluasi obat anti-diabetes oral dan insulin dengan metode ATC/DDD dan DU90% di RS Bala Keselamatan Bokor Turen bulan Januari hingga Desember 2022. Kriteria inklusi terdiri dari pasien DM tipe 1 dan tipe 2 dengan dan atau tanpa komorbid, menggunakan obat anti-diabetes oral dan insulin/kombinasi keduanya. Sedangkan, kriteria eksklusinya berupa data rekam medis tidak lengkap. Populasi penelitian ini sebanyak 238 pasien menggunakan teknik total sampling. Hasil: Data karakteristik pasien pengguna obat anti-diabetes oral dan insulin terbanyak berusia 46 hingga 65 tahun (71,34%); jenis kelamin perempuan (73,2%); Indeks Masa Tubuh normal sebesar 53,66% (IMT= 18,5-25); diagnose DM+HT (20,73%) dan DM Tipe II (18,29%). Penggunaan obat anti-diabetes tertinggi adalah obat glulisin sebesar 589 DDD/100 hari rawat inap dan DU90% sebesar 21,39%. Kesimpulan: Obat anti-diabetes yang paling banyak digunakan di RS Bokor Turen adalah glulisin.

Kata kunci: Obat Anti-diabetes Oral dan Insulin; Diabetes Melitus; DDD; DU 90%

INTRODUCTION

Diabetes Mellitus (DM) is a chronic metabolic disorder. The presence of risk factors, damage to insulin secretion and sensitivity leads to an increase in blood glucose and changes in fat and protein metabolism (DiPiro, 2020). The lack of insulin function contributes to the development of microvascular, macrovascular, and neuropathy as a chronic consequence of DM (Almasdy *et al.*, 2015).

Commented [AR1]: English to poor, please check proofreading and grammar

Commented [AR2]: not fit for purpose

The World Health Organization (WHO) in 2023 states that in 2014, 8.5% of adults aged 18 years and over have diabetes. In 2019, diabetes was the direct cause of 1.5 million deaths, and 48% of all deaths from diabetes occurred before the age of 70. Furthermore, 460,000 deaths from kidney disease are caused by diabetes, and elevated blood glucose causes about 20% of deaths from cardiovascular disease (WHO, 2023). DM is more common in low- and moderate-income countries and countries undergoing major economic and demographic transformations. DM is a significant global health problem and requires proper prevention and management measures. DM in Indonesia is currently a serious problem. Most DM sufferers are type 2 DM groups. Therefore, it is essential to evaluate the use of the drug as a basis for selection to ensure that the drug is used appropriately, safely, and efficiently (Pitasari, Andayani and Wijayanti, 2022).

There is a need to develop ATC/DDD systems to acquire knowledge about drug use worldwide to achieve equitable drug availability and prudent drug use, especially in developing countries. The primary purpose of the ATC/DDD system is to facilitate research on drug use and improve the overall quality of drug use (Tahar et al., 2020). A retrospective study using a *cross-sectional study*, which evaluated drug use patterns and costs associated with Type 2 DM in Saudi Arabia, showed that biguanide (metformin) was most widely prescribed as a monotherapy drug followed by a *fixed-dose combination*. The effectiveness of monotherapy drugs decreases with the duration of treatment; in these cases, combination drugs are prescribed. The most commonly prescribed combination drug is a biguanide with sulfonylurea/biguanide with thiazolidinedione, according to guidelines by the *American Diabetes Association* (ADA). The combination of sitagliptin and metformin is most widely preferred and widely prescribed in *fixed-dose combination therapy*, followed by vildagliptin and metformin (Ali et al., 2022). Combination drugs are used when a single pill cannot achieve the desired blood glucose level in diabetic patients (Okoro, Nmeka and Erah, 2018).

Evaluation of the use of anti-diabetic drugs with ATC/DDD and DU90% methods can provide insight into the dominant drug use patterns and the extent to which these drugs follow existing treatment recommendations and guidelines. Therefore, this study evaluated oral anti-diabetic drugs and insulin using ATC/DDD and DU90% in DM patients with and without comorbidities at the Bokor Turen Safety Army Hospital.

MATERIAL AND METHODS

Research Methods

The study design used a *cross-sectional study* with retrospective data collection through patient medical records. Quantitative evaluation of the use of anti-diabetic drugs using ATC / DDD and DU 90%

techniques. This research has gone through the health research ethics committee with no E.5.a/145/KEPKUMM/V/2023.

Population and Sample

This population is in the form of patients hospitalized with a diagnosis of DM and or without comorbidities at the Bokor Turen Salvation Army Hospital. The study sample included patients hospitalized with a diagnosis of DM and without comorbidities from January to December 2022 and who met the inclusion and exclusion criteria. Inclusion criteria include patients aged ≥ 17 years with a diagnosis of DM and without comorbidities. Exclusion criteria are patients whose medical record data is incomplete.

Sampling Techniques

The sampling technique is total sampling, where the number of samples is equal to a population of 164 patients and meets the criteria for inclusion and exclusion of the study.

Data Analysis

The calculation in evaluating the use of ATC / DDD method anti-diabetic drugs in the hospitalization of the Bokor Turen Salvation Army Hospital uses the formula:

$$\frac{DDD}{100} \text{ hari rawat inap} = \frac{\text{Jumlah antidiabetik (gram)}}{\text{Standar DDD WHO (gram)}} \times \frac{100}{LOS}$$

DU 90% is used to identify the amount of drug used as much as 90% of the total use of prescribed medicines and compare it with the amount of residual drug use (RI, 2017). The efficiency of drug use must be observed if the amount of drug use in 10% is more. The DU value of 90% is known after calculating DDD / 100 days of hospitalization per year. DU 90% is obtained by arranging the use of antibiotics from highest to lowest, then determining the cumulative percentage up to 90%.

RESULTS

a. Demographic Characteristics of Diabetes Mellitus Patients

The demographic characteristics of Diabetes Mellitus patients at the Bokor Turen Salvation Army Hospital from January to December 2022 were 164 patients in age, gender, BMI, diagnosis, comorbidities, drug names, administration intervals, and duration of administration. Judging from the most significant number, in the age characteristics of 117 patients aged 46-65, as many as 120 were female, and as many as 88 patients had a regular Body Mass Index (BMI) of 18.5-25 (Table 1).

Commented [AR3]: With english

Commented [AR4R3]: All table and picture with full english

119

Table 1. Data on the characteristics of diabetes mellitus patients

Karakteristik	Bulan (Tahun 2022)												Jumlah pasien (n=164)	%
	Jan 2022	Feb 2022	Mar 2022	Apr 2022	Mei 2022	Jun 2022	Jul 2022	Agust 2022	Sept 2022	Okt 2022	Nov 2022	Des 2022		
Usia														
17-45 thn	1	0	1	1	0	2	3	3	0	0	1	0	12	7,32
46 – 65 thn	11	6	13	12	7	12	13	8	8	14	8	5	117	71,34
66 – 95 thn	5	5	1	8	1	5	0	2	1	4	1	2	35	21,34
Mean ± SD	59,71 ± 9,18	63,54 ± 7,09	57,46 ± 8,46	62,28 ± 10,95	57,62 ± 6,47	57,57 ± 13,38	52,68 ± 7,88	58,46 ± 11,68	58,11 ± 4,93	61,38 ± 5,89	56,3 ± 6,97	60,29 ± 7,29	58,78 ± 2,52	
Total	17	11	15	21	8	19	16	13	9	18	10	7	164	100
Jenis kelamin														
Perempuan	12	9	11	14	5	16	13	8	7	10	10	5	120	73,2
Laki-laki	5	2	4	7	3	3	3	5	2	8	0	2	44	26,8
Total	17	11	15	21	8	19	16	13	9	18	10	7	164	100
IMT														
Kurus < 18,4	1	1	1	2	0	0	0	0	1	0	0	0	6	3,66
Normal 18,5 – 25	11	4	5	11	4	12	10	11	4	6	7	3	88	53,66
Gemuk > 25	5	6	9	8	4	7	6	2	4	12	3	4	70	42,68
Mean ± SD	23,07 ± 3,07	24,89 ± 4,93	25,94 ± 4,12	24,44 ± 4,94	24,85 ± 2,39	24 ± 4,16	24,15 ± 3,52	23,26 ± 2,33	25,48 ± 5,98	26,37 ± 3,92	23,71 ± 3,82	25,40 ± 3,06	24,63 ± 1,08	
Total	17	11	15	21	8	19	16	13	9	18	10	7	164	100

120

121

122

123

124

125

126

The description of DM and or without comorbidities at the Bokor Turen Salvation Army Hospital showed that the most patients with a diagnosis of DM and Hypertension (HT) were 34 patients (20.73%), followed by patients with a diagnosis of Type 2 DM as many as 28 patients (18.29%). At the same time, DM patients with other comorbidities have fewer than ten patients (Table 2).

Table 2. Description of Patients with Diabetes Mellitus and or Without Comorbidities

Diagnosis dengan Komorbid	Jumlah (n = 164)	%
DM (HT)	34	20,73
DM T2	28	18,29
DM (HT, CAD)	6	3,66
DM (Anemia)	5	3,05
DM (CAD)	5	3,05
DM (CKD)	5	3,05
DM (CVA)	5	3,05
DM (COPD)	5	3,05
DM (CVA, HT)	4	2,44
DM (Anemia, Thalasemia)	3	1,83
DM (Dispepsia)	3	1,83
DM (GERD, HT)	3	1,83
DM (HT, Anemia)	3	1,83
DM (Asma)	2	1,22
DM (DKD)	2	1,22
DM (DKD, HT)	2	1,22
DM (HT, CKD)	2	1,22
DM (HT, HF)	2	1,22
DM (HT, Vertigo)	2	1,22
DM (STEMI)	2	1,22
DM (Vertigo)	2	1,22
DM (GERD)	1	0,61
DM (Anemia, PAD)	1	0,61
DM (AKI)	1	0,61
DM (Anemia, CKD)	1	0,61
DM (Anemia, Dispepsia)	1	0,61
DM (Angina, Thalasemia)	1	0,61
DM (CAD, COPD)	1	0,61
DM (CAD, CVA, HT)	1	0,61
DM (CAD, HT)	1	0,61
DM (CAD, STEMI)	1	0,61
DM (CAD, TB)	1	0,61

127

Table 2. Description of Patients with Diabetes Mellitus and Without Comorbidities

(Continued)

Diagnosis dengan Komorbid	Jumlah (n = 164)	%
DM (CKD, HT, TB)	1	0,61
DM (CKD, STEMI)	1	0,61
DM (CKD, Thalassemia)	1	0,61
DM (COPD, Dispepsia)	1	0,61
DM (COPD, HT)	1	0,61
DM (CVA, susp covid)	1	0,61
DM (Gastritis)	1	0,61
DM (GERD, HF)	1	0,61
DM (Hepatitis)	1	0,61
DM (HF)	1	0,61
DM (HF, CAD, COPD)	1	0,61
DM (HF, Hepatitis)	1	0,61
DM (HT, HF, dispepsia)	1	0,61
DM (HT, CA infark)	1	0,61
DM (HT, CAD, Gerd)	1	0,61
DM (HT, CAD, Parkinson)	1	0,61
DM (HT, COPD, CVA)	1	0,61
DM (HT, COPD, Dispepsia)	1	0,61
DM (HT, Dispepsia)	1	0,61
DM (HT, HF, CAD)	1	0,61
DM (HT, STEMI)	1	0,61
DM (Myalgia)	1	0,61
DM (PAD)	1	0,61
DM (PAD, Vertigo)	1	0,61
DM T1 (HT)	1	0,61
Total	164	100

Commented [AR5]: Add abbreviations

Based on the DM treatment profile seen in the 12 months of 2022, it shows that the use of anti-diabetic drugs is the most in the use of glargine drugs with a dose strength of 100U/ml in as many as 71 patients. Followed by glulisine drugs with a dose strength of 100U / ml for as many as 57 patients. The third highest use was aspart 100U / ml, as many as 54 patients. Concerning the administration interval of most anti-diabetic drugs given every 24 hours a day with the duration of most drug administration for 4-6 days (Table 3).

Table 3. Profile of Anti-diabetic Drug Use

Karakteristik	Bulan (Tahun 2022)												Total (n=293)	%
	Jan 2022	Feb 2022	Mar 2022	Apr 2022	Mei 2022	Jun 2022	Jul 2022	Agust 2022	Sept 2022	Okt 2022	Nov 2022	Des 2022		
Nama obat														
Acarbos 50 mg	1	0	0	0	0	1	2	0	0	0	0	1	5	1,7
Glulisin 100U/ml	4	7	8	10	5	7	2	3	1	5	3	2	57	19,5
Glargine 100U/ml	8	9	8	8	4	9	4	4	5	7	3	2	71	24,2
Glibenklamid 5 mg	1	0	0	0	0	0	1	1	0	0	0	0	3	1,0
Glikuidon 30 mg	2	0	4	1	0	1	1	1	1	4	2	1	18	6,1
Glimepirid 2 mg	2	0	0	3	2	3	3	3	2	0	0	0		
Glimepirid 4 mg	0	0	0	1	0	1	1	0	2	0	0	0	18	6,1
Glimepirid 3 mg	0	0	0	0	1	0	0	0	0	0	0	0		
Lispro 100U/ml	1	0	0	0	0	1	2	0	1	0	0	2	7	2,4
Detemir 100U/ml	2	0	3	6	3	5	4	2	2	7	2	2	38	13,0
Metformin 500 mg	2	1	0	4	1	2	3	3	1	1	1	1	20	6,8
Aspart 100U/ml	6	3	3	4	2	8	7	4	5	8	4	0	54	18,4
Pioglitazon 30 mg	0	0	0	0	0	0	0	1	0	0	0	0	1	0,3
Degludec-Aspart 100U/ml	0	1	0	0	0	0	0	0	0	0	0	0	1	0,3
Total	29	21	26	36	17	37	29	22	18	32	15	11	293	100
Interval pemberian														
24 jam	15	10	16	20	9	18	14	11	9	20	9	6	157	53,6
12 jam	2	1	0	3	1	3	3	2	3	2	1	0	21	7,2
8 jam	12	10	10	13	7	16	12	9	6	10	4	5	114	38,9
6 jam	0	0	0	0	0	0	0	0	0	0	1	0	1	0,3
Total	29	21	26	36	17	37	29	22	18	32	15	11	293	100

Karakteristik	Bulan (Tahun 2022)												Total (n=293)	%
	Jan 2022	Feb 2022	Mar 2022	Apr 2022	Mei 2022	Jun 2022	Jul 2022	Agust 2022	Sept 2022	Okt 2022	Nov 2022	Des 2022		
Durasi														
1-3 Hari	7	15	12	10	5	20	14	13	9	15	4	3	127	43,3
4-6 Hari	22	6	14	26	12	17	15	9	9	17	11	8	166	56,7
Total	29	21	26	36	17	37	29	22	18	32	15	11	293	100

b. Evaluation of the Use of Anti-diabetic Drugs with ATC/DDD and DU 90% Methods

The results of the evaluation of the use of anti-diabetic drugs using the ATC / DDD method showed that the use of antidiabetics that are often used is glulisine drugs with a dose strength of 100U / ml of 589.02 DDD / 100 days of hospitalization, which means that during 100 days of treatment in the hospital around 589 diabetic patients received glulisine amounting to 40 IU every day. The total number of patients hospitalized and using anti-diabetic drugs from January to December 2022 was 568 days. The second most significant use of anti-diabetic medications is aspart with a dose strength of 100U / ml of 519.72 DDD / 100 days of hospitalization, which means that during 100 days of hospitalization in the hospital around 520 DM patients received aspart drugs of 40 IU (Table 4).

162

Table 4. Analysis of the Use of Anti-diabetic Drugs using the ATC / DDD Method

No.	Kode ATC	Nama Obat	DDD (WHO)	LOS (Hari)	DDD/100 Hari rawat inap
1	A10AB06	Glulisin 100 U/ml	40 UI	568 Hari	589,02
2	A10AB05	Aspart 100 U/ml	40 UI		519,72
3	A10BB12	Glimepirid 2 mg, 3 mg, 4 mg	2 mg		474,28
4	A10AE04	Glargine 100 U/ml	40 UI		426,85
5	A10AE05	Detemir 100 U/ml	40 UI		221,38
6	A10BA02	Metformin 500 mg	2 g		198,80
7	A10BB08	Glikuidon 30 mg	60 mg		129,22
8	A10AB04	Lispro 100 U/ml	40 UI		85,20
9	A10BF01	Acarbos 50 mg	0,3 g		43,55
10	A10BB01	Glibenklamid 5 mg	10 mg		31,24
11	A10BG03	Pioglitazon 30 mg	30 mg		22,72
12	A10AD06	Degludec-Aspart 100 U/ml	40 UI		11,36

163

164 The 90% DU in this study was used to evaluate the use of anti-diabetic drugs in the top 90% of services
 165 in the population. There were 12 anti-diabetic assessed drugs in 90% DU and showed that glulycin,
 166 aspart, glimepiride 2 mg, glimepiride 3 mg, glimepiride 4 mg, glargine 100U/ml, detemir 100U/ml, and
 167 metformin 500 mg were anti-diabetic drugs that accounted for 90% of the highest anti-diabetic drug use
 168 in the DM patient population at the Bokor Turen Salvation Army Hospital (Table 5).

169

170

Table 5. Profile of Anti-diabetic Drug Use with DU Method 90%

Kode ATC	Nama Obat	DDD/100 Hari rawat inap	Persentase	Persentase Kumulatif	Segmen
A10AB06	Glulisin 100 U/ml	589,02	21,39%	21,39%	Du 90%
A10AB05	Aspart 100 U/ml	519,72	18,88%	40,27%	

171

172

173

174

Table 5. Profile of Anti-diabetic Drug Use with DU Method 90% (Continued)

	Glimepirid 2 mg			
A10BB12	Glimepirid 3 mg	474,28	17,23%	57,49%
	Glimepirid 4 mg			
A10AE04	Glargine 100 U/ml	426,85	15,50%	73,00%
A10AE05	Detemir 100 U/ml	221,38	8,04%	81,04%
A10BA02	Metformin 500 mg	198,80	7,22%	88,26%
A10BB08	Glikuidon 30 mg	129,22	4,69%	92,95%
A10AB04	Lispro 100 U/ ml	85,20	3,09%	96,05%
A10BF01	Acarbos 50 mg	43,55	1,58%	97,63%
A10BB01	Glibenklamid 5 mg	31,24	1,13%	98,76%
A10BG03	Pioglitazon 30 mg	22,72	0,83%	99,59%
A10AD06	Degludec- Aspart 100 U/ ml	11,36	0,41%	100,00%
		2753,33	99,99	

175

176

177 DISCUSSION

178 Insulin glulisine (Apidra) in this study occurred in patients with Type 2 diabetes with comorbidities,
 179 such as hypertension, coronary artery disease (CAD), and chronic kidney disease (CKD). On average,
 180 DM patients who get apidra drugs are given every 8 hours, and this drug is widely presented to 76% of
 181 female patients with an age range of 58-61 years. This insulin is widely recommended at the Bokor
 182 Turen Salvation Army Hospital because it can lower blood sugar levels quickly and has a more
 183 negligible risk of hypoglycemia. Following the 2021 PERKENI guidelines, Apidra is insulin rapid-
 184 acting, generally used with food. Apidra is designed to decrease glucose levels in the blood after a meal
 185 rapidly or when blood glucose levels are high. Apidra is commonly combined with basal insulins such
 186 as Lantus (insulin glargine) and Levemir (insulin detemir) (Soelistijo et al., 2021). A study examining
 187 the clinical effects of Type 2 DM patients with cardiovascular comorbidities where patients used insulin
 188 rapid-acting showed that it could have beneficial effects from insulin glulisine administration associated
 189 with death and stroke. Still, there was no difference in coronary heart disease (CHD) or cardiovascular
 190 disease (CVD) (Svensson et al., 2017).

191

Insulin aspart (novorapid) is given to patients with type 2 diabetes with comorbid HT accompanied by CAD, ST-Elevation Myocardial Infarction (STEMI), and Heart Failure (HF). The interval of insulin administration averaged every 8 hours per day and was used by 66% of female patients with an average of 58-61 years. Like glulisine insulin, this insulin can also quickly lower blood sugar levels and has a low risk of hypoglycemia. This insulin is also rapid-acting, which provides therapeutic effectiveness after 15 minutes, with the peak of therapeutic efficacy occurring within 1-2 hours and can last up to 4-6 hours. In the case of the population in Japan, administering insulin aspart to type 2 DM patients can significantly reduce cardiovascular complications within 5 to 10 years, resulting in improved quality of life and lower costs compared to *human insulin* (Pollock et al., 2011).

The third most significant use of anti-diabetic drugs is glimepiride doses of 2mg, 3 mg, and 4 mg. This drug is given to patients with type 2 diabetes with cardiovascular comorbidities and an interval of 24 hours per day and primarily female patients (83%) with an average age of 58 years. Regular glimepiride is combined with insulin or another oral medication such as metformin. In the case study of type 2 DM patients with CVD, the average patient suffering from DM was around 5.7 ± 4.8 years. CVD suffered by type 2 DM patients in the form of hypertension (68.5% of patients); dyslipidemia (47.9% of patients); CAD (25.4% of patients); Transient Ischemic Attack (TIA) in 3.6% of patients; peripheral artery disease (PAD) accounted for 4.8% of patients and heart failure in 2.9% of patients. Type 2 DM patients with various comorbidities receive the drug glimepiride/metformin Fixed Dose Combination (FDC) as a first-line therapy. As many as 68.2% of FDC patients achieved blood pressure within optimal limits. Most of the other patients experienced an increase in glycemic parameters and a change in body weight of about 18.4%. 59.2% of patients experienced weight loss (Ray et al., 2022). Glimepiride is a sulfonylurea class drug that has pharmacological effects to increase insulin production by pancreatic beta cells. The most common side effects are hypoglycemia and weight gain.

The limitation of this study is that data were taken only retrospectively from patient medical record data and did not make direct observations on DM patients.

CONCLUSION

Based on the analysis of demographic data and treatment profiles of DM patients, DM patients are dominated by the age group of 46-65 years, with women who use anti-diabetic drugs more. Patients with comorbidities use more anti-diabetic medications than patients with non-comorbid DM. Evaluation of anti-diabetic drugs ATC / DDD method shows that insulin glulisine (Apidra) is most used in DM patients

with and without comorbidities. Through the DU method, 90% of glulisine, aspart, glargine, glimepiride, detemir, and metformin drugs were most widely used in DM patients.

CONFLICT OF INTEREST

All authors declare no conflict of interest.

REFERENCES

- Ali, M.D. et al. (2022) 'Evaluation of drug utilization pattern and cost associated with diabetes mellitus Type two management in Saudi Arabia', *Brazilian Journal of Pharmaceutical Sciences*, 58, p. E20681. Available at: <https://doi.org/10.1590/s2175-97902022e20681>.
- Almasdy, D. et al. (2015) 'Evaluation of the Use of Anti-diabetic Drugs in Type-2 Diabetes Mellitus Patients in a Government Hospital of Padang City – West Sumatra', *Journal of Pharmaceutical & Clinical Science*, 2(1), p. 104. Available at: <https://doi.org/10.29208/jsfk.2015.2.1.58>.
- DiPiro, J.T. (ed.) (2020) *Pharmacotherapy: a pathophysiologic approach*. Eleventh edition. New York: McGraw Hill Medical.
- Okoro, R.N., Nmeka, C. and Erah, P.O. (2018) 'Utilization study of antidiabetes medicines at a tertiary care hospital in Nigeria', *Future Journal of Pharmaceutical Sciences*, 4(2), pp. 109–115. Available at: <https://doi.org/10.1016/j.fjps.2017.11.004>.
- Pitasari, N.W.N., Andayani, T.M. and Wijayanti, T. (2022) 'Evaluation of the Use of Antidiabetic Drugs in Patients of Back-Referral Program at the Demak District Pharmacy', *Journal of Management and Pharmacy Practice*, 12(2), p. 125. Available at: <https://doi.org/10.22146/jmpf.73841>.
- Pollock, R.F. et al. (2011) 'The cost effectiveness of rapid-acting insulin aspart compared with human insulin in type 2 diabetes patients: an analysis from the Japanese third-party payer perspective', *Journal of Medical Economics*, 14(1), pp. 36–46. Available at: <https://doi.org/10.3111/13696998.2010.541045>.
- Ray, S. et al. (2022) 'Usage Pattern of Glimepiride/Metformin Fixed-dose Combination in Type 2 Diabetes Patients with CVD or at Risk of CVD: An Experience in Indian Setting', *Asian Journal of Diabetology*, 23(2), pp. 13–19.
- RI, K. (2017) 'Technical Guidelines for Evaluation of Drug Use in Health Facilities'.
- Soelistijo, S.A. dkk. (2021) 'Guidelines for the management and prevention of adult type 2 diabetes mellitus in INDONESIA - 2021', *PB. PERKENI*, p. 119.
- Svensson, A.-M. et al. (2017) 'Clinical effects, cardiovascular and renal outcomes associated with rapid-acting insulin analogs among individuals with type 2 diabetes: a nation-wide observational cohort study', *Clinical Diabetes and Endocrinology*, 3(1), p. 5. Available at: <https://doi.org/10.1186/s40842-017-0043-2>.

Commented [AR6]: All English and all authors in references must be written in full

- 260 Tahar, N.dkk. (2020) 'EVALUATION OF THE USE OF ORAL ANTI-DIABETIC DRUGS IN TYPE
261 2 DIABETES MELLITUS PATIENTS USING ATC/DDD AND DU 90% METHODS', *The*
262 *2nd Alauddin Pharmaceutical Conference and Expo (ALPHA-C) 2020* [Preprint]. Available at:
263 <https://doi.org/10.24252/kesehatan.v1i1.18380>.
- 264 WHO (2023) *Diabetes*. Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes>
265 (Accessed: September 21, 2023).

Lampiran Reviewer Kedua Tahap Pertama

27 November 2023

Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients

ABSTRACT

Background: Diabetes Mellitus (DM), which results from insulin resistance, is closely linked to long-term damage to pancreatic beta cells, organ dysfunction, and organ failure, particularly in the eyes, kidneys, nerves, heart, and blood vessels. The development of the ATC/DDD system is necessary to increase global drug knowledge, ensure equitable drug availability, and promote responsible drug use. **Objectives:** This study aimed to identify data on patient characteristics and oral anti-diabetic drugs and insulin using the ATC/DDD and DU90% methods. **Material and Methods:** This study is a cross-sectional evaluation of oral anti-diabetic medications and insulin using the ATC/DDD method and DU90% at the Bala Keselamatan Bokor Turen Hospital from January to December 2022. The inclusion criteria were individuals diagnosed with type 1 or type 2 diabetes mellitus, with or without comorbidities, who were treated with oral anti-diabetic medications and a combination of insulin and oral medications. In the present study, the incomplete medical record data served as the exclusion criteria. The sample for this investigation consisted of 238 patients selected using the complete sampling technique. **Results:** The majority of patients who used oral anti-diabetic medications and insulin were between the ages of 46 and 65 (71.34%), were female (73.2%), had standard body mass indices (BMIs between 18.5 and 25), and had diagnoses of DM + HT (20.73%) and Type II DM (18.29%). Glulisin was the most commonly used anti-diabetic drug, accounting for 589 DDD/100 days of hospitalization, and DU90% was 21.39%. **Conclusion:** Glulisin is the most frequently prescribed anti-diabetic medication at Bokor Turen Hospital.

Keywords: Antidiabetic Oral and Insulin; Diabetes Mellitus; ATC/DDD; DU 90%

ABSTRAK

Latar Belakang: Resistensi insulin yang berkembang menjadi Diabetes Melitus (DM) erat hubungannya dengan terjadinya kerusakan sel beta pankreas dalam jangka panjang, disfungsi organ, dan kegagalan organ terutama pada mata, ginjal, saraf, jantung, dan pembuluh darah. Pengembangan sistem ATC/DDD diperlukan untuk meningkatkan pengetahuan tentang penggunaan obat di seluruh dunia, memastikan ketersediaan obat secara merata, dan mendorong penggunaan obat yang bijak. Tujuan penelitian ini adalah mengidentifikasi data karakteristik pasien serta obat anti-diabetes oral dan insulin menggunakan metode ATC/DDD dan DU90%. Bahan dan Metode: Rancangan penelitian ini menggunakan *cross-sectional study* yang mengevaluasi obat anti-diabetes oral dan insulin dengan metode ATC/DDD dan DU90% di RS Bala Keselamatan Bokor Turen bulan Januari hingga Desember 2022. Kriteria inklusi terdiri dari pasien DM tipe 1 dan tipe 2 dengan dan atau tanpa komorbid, menggunakan obat anti-diabetes oral dan insulin/kombinasi keduanya. Sedangkan, kriteria eksklusinya berupa data rekam medis tidak lengkap. Populasi penelitian ini sebanyak 238 pasien menggunakan teknik total sampling. Hasil: Data karakteristik pasien pengguna obat anti-diabetes oral dan insulin terbanyak berusia 46 hingga 65 tahun (71,34%); jenis kelamin perempuan (73,2%); Indeks Masa Tubuh normal sebesar 53,66% (IMT= 18,5-25); diagnose DM+HT (20,73%) dan DM Tipe II (18,29%). Penggunaan obat anti-diabetes tertinggi adalah obat glulisin sebesar 589 DDD/100 hari rawat inap dan DU90% sebesar 21,39%. Kesimpulan: Obat anti-diabetes yang paling banyak digunakan di RS Bokor Turen adalah glulisin.

Kata kunci: Obat Anti-diabetes Oral dan Insulin; Diabetes Melitus; DDD; DU 90%

INTRODUCTION

Diabetes Mellitus (DM) is a chronic metabolic disorder. The presence of risk factors, damage to insulin secretion and sensitivity leads to an increase in blood glucose and changes in fat and protein metabolism (DiPiro, 2020). The lack of insulin function contributes to the development of microvascular, macrovascular, and neuropathy as a chronic consequence of DM (Almasdy *et al.*, 2015).

Commented [MOU1]: Past tenses

Commented [MOU2]: Index?

The World Health Organization (WHO) in 2023 states that in 2014, 8.5% of adults aged 18 years and over have diabetes. In 2019, diabetes was the direct cause of 1.5 million deaths, and 48% of all deaths from diabetes occurred before the age of 70. Furthermore, 460,000 deaths from kidney disease are caused by diabetes, and elevated blood glucose causes about 20% of deaths from cardiovascular disease (WHO, 2023). DM is more common in low- and moderate-income countries and countries undergoing major economic and demographic transformations. DM is a significant global health problem and requires proper prevention and management measures. DM in Indonesia is currently a serious problem. Most DM sufferers are type 2 DM groups. Therefore, it is essential to evaluate the use of the drug as a basis for selection to ensure that the drug is used appropriately, safely, and efficiently (Pitasari, Andayani and Wijayanti, 2022).

There is a need to develop ATC/DDD systems to acquire knowledge about drug use worldwide to achieve equitable drug availability and prudent drug use, especially in developing countries. The primary purpose of the ATC/DDD system is to facilitate research on drug use and improve the overall quality of drug use (Tahar et al., 2020). A retrospective study using a *cross-sectional study*, which evaluated drug use patterns and costs associated with Type 2 DM in Saudi Arabia, showed that biguanide (metformin) was most widely prescribed as a monotherapy drug followed by a *fixed-dose combination*. The effectiveness of monotherapy drugs decreases with the duration of treatment; in these cases, combination drugs are prescribed. The most commonly prescribed combination drug is a biguanide with sulfonylurea/biguanide with thiazolidinedione, according to guidelines by the *American Diabetes Association* (ADA). The combination of sitagliptin and metformin is most widely preferred and widely prescribed in *fixed-dose combination therapy*, followed by vildagliptin and metformin (Ali et al., 2022). Combination drugs are used when a single pill cannot achieve the desired blood glucose level in diabetic patients (Okoro, Nmeko and Erah, 2018).

Evaluation of the use of anti-diabetic drugs with ATC/DDD and DU90% methods can provide insight into the dominant drug use patterns and the extent to which these drugs follow existing treatment recommendations and guidelines. Therefore, this study evaluated oral anti-diabetic drugs and insulin using ATC/DDD and DU90% in DM patients with and without comorbidities at the Bokor Turen Safety Army Hospital.

MATERIAL AND METHODS

Research Methods

The study design used a *cross-sectional study* with retrospective data collection through patient medical records. Quantitative evaluation of the use of anti-diabetic drugs using ATC / DDD and DU 90%

Commented [MOU3]: Please explain about the hospital, namely its level of accreditation, etc

techniques. This research has gone through the health research ethics committee with no E.5.a/145/KEPKUMM/V/2023.

Population and Sample

This population is in the form of patients hospitalized with a diagnosis of DM and or without comorbidities at the Bokor Turen Salvation Army Hospital. The study sample included patients hospitalized with a diagnosis of DM and without comorbidities from January to December 2022 and who met the inclusion and exclusion criteria. Inclusion criteria include patients aged ≥ 17 years with a diagnosis of DM and without comorbidities. Exclusion criteria are patients whose medical record data is incomplete.

Sampling Techniques

The sampling technique is total sampling, where the number of samples is equal to a population of 164 patients and meets the criteria for inclusion and exclusion of the study.

Data Analysis

The calculation in evaluating the use of ATC / DDD method anti-diabetic drugs in the hospitalization of the Bokor Turen Salvation Army Hospital uses the formula:

$$\frac{DDD}{100} \text{ hari rawat inap} = \frac{\text{Jumlah antidiabetik (gram)}}{\text{Standar DDD WHO (gram)}} \times \frac{100}{LOS}$$

DU 90% is used to identify the amount of drug used as much as 90% of the total use of prescribed medicines and compare it with the amount of residual drug use (RI, 2017). The efficiency of drug use must be observed if the amount of drug use in 10% is more. The DU value of 90% is known after calculating DDD / 100 days of hospitalization per year. DU 90% is obtained by arranging the use of antibiotics from highest to lowest, then determining the cumulative percentage up to 90%.

RESULTS

a. Demographic Characteristics of Diabetes Mellitus Patients

The demographic characteristics of Diabetes Mellitus patients at the Bokor Turen Salvation Army Hospital from January to December 2022 were 164 patients in age, gender, BMI, diagnosis, comorbidities, drug names, administration intervals, and duration of administration. Judging from the most significant number, in the age characteristics of 117 patients aged 46-65, as many as 120 were female, and as many as 88 patients had a regular Body Mass Index (BMI) of 18.5-25 (Table 1).

Commented [MOU4]: Past tenses

Commented [MOU5]:

Commented [MOU6]:

Commented [MOU7]: Do not use a picture

Commented [MOU8]: ???

Commented [MOU9]: Is this the number population in 1 year that meet the criteria? What is the total population in 1 year?is it only 164?

119

Table 1. Data on the characteristics of diabetes mellitus patients

Karakteristik	Bulan (Tahun 2022)												Jumlah pasien (n=164)	%
	Jan 2022	Feb 2022	Mar 2022	Apr 2022	Mei 2022	Jun 2022	Jul 2022	Agust 2022	Sept 2022	Okt 2022	Nov 2022	Des 2022		
Usia														
17-45 thn	1	0	1	1	0	2	3	3	0	0	1	0	12	7,32
46 – 65 thn	11	6	13	12	7	12	13	8	8	14	8	5	117	71,34
66 – 95 thn	5	5	1	8	1	5	0	2	1	4	1	2	35	21,34
Mean ± SD	59,71 ± 9,18	63,54 ± 7,09	57,46 ± 8,46	62,28 ± 10,95	57,62 ± 6,47	57,57 ± 13,38	52,68 ± 7,88	58,46 ± 11,68	58,11 ± 4,93	61,38 ± 5,89	56,3 ± 6,97	60,29 ± 7,29	58,78 ± 2,52	
Total	17	11	15	21	8	19	16	13	9	18	10	7	164	100
Jenis kelamin														
Perempuan	12	9	11	14	5	16	13	8	7	10	10	5	120	73,2
Laki-laki	5	2	4	7	3	3	3	5	2	8	0	2	44	26,8
Total	17	11	15	21	8	19	16	13	9	18	10	7	164	100
IMT														
Kurus < 18,4	1	1	1	2	0	0	0	0	1	0	0	0	6	3,66
Normal 18,5 – 25	11	4	5	11	4	12	10	11	4	6	7	3	88	53,66
Gemuk > 25	5	6	9	8	4	7	6	2	4	12	3	4	70	42,68
Mean ± SD	23,07 ± 3,07	24,89 ± 4,93	25,94 ± 4,12	24,44 ± 4,94	24,85 ± 2,39	24 ± 4,16	24,15 ± 3,52	23,26 ± 2,33	25,48 ± 5,98	26,37 ± 3,92	23,71 ± 3,82	25,40 ± 3,06	24,63 ± 1,08	
Total	17	11	15	21	8	19	16	13	9	18	10	7	164	100

120

121

122

123

124

125

126

Table 2. Description of Patients with Diabetes Mellitus and or Without Comorbidities

Diagnosis dengan Komorbid	Jumlah (n = 164)	%
DM (HT)	34	20,73
DM T2	28	18,29
DM (HT, CAD)	6	3,66
DM (Anemia)	5	3,05
DM (CAD)	5	3,05
DM (CKD)	5	3,05
DM (CVA)	5	3,05
DM (COPD)	5	3,05
DM (CVA, HT)	4	2,44
DM (Anemia, Thalasemia)	3	1,83
DM (Dispepsia)	3	1,83
DM (GERD, HT)	3	1,83
DM (HT, Anemia)	3	1,83
DM (Asma)	2	1,22
DM (DKD)	2	1,22
DM (DKD, HT)	2	1,22
DM (HT, CKD)	2	1,22
DM (HT, HF)	2	1,22
DM (HT, Vertigo)	2	1,22
DM (STEMI)	2	1,22
DM (Vertigo)	2	1,22
DM (GERD)	1	0,61
DM (Anemia, PAD)	1	0,61
DM (AKI)	1	0,61
DM (Anemia, CKD)	1	0,61
DM (Anemia, Dispepsia)	1	0,61
DM (Angina, Thalasemia)	1	0,61
DM (CAD, COPD)	1	0,61
DM (CAD, CVA, HT)	1	0,61
DM (CAD, HT)	1	0,61
DM (CAD, STEMI)	1	0,61
DM (CAD, TB)	1	0,61

127

Commented [MOU10]: Use the table format, not a images

Please present in English and the numbers of patient in 1 years, not every month

Commented [MOU11]: Use the table format, not a images

Please present in English

Table 2. Description of Patients with Diabetes Mellitus and Without Comorbidities

(Continued)

Diagnosis dengan Komorbid	Jumlah (n = 164)	%
DM (CKD, HT, TB)	1	0,61
DM (CKD, STEMI)	1	0,61
DM (CKD, Thalassemia)	1	0,61
DM (COPD, Dispepsia)	1	0,61
DM (COPD, HT)	1	0,61
DM (CVA, susp covid)	1	0,61
DM (Gastritis)	1	0,61
DM (GERD, HF)	1	0,61
DM (Hepatitis)	1	0,61
DM (HF)	1	0,61
DM (HF, CAD, COPD)	1	0,61
DM (HF, Hepatitis)	1	0,61
DM (HT, HF, dispepsia)	1	0,61
DM (HT, CA infark)	1	0,61
DM (HT, CAD, Gerd)	1	0,61
DM (HT, CAD, Parkinson)	1	0,61
DM (HT, COPD, CVA)	1	0,61
DM (HT, COPD, Dispepsia)	1	0,61
DM (HT, Dispepsia)	1	0,61
DM (HT, HF, CAD)	1	0,61
DM (HT, STEMI)	1	0,61
DM (Myalgia)	1	0,61
DM (PAD)	1	0,61
DM (PAD, Vertigo)	1	0,61
DM T1 (HT)	1	0,61
Total	164	100

Commented [MOU12]: Use the table format, not a images

Please present in English

Based on the DM treatment profile seen in the 12 months of 2022, it shows that the use of anti-diabetic drugs is the most in the use of glargine drugs with a dose strength of 100U/ml in as many as 71 patients. Followed by glulisine drugs with a dose strength of 100U / ml for as many as 57 patients. The third highest use was aspart 100U / ml, as many as 54 patients. Concerning the administration interval of most anti-diabetic drugs given every 24 hours a day with the duration of most drug administration for 4-6 days (Table 3).

Table 3. Profile of Anti-diabetic Drug Use

Karakteristik	Bulan (Tahun 2022)												Total (n=293)	%
	Jan 2022	Feb 2022	Mar 2022	Apr 2022	Mei 2022	Jun 2022	Jul 2022	Agust 2022	Sept 2022	Okt 2022	Nov 2022	Des 2022		
Nama obat														
Acarbos 50 mg	1	0	0	0	0	1	2	0	0	0	0	1	5	1,7
Glulisin 100U/ml	4	7	8	10	5	7	2	3	1	5	3	2	57	19,5
Glargine 100U/ml	8	9	8	8	4	9	4	4	5	7	3	2	71	24,2
Glibenklamid 5 mg	1	0	0	0	0	0	1	1	0	0	0	0	3	1,0
Glikuidon 30 mg	2	0	4	1	0	1	1	1	1	4	2	1	18	6,1
Glimepirid 2 mg	2	0	0	3	2	3	3	3	2	0	0	0		
Glimepirid 4 mg	0	0	0	1	0	1	1	0	2	0	0	0	18	6,1
Glimepirid 3 mg	0	0	0	0	1	0	0	0	0	0	0	0		
Lispro 100U/ml	1	0	0	0	0	1	2	0	1	0	0	2	7	2,4
Detemir 100U/ml	2	0	3	6	3	5	4	2	2	7	2	2	38	13,0
Metformin 500 mg	2	1	0	4	1	2	3	3	1	1	1	1	20	6,8
Aspart 100U/ml	6	3	3	4	2	8	7	4	5	8	4	0	54	18,4
Pioglitazon 30 mg	0	0	0	0	0	0	0	1	0	0	0	0	1	0,3
Degludec-Aspart 100U/ml	0	1	0	0	0	0	0	0	0	0	0	0	1	0,3
Total	29	21	26	36	17	37	29	22	18	32	15	11	293	100
Interval pemberian														
24 jam	15	10	16	20	9	18	14	11	9	20	9	6	157	53,6
12 jam	2	1	0	3	1	3	3	2	3	2	1	0	21	7,2
8 jam	12	10	10	13	7	16	12	9	6	10	4	5	114	38,9
6 jam	0	0	0	0	0	0	0	0	0	0	1	0	1	0,3
Total	29	21	26	36	17	37	29	22	18	32	15	11	293	100

Karakteristik	Bulan (Tahun 2022)												Total (n=293)	%
	Jan 2022	Feb 2022	Mar 2022	Apr 2022	Mei 2022	Jun 2022	Jul 2022	Agust 2022	Sept 2022	Okt 2022	Nov 2022	Des 2022		
Durasi														
1-3 Hari	7	15	12	10	5	20	14	13	9	15	4	3	127	43,3
4-6 Hari	22	6	14	26	12	17	15	9	9	17	11	8	166	56,7
Total	29	21	26	36	17	37	29	22	18	32	15	11	293	100

b. Evaluation of the Use of Anti-diabetic Drugs with ATC/DDD and DU 90% Methods

The results of the evaluation of the use of anti-diabetic drugs using the ATC / DDD method showed that the use of antidiabetics that are often used is glulisine drugs with a dose strength of 100U / ml of 589.02 DDD / 100 days of hospitalization, which means that during 100 days of treatment in the hospital around 589 diabetic patients received glulisine amounting to 40 IU every day. The total number of patients hospitalized and using anti-diabetic drugs from January to December 2022 was 568 days. The second most significant use of anti-diabetic medications is aspart with a dose strength of 100U / ml of 519.72 DDD / 100 days of hospitalization, which means that during 100 days of hospitalization in the hospital around 520 DM patients received aspart drugs of 40 IU (Table 4).

Commented [MOU13]: Use the table format, not a images

Please present in English

162

Table 4. Analysis of the Use of Anti-diabetic Drugs using the ATC / DDD Method

No.	Kode ATC	Nama Obat	DDD (WHO)	LOS (Hari)	DDD/100 Hari rawat inap
1	A10AB06	Glulisin 100 U/ml	40 UI	568 Hari	589,02
2	A10AB05	Aspart 100 U/ml	40 UI		519,72
3	A10BB12	Glimepirid 2 mg, 3 mg, 4 mg	2 mg		474,28
4	A10AE04	Glargine 100 U/ml	40 UI		426,85
5	A10AE05	Detemir 100 U/ml	40 UI		221,38
6	A10BA02	Metformin 500 mg	2 g		198,80
7	A10BB08	Glikuidon 30 mg	60 mg		129,22
8	A10AB04	Lispro 100 U/ml	40 UI		85,20
9	A10BF01	Acarbos 50 mg	0,3 g		43,55
10	A10BB01	Glibenklamid 5 mg	10 mg		31,24
11	A10BG03	Pioglitazon 30 mg	30 mg		22,72
12	A10AD06	Degludec-Aspart 100 U/ml	40 UI		11,36

Commented [MOU14]: Use the table format, not a images

Please present in English

163

164 The 90% DU in this study was used to evaluate the use of anti-diabetic drugs in the top 90% of services
 165 in the population. There were 12 anti-diabetic assessed drugs in 90% DU and showed that glulycin,
 166 aspart, glimepiride 2 mg, glimepiride 3 mg, glimepiride 4 mg, glargine 100U/ml, detemir 100U/ml, and
 167 metformin 500 mg were anti-diabetic drugs that accounted for 90% of the highest anti-diabetic drug use
 168 in the DM patient population at the Bokor Turen Salvation Army Hospital (Table 5).

169

170

Table 5. Profile of Anti-diabetic Drug Use with DU Method 90%

Kode ATC	Nama Obat	DDD/100 Hari rawat inap	Persentase	Persentase Kumulatif	Segmen
A10AB06	Glulisin 100 U/ml	589,02	21,39%	21,39%	Du 90%
A10AB05	Aspart 100 U/ml	519,72	18,88%	40,27%	

171

172

173

174

Table 5. Profile of Anti-diabetic Drug Use with DU Method 90% (Continued)

	Glimepirid 2 mg			
A10BB12	Glimepirid 3 mg	474,28	17,23%	57,49%
	Glimepirid 4 mg			
A10AE04	Glargine 100 U/ml	426,85	15,50%	73,00%
A10AE05	Detemir 100 U/ml	221,38	8,04%	81,04%
A10BA02	Metformin 500 mg	198,80	7,22%	88,26%
A10BB08	Glikuidon 30 mg	129,22	4,69%	92,95%
A10AB04	Lispro 100 U/ ml	85,20	3,09%	96,05%
A10BF01	Acarbos 50 mg	43,55	1,58%	97,63%
A10BB01	Glibenklamid 5 mg	31,24	1,13%	98,76%
A10BG03	Pioglitazon 30 mg	22,72	0,83%	99,59%
A10AD06	Degludec- Aspart 100 U/ ml	11,36	0,41%	100,00%
		2753,33	99,99	

175

176

177 DISCUSSION

178 Insulin glulisine (Apidra) in this study occurred in patients with Type 2 diabetes with comorbidities,
 179 such as hypertension, coronary artery disease (CAD), and chronic kidney disease (CKD). On average,
 180 DM patients who get apidra drugs are given every 8 hours, and this drug is widely presented to 76% of
 181 female patients with an age range of 58-61 years. This insulin is widely recommended at the Bokor
 182 Turen Salvation Army Hospital because it can lower blood sugar levels quickly and has a more
 183 negligible risk of hypoglycemia. Following the 2021 PERKENI guidelines, Apidra is insulin rapid-
 184 acting, generally used with food. Apidra is designed to decrease glucose levels in the blood after a meal
 185 rapidly or when blood glucose levels are high. Apidra is commonly combined with basal insulins such
 186 as Lantus (insulin glargine) and Levemir (insulin detemir) (Soelistijo et al., 2021). A study examining
 187 the clinical effects of Type 2 DM patients with cardiovascular comorbidities where patients used insulin
 188 rapid-acting showed that it could have beneficial effects from insulin glulisine administration associated
 189 with death and stroke. Still, there was no difference in coronary heart disease (CHD) or cardiovascular
 190 disease (CVD) (Svensson et al., 2017).

191

Insulin aspart (novorapid) is given to patients with type 2 diabetes with comorbid HT accompanied by CAD, ST-Elevation Myocardial Infarction (STEMI), and Heart Failure (HF). The interval of insulin administration averaged every 8 hours per day and was used by 66% of female patients with an average of 58-61 years. Like glulisine insulin, this insulin can also quickly lower blood sugar levels and has a low risk of hypoglycemia. This insulin is also rapid-acting, which provides therapeutic effectiveness after 15 minutes, with the peak of therapeutic efficacy occurring within 1-2 hours and can last up to 4-6 hours. In the case of the population in Japan, administering insulin aspart to type 2 DM patients can significantly reduce cardiovascular complications within 5 to 10 years, resulting in improved quality of life and lower costs compared to human insulin (Pollock et al., 2011).

Commented [MOU15]: Why is insulin glulisine more often choice than aspart?

The third most significant use of anti-diabetic drugs is glimepiride doses of 2mg, 3 mg, and 4 mg. This drug is given to patients with type 2 diabetes with cardiovascular comorbidities and an interval of 24 hours per day and primarily female patients (83%) with an average age of 58 years. Regular glimepiride is combined with insulin or another oral medication such as metformin. In the case study of type 2 DM patients with CVD, the average patient suffering from DM was around 5.7 ± 4.8 years. CVD suffered by type 2 DM patients in the form of hypertension (68.5% of patients); dyslipidemia (47.9% of patients); CAD (25.4% of patients); Transient Ischemic Attack (TIA) in 3.6% of patients; peripheral artery disease (PAD) accounted for 4.8% of patients and heart failure in 2.9% of patients. Type 2 DM patients with various comorbidities receive the drug glimepiride/metformin Fixed Dose Combination (FDC) as a first-line therapy. As many as 68.2% of FDC patients achieved blood pressure within optimal limits. Most of the other patients experienced an increase in glycemic parameters and a change in body weight of about 18.4%. 59.2% of patients experienced weight loss (Ray et al., 2022). Glimepiride is a sulfonylurea class drug that has pharmacological effects to increase insulin production by pancreatic beta cells. The most common side effects are hypoglycemia and weight gain.

Commented [MOU16]: Why is metformin not among the top three most widely used antidiabetic drug? Wasn't it the first choice?

Commented [MOU17]: Please compare the result of this research with the previous research

The limitation of this study is that data were taken only retrospectively from patient medical record data and did not make direct observations on DM patients.

Commented [MOU18]: Please explain benefit and follow up from the result of research for the hospital or science?

CONCLUSION

Based on the analysis of demographic data and treatment profiles of DM patients, DM patients are dominated by the age group of 46-65 years, with women who use anti-diabetic drugs more. Patients with comorbidities use more anti-diabetic medications than patients with non-comorbid DM. Evaluation of anti-diabetic drugs ATC / DDD method shows that insulin glulisine (Apidra) is most used in DM patients

with and without comorbidities. Through the DU method, 90% of glulisine, aspart, glargine, glimepiride, detemir, and metformin drugs were most widely used in DM patients.

CONFLICT OF INTEREST

All authors declare no conflict of interest.

REFERENCES

- Ali, M.D. et al. (2022) 'Evaluation of drug utilization pattern and cost associated with diabetes mellitus Type two management in Saudi Arabia', *Brazilian Journal of Pharmaceutical Sciences*, 58, p. E20681. Available at: <https://doi.org/10.1590/s2175-97902022e20681>.
- Almasdy, D. et al. (2015) 'Evaluation of the Use of Anti-diabetic Drugs in Type-2 Diabetes Mellitus Patients in a Government Hospital of Padang City – West Sumatra', *Journal of Pharmaceutical & Clinical Science*, 2(1), p. 104. Available at: <https://doi.org/10.29208/jsfk.2015.2.1.58>.
- DiPiro, J.T. (ed.) (2020) *Pharmacotherapy: a pathophysiologic approach*. Eleventh edition. New York: McGraw Hill Medical.
- Okoro, R.N., Nmeka, C. and Erah, P.O. (2018) 'Utilization study of antidiabetes medicines at a tertiary care hospital in Nigeria', *Future Journal of Pharmaceutical Sciences*, 4(2), pp. 109–115. Available at: <https://doi.org/10.1016/j.fjps.2017.11.004>.
- Pitasari, N.W.N., Andayani, T.M. and Wijayanti, T. (2022) 'Evaluation of the Use of Antidiabetic Drugs in Patients of Back-Referral Program at the Demak District Pharmacy', *Journal of Management and Pharmacy Practice*, 12(2), p. 125. Available at: <https://doi.org/10.22146/jmpf.73841>.
- Pollock, R.F. et al. (2011) 'The cost effectiveness of rapid-acting insulin aspart compared with human insulin in type 2 diabetes patients: an analysis from the Japanese third-party payer perspective', *Journal of Medical Economics*, 14(1), pp. 36–46. Available at: <https://doi.org/10.3111/13696998.2010.541045>.
- Ray, S. et al. (2022) 'Usage Pattern of Glimepiride/Metformin Fixed-dose Combination in Type 2 Diabetes Patients with CVD or at Risk of CVD: An Experience in Indian Setting', *Asian Journal of Diabetology*, 23(2), pp. 13–19.
- RI, K. (2017) 'Technical Guidelines for Evaluation of Drug Use in Health Facilities'.
- Soelistijo, S.A. dkk. (2021) 'Guidelines for the management and prevention of adult type 2 diabetes mellitus in INDONESIA - 2021', *PB. PERKENI*, p. 119.
- Svensson, A.-M. et al. (2017) 'Clinical effects, cardiovascular and renal outcomes associated with rapid-acting insulin analogs among individuals with type 2 diabetes: a nation-wide observational cohort study', *Clinical Diabetes and Endocrinology*, 3(1), p. 5. Available at: <https://doi.org/10.1186/s40842-017-0043-2>.

Commented [MOU19]: The number of references not enough

- 260 Tahar, N.dkk. (2020) 'EVALUATION OF THE USE OF ORAL ANTI-DIABETIC DRUGS IN TYPE
261 2 DIABETES MELLITUS PATIENTS USING ATC/DDD AND DU 90% METHODS', *The*
262 *2nd Alauddin Pharmaceutical Conference and Expo (ALPHA-C) 2020* [Preprint]. Available at:
263 <https://doi.org/10.24252/kesehatan.v1i1.18380>.
- 264 WHO (2023) *Diabetes*. Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes>
265 (Accessed: September 21, 2023).

3. Lampiran reviewer kedua tahap pertama 27 November 2023

Jurnal Farmasi Galenika (Galenika Journal of Pharma... Tasks 0 English View Site godoliva13

Reviewer's Attachments

Q Search

57083-1	16624-Article Text-56939-1-4-20231020 review 1.docx	November 20, 2023
57216-1	16624-Article Text-56939-1-4-20231020.docx	November 27, 2023

Revisions

Q Search Upload File

57681-1	Article Text, Revisi ATC:DDD Diabetes Mellitus.docx	December 27, 2023	Article Text
---------	---	-------------------	--------------

Review Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
No Items				

[https://bestjournal.untad.ac.id/index.php/Galenika/\\$?call\\$/\\$\\$tab/author-dashboard/author-dashboard-review-round-tab/fetch-review-round-info?submissionId=16624&stageId=3&reviewRoundId=11843](https://bestjournal.untad.ac.id/index.php/Galenika/$?call$/$$tab/author-dashboard/author-dashboard-review-round-tab/fetch-review-round-info?submissionId=16624&stageId=3&reviewRoundId=11843)

4. Submit revisi artikel tahap pertama 27 Desember 2023

Jurnal Farmasi Galenika (Galenika Journal of Pharma... Tasks 0 English View Site godoliva13

Reviewer's Attachments

Q Search

57083-1	16624-Article Text-56939-1-4-20231020 review 1.docx	November 20, 2023
57216-1	16624-Article Text-56939-1-4-20231020.docx	November 27, 2023

Revisions

Q Search Upload File

57681-1	Article Text, Revisi ATC:DDD Diabetes Mellitus.docx	December 27, 2023	Article Text
---------	---	-------------------	--------------

Review Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
No Items				

[https://bestjournal.untad.ac.id/index.php/Galenika/\\$?call\\$/\\$\\$tab/author-dashboard/author-dashboard-review-round-tab/fetch-review-round-info?submissionId=16624&stageId=3&reviewRoundId=11843](https://bestjournal.untad.ac.id/index.php/Galenika/$?call$/$$tab/author-dashboard/author-dashboard-review-round-tab/fetch-review-round-info?submissionId=16624&stageId=3&reviewRoundId=11843)

5. Lampiran reviewer tahap kedua 5 Januari 2024

The screenshot displays the 'JFG Editor Decision' page. At the top, there's a header with 'Jurnal Farmasi Galenika (Galenika Journal of Pharmacy)', 'Tasks 0', and user information 'English', 'View Site', and 'godeliva13'. Below the header, there's a section for 'Reviewer's Attachments' with a search bar. It lists one attachment: '57954-1', '16624-Article Text-57791-1-4-20231227 review 2.docx', dated 'January 5, 2024'. Below this is a 'Revisions' section with a search bar and an 'Upload File' button. It lists one revision: '58212-1', 'Article Text, Revisi ATC/DDD.docx', dated 'January 31, 2024', with the type 'Article Text'. At the bottom is a 'Review Discussions' section with an 'Add discussion' button. It shows a table with columns: Name, From, Last Reply, Replies, and Closed. The table is currently empty, showing 'No Items'.

6. Keputusan Editor “Revisions Required” 26 Januari 2024

The screenshot shows a 'Notifications' window titled '[JFG] Editor Decision' dated '2024-01-26 01:54 AM'. The notification is from 'Godeliva Adriani Hendra'. The text of the notification reads: 'We have reached a decision regarding your submission to Jurnal Farmasi Galenika (Galenika Journal of Pharmacy) (e-Journal), "Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients". Our decision is: Revisions Required. Please note that you have to submit the revised version of your manuscript as soon as possible with the respond to reviewers comments by answering their comments point by point in a separated file (rebuttal letter). We will not process the manuscript if the manuscript was not revised as reviewer suggestion or if there is no a rebuttal letter. Thanks and regards. Amelia Rumi, Universitas Tadulako, amelia.rumi@gmail.com'. The background shows a sidebar with 'Submissions' and 'Round 1' selected, and a list of notifications on the right.

Lampiran Reviewer Tahap Kedua

5 Januari 2024

Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients

ABSTRACT

Background: Diabetes Mellitus (DM), which results from insulin resistance, is closely linked to long-term damage to pancreatic beta cells, organ dysfunction, and organ failure, particularly in the eyes, kidneys, nerves, heart, and blood vessels. The development of the ATC/DDD system is necessary to increase global drug knowledge, ensure equitable drug availability, and promote responsible drug use. **Objectives:** This study aimed to identify data on patient characteristics and oral anti-diabetic drugs and insulin using the ATC/DDD and DU90% methods. **Material and Methods:** This study was a cross-sectional evaluation of oral anti-diabetic medications and insulin using the ATC/DDD method and DU90% at the Bala Keselamatan Bokor Turen Hospital from January to December 2022. The inclusion criteria were individuals diagnosed with type 1 or type 2 diabetes mellitus, with or without comorbidities, who were treated with oral anti-diabetic medications and a combination of insulin and oral medications. In the present study, the incomplete medical record data served as the exclusion criteria. The sample for this investigation consisted of 238 patients selected using the complete sampling technique. **Results:** The majority of patients who used oral anti-diabetic medications and insulin were between the ages of 46 and 65 (71.34%), were female (73.2%), had standard body mass index (BMIs between 18.5 and 25), and had diagnoses of DM + HT (20.73%) and Type II DM (18.29%). Glulisin was the most commonly used anti-diabetic drug, accounting for 589 DDD/100 days of hospitalization, and DU90% was 21.39%. **Conclusion:** Glulisin is the most frequently prescribed anti-diabetic medication at Bokor Turen Hospital.

Keywords: Antidiabetic Oral and Insulin; Diabetes Mellitus; ATC/DDD; DU 90%

ABSTRAK

Latar Belakang: Resistensi insulin yang berkembang menjadi Diabetes Melitus (DM) erat hubungannya dengan terjadinya kerusakan sel beta pankreas dalam jangka panjang, disfungsi organ, dan kegagalan organ terutama pada mata, ginjal, saraf, jantung, dan pembuluh darah. Pengembangan sistem ATC/DDD diperlukan untuk meningkatkan pengetahuan tentang penggunaan obat di seluruh dunia, memastikan ketersediaan obat secara merata, dan mendorong penggunaan obat yang bijak. Tujuan penelitian ini adalah mengidentifikasi data karakteristik pasien serta obat anti-diabetes oral dan insulin menggunakan metode ATC/DDD dan DU90%. Bahan dan Metode: Rancangan penelitian ini menggunakan *cross-sectional study* yang mengevaluasi obat anti-diabetes oral dan insulin dengan metode ATC/DDD dan DU90% di RS Bala Keselamatan Bokor Turen bulan Januari hingga Desember 2022. Kriteria inklusi terdiri dari pasien DM tipe 1 dan tipe 2 dengan dan atau tanpa komorbid, menggunakan obat anti-diabetes oral dan insulin/kombinasi keduanya. Sedangkan, kriteria eksklusinya berupa data rekam medis tidak lengkap. Populasi penelitian ini sebanyak 238 pasien menggunakan teknik total sampling. Hasil: Data karakteristik pasien pengguna obat anti-diabetes oral dan insulin terbanyak berusia 46 hingga 65 tahun (71,34%); jenis kelamin perempuan (73,2%); Indeks Masa Tubuh normal sebesar 53,66% (IMT= 18,5-25); diagnose DM+HT (20,73%) dan DM Tipe II (18,29%). Penggunaan obat anti-diabetes tertinggi adalah obat glulisin sebesar 589 DDD/100 hari rawat inap dan DU90% sebesar 21,39%. Kesimpulan: Obat anti-diabetes yang paling banyak digunakan di RS Bokor Turen adalah glulisin.

Kata kunci: Obat Anti-diabetes Oral dan Insulin; Diabetes Melitus; DDD; DU 90%

INTRODUCTION

Diabetes Mellitus (DM) is a chronic metabolic disorder. Risk factors, damage to insulin secretion, and sensitivity lead to increased blood glucose and changes in fat and protein metabolism (DiPiro, 2020). The lack of insulin function contributes to the development of microvascular, macrovascular, and neuropathy as a chronic consequence of DM (Almasdy *et al.*, 2015).

The World Health Organization (WHO) in 2023 states that in 2014, 8.5% of adults aged 18 years and over have diabetes. In 2019, diabetes was the direct cause of 1.5 million deaths, and 48% of all deaths from diabetes occurred before the age of 70. Furthermore, 460,000 deaths from kidney disease are caused by diabetes, and elevated blood glucose causes about 20% of deaths from cardiovascular disease (WHO, 2023). DM is more common in low- and moderate-income countries and countries undergoing major economic and demographic transformations. DM is a significant global health problem and requires proper prevention and management measures. DM in Indonesia is currently a serious problem. Most DM sufferers are type 2 DM groups. Therefore, it is essential to evaluate the use of the drug as a basis for selection to ensure that the drug is used appropriately, safely, and efficiently (Pitasari, Andayani, and Wijayanti, 2022).

There is a need to develop ATC/DDD systems to acquire knowledge about drug use worldwide to achieve equitable drug availability and prudent drug use, especially in developing countries. The primary purpose of the ATC/DDD system is to facilitate research on drug use and improve the overall quality of drug use (Tahar et al., 2020). A retrospective study using a *cross-sectional study*, which evaluated drug use patterns and costs associated with Type 2 DM in Saudi Arabia, showed that biguanide (metformin) was most widely prescribed as a monotherapy drug followed by a *fixed-dose combination*. The effectiveness of monotherapy drugs decreases with the duration of treatment; in these cases, combination drugs are prescribed. The most commonly prescribed combination drug is a biguanide with sulfonylurea/biguanide with thiazolidinedione, according to guidelines by the *American Diabetes Association* (ADA). The combination of sitagliptin and metformin is most widely preferred and widely prescribed in *fixed-dose combination therapy*, followed by vildagliptin and metformin (Ali et al., 2022). Combination drugs are used when a single pill cannot achieve the desired blood glucose level in diabetic patients (Okoro, Nmeka, and Erah, 2018).

Evaluation of the use of anti-diabetic drugs with ATC/DDD and DU90% methods can provide insight into the dominant drug use patterns and the extent to which these drugs follow existing treatment recommendations and guidelines. Therefore, this study evaluated oral anti-diabetic drugs and insulin using ATC/DDD and DU90% in DM patients with and without comorbidities at the Bokor Turen Salvation Army Hospital, a hospital with type D accreditation.

MATERIAL AND METHODS

Research Methods

The study design used a *cross-sectional study* with retrospective data collection through patient medical records. Quantitative evaluation of the use of anti-diabetic drugs using ATC / DDD and DU 90%

Commented [MOU1]: Explain further, why this research carried out at this hospital. Author can explain the prevalence Dm in this hospital, or the other reason

techniques. This research has gone through the health research ethics committee with no E.5.a/145/KEPKUMM/V/2023.

Population and Sample

This population consists of patients hospitalized with a diagnosis of DM and without comorbidities at the Bokor Turen Salvation Army Hospital. The study sample included patients hospitalized with a diagnosis of DM and without comorbidities from January to December 2022 and who met the inclusion and exclusion criteria. Inclusion criteria include patients aged ≥ 17 years with a diagnosis of DM and without comorbidities. Exclusion criteria were patients whose medical record data was incomplete.

Sampling Techniques

The sampling technique was total sampling, where the number of samples was equal to a population of 164 patients and met the criteria for inclusion and exclusion of the study.

Data Analysis

The calculation in evaluating the use of ATC / DDD method anti-diabetic drugs in the hospitalization of the Bokor Turen Salvation Army Hospital uses the formula:

$$\frac{\text{DDD}}{100} \text{ bed} - \text{days} = \frac{\text{total antibiotics (gram)}}{\text{DDD WHO (gram)}} \times \frac{100}{\text{LOS}}$$

DU 90% is used to identify the amount of drug used as much as 90% of the total use of prescribed medicines and compare it with the amount of residual drug use (RI, 2017). The efficiency of drug use must be observed if the amount of drug use in 10% is more. The DU value of 90% is known after calculating DDD / 100 days of hospitalization per year.

RESULTS

a. Demographic Characteristics of Diabetes Mellitus Patients

The demographic characteristics of Diabetes Mellitus patients at the Bokor Turen Salvation Army Hospital from January to December 2022 were 238 patients in age, gender, BMI, diagnosis, comorbidities, drug names, administration intervals, and duration of administration. This study's sample size was 164 patients. Judging from the most significant number, in the age characteristics of 117 patients aged 46-65, as many as 120 were female, and as many as 88 patients had a regular Body Mass Index (BMI) of 18.5-25 (Table 1).

Commented [MOU2]: Give information that this sample meets the inclusion criteria

Table 1. Data on the Characteristics of Diabetes Mellitus Patients

Characteristic of Patient	Parameter	Total N= (164)	%
Age	17-45	12	7,32
	46-65	117	71,34
	66-95	35	21,34
Mean \pm SD	58,78 \pm 2,52		
Gender	Female	120	73,2
	Male	44	26,8
BMI	Underweight <18,5	6	3,66
	Healthy weight 18,5-25	88	53,66
	Overweight >25	70	42,68
Mean \pm SD	24,63 \pm 1,08		

The description of DM and or without comorbidities at the Bokor Turen Salvation Army Hospital showed that the most patients with a diagnosis of DM and Hypertension (HT) were 34 patients (20.73%), followed by patients with a diagnosis of Type 2 DM as many as 28 patients (18.29%). At the same time, DM patients with other comorbidities have fewer than ten patients (Table 2).

Table 2. Description of Patients with Diabetes Mellitus and or Without Comorbidities

Diagnosis with comorbidities	Number of Patients (n= 164)	%
DM (HT)	34	20,73
DM Type 2	28	18,29
DM (HT, CAD)	6	3,66
DM (anemia)	5	3,05
DM (CAD)	5	3,05
DM (CKD)	5	3,05
DM (CVA)	5	3,05
DM (COPD)	5	3,05
DM (CVA, HT)	4	2,44
DM (anemia, thalassemia)	3	1,83
DM (dyspepsia)	3	1,83
DM (GERD, HT)	3	1,83
DM (HT, anemia)	3	1,83
DM (asthma)	2	1,22
DM (DKD)	2	1,22
DM (DKD, HT)	2	1,22
DM (HT, CKD)	2	1,22
DM (HT, HF)	2	1,22
DM (HT, Vertigo)	2	1,22
DM (STEMI)	2	1,22

CAD= coronary artery disease, CKD= chronic kidney disease, CVA= Cerebrovascular Accident, GERD= Gastroesophageal Reflux Disease, DKD= Diabetic Kidney Disease, HF= Heart Failure

132 **Table 2.** Description of Patients with Diabetes Mellitus and or Without Comorbidities (continued)

Diagnosis with comorbidities	Number of Patients (n= 164)	%
DM (Vertigo)	2	1,22
DM (GERD)	1	0,61
DM (Anemia, PAD)	1	0,61
DM (AKI)	1	0,61
DM (Anemia, CKD)	1	0,61
DM (Anemia, Dyspepsia)	1	0,61
DM (Angina, Thalassemia)	1	0,61
DM (CAD, COPD)	1	0,61
DM (CAD, CVA, HT)	1	0,61
DM (CAD, HT)	1	0,61
DM (CAD, STEMI)	1	0,61
DM (CAD, TB)	1	0,61
DM (CKD, HT, TB)	1	0,61
DM (CKD, STEMI)	1	0,61
DM (CKD, Thalassemia)	1	0,61
DM (COPD, Dyspepsia)	1	0,61
DM (COPD, HT)	1	0,61
DM (CVA, susp. COVID-19)	1	0,61
DM (Gastritis)	1	0,61
DM (GERD, HF)	1	0,61
DM (Hepatitis)	1	0,61
DM (HF)	1	0,61
DM (HF, CAD, COPD)	1	0,61
DM (HF, Hepatitis)	1	0,61
DM (HT, HF, dyspepsia)	1	0,61
DM (HT, CA infark)	1	0,61
DM (HT, CAD, GERD)	1	0,61
DM (HT, CAD, Parkinson)	1	0,61
DM (HT, COPD, CVA)	1	0,61
DM (HT, COPD, Dyspepsia)	1	0,61
DM (HT, Dyspepsia)	1	0,61
DM (HT, HF, CAD)	1	0,61
DM (HT, STEMI)	1	0,61
DM (myalgia)	1	0,61
DM (PAD)	1	0,61
DM (PAD, Vertigo)	1	0,61
DM Type 1 (HT)	1	0,61

133 PAD= Peripheral Artery Disease, AKI= Acute Kidney Injury, COPD= Coronary
 134 Obstructive Pulmonary Disease, STEMI= ST-Elevation Myocardial Infarction
 135

136 Based on the DM treatment profile seen in the 12 months of 2022, it shows that the use of anti-diabetic
 137 drugs is the most in the use of glargine drugs with a dose strength of 100U/ml in as many as 71 patients.
 138 This was followed by glulisine drugs with a dose strength of 100U / ml for as many as 57 patients. The
 139 third highest use was aspart 100U / ml, as many as 54 patients. Concerning the administration interval
 140 of most anti-diabetic drugs, they are given every 24 hours a day, with the duration of most drug
 141 administration being 4-6 days (Table 3).

Table 3. Profile of Anti-diabetic Drug Use in Year 2022

Characteristic of Patient	Parameter	Number of Patients (N= 293)	%
Type of drug	Acarbose 50 mg	5	1,7
	Glulisine 100 IU/ml	57	19,5
	Glargine 100 IU/ml	71	24,2
	Glibenclamide 5 mg	3	1
	Gliquadone 30 mg	18	6,1
	Glimepiride 2 mg	18	6,1
	Glimepiride 3 mg	18	6,1
	Glimepiride 4 mg	18	6,1
	Lispro 100 IU/ml	7	2,4
	Detemir 100 IU/ml	38	13
	Metformin 500 mg	20	6,8
	Aspart 100 IU/ml	54	18,4
	Pioglitazone 30mg	1	0,3
	Degludec + Aspart 100 IU/ml	1	0,3
Interval of administration	24 hours	157	53,6
	12 hours	21	7,2
	8 hours	114	38,9
	6 hours	1	0,3
Duration of administration	1-3 days	127	43,3
	4-6 days	166	56,7

are often used is glulisine drugs with a dose strength of 100U / ml of 589.02 DDD / 100 days of hospitalization, which means that during 100 days of treatment in the hospital around 589 diabetic patients received glulisine amounting to 40 IU every day. The total number of patients hospitalized and using anti-diabetic drugs from January to December 2022 was 568 days. The second most significant use of anti-diabetic medications is aspart with a dose strength of 100U / ml of 519.72 DDD / 100 days of hospitalization, which means that during 100 days of hospitalization in the hospital around 520 DM patients received aspart drugs of 40 IU (Table 4).

Commented [MOU3]: correct the format

Table 4. Analysis of the Use of Anti-diabetic Drugs using the ATC / DDD Method

No.	ATC code	Type of Drugs	DDD (WHO)	Length of Stay (LOS)	DDD/100 bed-days
1	A10AB06	Glulisine 100 IU/ml	40 IU	568 days	589,02

2	A10AB05	Aspart 100 IU/ml	40 IU	519,72
		Glimepirid 2 mg		474,28
3	A10BB12	Glimepirid 3 mg	2 mg	426,85
		Glimepirid 4 mg		221,38
4	A10AE04	Glargine 100 IU/ml	40 IU	426,85
5	A10AE05	Detemir 100 IU/ml	40 IU	221,38
6	A10BA02	Metformin 500 mg	2 g	198,80
7	A10BB08	Gliquadone 30 mg	60 mg	129,22
8	A10AB04	Lispro 100 IU/ml	40 IU	85,20
9	A10BF01	Acarbose 50 mg	0,3 g	43,55
10	A10BB01	Glibenclamide 5 mg	10 mg	31,24
11	A10BG03	Pioglitazone 30 mg	30 mg	22,72
12	A10AD06	Degludec + aspart 100 IU/ml	40 IU	11,36

Commented [MOU4]:

Commented [MOU5]:

The 90% DU in this study was used to evaluate the use of anti-diabetic drugs in the top 90% of services in the population. There were 12 anti-diabetic assessed drugs in 90% DU and showed that glulycin, aspart, glimepiride 2 mg, glimepiride 3 mg, glimepiride 4 mg, glargine 100U/ml, detemir 100U/ml, and metformin 500 mg were anti-diabetic drugs that accounted for 90% of the highest anti-diabetic drug use in the DM patient population at the Bokor Turen Salvation Army Hospital (Table 5).

Commented [MOU6]: ??

Commented [MOU7]: Milligrams and please correct all this abbreviations in this article, in the table also

Commented [MOU8]:

Commented [MOU9]:

Commented [MOU10]:

Table 5. Profile of Anti-diabetic Drug Use with DU Method 90%

ATC code	Type of Drugs	DDD/100 bed-days	Percentage	Cumulative	DU 90% segment
A10AB06	Glulisine 100 IU/ml	589,02	21,39%	21,39%	
A10AB05	Aspart 100 IU/ml	519,72	18,88%	40,27%	
	Glimepiride 2 mg				
A10BB12	Glimepiride 3 mg	474,28	17,23%	57,49%	DU90%
	Glimepiride 4 mg				
A10AE04	Glargine 100 IU/ml	426,85	15,50%	73,00%	
A10AE05	Detemir 100 IU/ml	221,38	8,04%	81,04%	
A10BA02	Metformin 500 mg	198,80	7,22%	88,26%	
A10BB08	Gliquadone 30 mg	129,22	4,69%	92,95%	
A10AB04	Lispro 100 IU/ml	85,20	3,09%	96,05%	
A10BF01	Acarbose 50 mg	43,55	1,58%	97,63%	
A10BB01	Glibenclamide 5 mg	31,24	1,13%	98,76%	DU 10%
A10BG03	Pioglitazone 30 mg	22,72	0,83%	99,59%	
A10AD06	Degludec+Aspart 100 IU/ml	11,36	0,41%	100%	

Commented [MOU11]:

DISCUSSION

Insulin glulisine (Apidra) in this study occurred in patients with Type 2 diabetes with comorbidities, such as hypertension, coronary artery disease (CAD), and chronic kidney disease (CKD). On average, DM patients who get apidra drugs are given every 8 hours, and this drug is widely presented to 76% of

female patients with an age range of 58-61 years. This insulin is widely recommended at the Bokor Turen Salvation Army Hospital because it can lower blood sugar levels quickly and has a more negligible risk of hypoglycemia. Following the 2021 PERKENI guidelines, Apidra is a rapid-acting insulin generally used with food. Apidra is designed to decrease glucose levels in the blood rapidly after a meal or when blood glucose levels are high. Apidra is commonly combined with basal insulins such as Lantus (insulin glargine) and Levemir (insulin detemir) (Soelistijo et al., 2021). A study examining the clinical effects of Type 2 DM patients with cardiovascular comorbidities where patients used insulin rapid-acting showed that it could have beneficial effects from insulin glulisine administration associated with death and stroke. Still, there was no difference in coronary heart disease (CHD) or cardiovascular disease (CVD) (Svensson et al., 2017).

Insulin aspart (novorapid) is given to patients with type 2 diabetes with comorbid HT accompanied by CAD, ST-Elevation Myocardial Infarction (STEMI), and Heart Failure (HF). The interval of insulin administration averaged every 8 hours per day and was used by 66% of female patients with an average of 58-61 years. Like glulisine insulin, this insulin can also quickly lower blood sugar levels and has a low risk of hypoglycemia. This insulin is also rapid-acting, which provides therapeutic effectiveness after 15 minutes, with the peak of therapeutic efficacy occurring within 1-2 hours and can last up to 4-6 hours. In the case of the population in Japan, administering insulin aspart to type 2 DM patients can significantly reduce cardiovascular complications within 5 to 10 years, resulting in improved quality of life and lower costs compared to *human insulin* (Pollock et al., 2011).

The third most significant use of anti-diabetic drugs is glimepiride doses of 2mg, 3 mg, and 4 mg. This drug is given to patients with type 2 diabetes with cardiovascular comorbidities and an interval of 24 hours per day and primarily female patients (83%) with an average age of 58 years. Regular glimepiride is combined with insulin or another oral medication such as metformin. In the case study of type 2 DM patients with CVD, the average patient suffering from DM was around 5.7 ± 4.8 years. CVD suffered by type 2 DM patients in the form of hypertension (68.5% of patients); dyslipidemia (47.9% of patients); CAD (25.4% of patients); Transient Ischemic Attack (TIA) in 3.6% of patients; peripheral artery disease (PAD) accounted for 4.8% of patients and heart failure in 2.9% of patients. Type 2 DM patients with various comorbidities receive the drug glimepiride/metformin Fixed Dose Combination (FDC) as a first-line therapy because glimepiride/metformin FDC is more effective than metformin alone for type 2 DM patients with comorbidities. As many as 68.2% of FDC patients achieved blood pressure within optimal limits. Most of the other patients experienced an increase in glycemic parameters and a change in body weight of about 18.4%. 59.2% of patients experienced weight loss (Ray et al., 2022). Glimepiride is a

Commented [MOU12]: Past tenses

Commented [MOU13]: Why is insulin glulisine more often choice than aspart?

Commented [MOU14]:

Commented [MOU15]: Why is metformin not among the top three most widely used antidiabetic drug? Wasn't it the first choice?

sulfonylurea class drug that has pharmacological effects to increase insulin production by pancreatic beta cells. The most common side effects are hypoglycemia and weight gain.

Commented [MOU16]: Add discussion for the other drugs that include DU90 % : glargine, detemir and metformin. Please compare the result of this research with the previous research.

The limitation of this study is that data were taken only retrospectively from patients' medical record data, and no direct observations were made on DM patients. For Bala Keselamatan Bokor Turen Hospital to provide an overview of drug use patterns in patients with diabetes mellitus. The results of this study can later be used as a basis for determining the selection of the use of antidiabetic drugs for patients in accordance with applicable standards for patients in accordance with applicable standards.

CONCLUSION

Based on the analysis of demographic data and treatment profiles of DM patients, DM patients are dominated by the age group of 46-65 years, with women who use anti-diabetic drugs more. Patients with comorbidities use more anti-diabetic medications than patients with non-comorbid DM. Evaluation of anti-diabetic drugs ATC / DDD method shows that insulin glulisine (Apidra) is most used in DM patients with and without comorbidities. Through the DU method, 90% of glulisine, aspart, glargine, glimepiride, detemir, and metformin drugs were most widely used in DM patients.

CONFLICT OF INTEREST

All authors declare no conflict of interest.

REFERENCES

Commented [MOU17]: The number of references not enough, please add more

- Ali, M.D. et al. (2022) 'Evaluation of drug utilization pattern and cost associated with diabetes mellitus Type two management in Saudi Arabia', *Brazilian Journal of Pharmaceutical Sciences*, 58, p. E20681. Available at: <https://doi.org/10.1590/s2175-97902022e20681>.
- Almasdy, D. et al. (2015) 'Evaluation of the Use of Anti-diabetic Drugs in Type-2 Diabetes Mellitus Patients in a Government Hospital of Padang City – West Sumatra', *Journal of Pharmaceutical & Clinical Science*, 2(1), p. 104. Available at: <https://doi.org/10.29208/jsfk.2015.2.1.58>.
- DiPiro, J.T. (ed.) (2020) *Pharmacotherapy: a pathophysiologic approach*. Eleventh edition. New York: McGraw Hill Medical.
- Okoro, R.N., Nmeka, C. and Erah, P.O. (2018) 'Utilization study of antidiabetes medicines at a tertiary care hospital in Nigeria', *Future Journal of Pharmaceutical Sciences*, 4(2), pp. 109–115. Available at: <https://doi.org/10.1016/j.fjps.2017.11.004>.

- 255 Pitasari, N.W.N., Andayani, T.M. and Wijayanti, T. (2022) 'Evaluation of the Use of Antidiabetic Drugs
256 in Patients of Back-Referral Program at the Demak District Pharmacy', *Journal of Management
257 and Pharmacy Practice*, 12(2), p. 125. Available at: <https://doi.org/10.22146/jmpf.73841>.
- 258 Pollock, R.F. et al. (2011) 'The cost-effectiveness of rapid-acting insulin aspart compared with human
259 insulin in type 2 diabetes patients: an analysis from the Japanese third-party payer perspective',
260 *Journal of Medical Economics*, 14(1), pp. 36–46. Available at:
261 <https://doi.org/10.3111/13696998.2010.541045>.
- 262 Ray, S. et al. (2022) 'Usage Pattern of Glimepiride/Metformin Fixed-dose Combination in Type 2
263 Diabetes Patients with CVD or at Risk of CVD: An Experience in Indian Setting', *Asian Journal
264 of Diabetology*, 23(2), pp. 13–19.
- 265 RI, K. (2017) 'Technical Guidelines for Evaluation of Drug Use in Health Facilities'.
- 266 Soelistijo, S.A. dkk. (2021) 'Guidelines for the management and prevention of adult type 2 diabetes
267 mellitus in INDONESIA - 2021', *PB. PERKENI*, p. 119.
- 268 Svensson, A.-M. et al. (2017) 'Clinical effects, cardiovascular and renal outcomes associated with rapid-
269 acting insulin analogs among individuals with type 2 diabetes: a nation-wide observational
270 cohort study', *Clinical Diabetes and Endocrinology*, 3(1), p. 5. Available at:
271 <https://doi.org/10.1186/s40842-017-0043-2>.
- 272 Tahar, N.dkk. (2020) 'EVALUATION OF THE USE OF ORAL ANTI-DIABETIC DRUGS IN TYPE
273 2 DIABETES MELLITUS PATIENTS USING ATC/DDD AND DU 90% METHODS', *The
274 2nd Alauddin Pharmaceutical Conference and Expo (ALPHA-C) 2020* [Preprint]. Available at:
275 <https://doi.org/10.24252/kesehatan.v1i1.18380>.
- 276 WHO (2023) *Diabetes*. Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes>
277 (Accessed: September 21, 2023).

7. Submit revisi artikel tahap kedua 31 Januari 2024

Jurnal Farmasi Galenika (Galenika Journal of Pharma...Tasks0EnglishView Sitegodeliva13


[\[JFG\] Editor Decision](#)

2024-02-23 08:24 AM

[\[JFG\] Editor Decision](#)

2024-03-04 12:33 AM


Reviewer's AttachmentsSearch

 57954-1

16624-Article Text-57791-1-4-20231227 review 2.docx

January 5, 2024

RevisionsSearchUpload File

 58212-1

Article Text, Revisi ATC:DDD.docx

January 31, 2024

Article Text

Review DiscussionsAdd discussion

Name	From	Last Reply	Replies	Closed
No Items				

8. Lampiran reviewer tahap ketiga 9 Februari 2024

Jurnal Farmasi Galenika (Galenika Journal of Pharma...Tasks0EnglishView Sitegodeliva13


[\[JFG\] Editor Decision](#)

2024-02-23 08:24 AM

[\[JFG\] Editor Decision](#)

2024-03-04 12:33 AM


Reviewer's AttachmentsSearch

 58259-1

16624-Article Text-58244-1-4-20240131.docx

February 9, 2024

RevisionsSearchUpload File

 58277-1

Article Text, Revisi anti-diabetic drug.docx

February 15, 2024

Article Text

Review DiscussionsAdd discussion

Name	From	Last Reply	Replies	Closed
No Items				

Lampiran Reviewer Tahap Ketiga

9 Februari 2024

Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients

ABSTRACT

Background: Diabetes Mellitus (DM), which results from insulin resistance, is closely linked to long-term damage to pancreatic beta cells, organ dysfunction, and organ failure, particularly in the eyes, kidneys, nerves, heart, and blood vessels. The development of the ATC/DDD system is necessary to increase global drug knowledge, ensure equitable drug availability, and promote responsible drug use. **Objectives:** This study aimed to identify data on patient characteristics and oral anti-diabetic drugs and insulin using the ATC/DDD and DU90% methods. **Material and Methods:** This study was a cross-sectional evaluation of oral anti-diabetic medications and insulin using the ATC/DDD method and DU90% at the Bala Keselamatan Bokor Turen Hospital from January to December 2022. The inclusion criteria were individuals diagnosed with type 1 or type 2 diabetes mellitus, with or without comorbidities, who were treated with oral anti-diabetic medications and a combination of insulin and oral medications. In the present study, the incomplete medical record data served as the exclusion criteria. The sample for this investigation consisted of 238 patients selected using the complete sampling technique. **Results:** The majority of patients who used oral anti-diabetic medications and insulin were between the ages of 46 and 65 (71.34%), were female (73.2%), had standard body mass index (BMIs between 18.5 and 25), and had diagnoses of DM + HT (20.73%) and Type II DM (18.29%). Glulisin was the most commonly used anti-diabetic drug, accounting for 589 DDD/100 days of hospitalization, and DU90% was 21.39%. **Conclusion:** Glulisin is the most frequently prescribed anti-diabetic medication at Bokor Turen Hospital.

Keywords: Antidiabetic Oral and Insulin; Diabetes Mellitus; ATC/DDD; DU 90%

ABSTRAK

Latar Belakang: Resistensi insulin yang berkembang menjadi Diabetes Melitus (DM) erat hubungannya dengan terjadinya kerusakan sel beta pankreas dalam jangka panjang, disfungsi organ, dan kegagalan organ terutama pada mata, ginjal, saraf, jantung, dan pembuluh darah. Pengembangan sistem ATC/DDD diperlukan untuk meningkatkan pengetahuan tentang penggunaan obat di seluruh dunia, memastikan ketersediaan obat secara merata, dan mendorong penggunaan obat yang bijak. Tujuan penelitian ini adalah mengidentifikasi data karakteristik pasien serta obat anti-diabetes oral dan insulin menggunakan metode ATC/DDD dan DU90%. Bahan dan Metode: Rancangan penelitian ini menggunakan *cross-sectional study* yang mengevaluasi obat anti-diabetes oral dan insulin dengan metode ATC/DDD dan DU90% di RS Bala Keselamatan Bokor Turen bulan Januari hingga Desember 2022. Kriteria inklusi terdiri dari pasien DM tipe 1 dan tipe 2 dengan dan atau tanpa komorbid, menggunakan obat anti-diabetes oral dan insulin/kombinasi keduanya. Sedangkan, kriteria eksklusinya berupa data rekam medis tidak lengkap. Populasi penelitian ini sebanyak 238 pasien menggunakan teknik total sampling. Hasil: Data karakteristik pasien pengguna obat anti-diabetes oral dan insulin terbanyak berusia 46 hingga 65 tahun (71,34%); jenis kelamin perempuan (73,2%); Indeks Masa Tubuh normal sebesar 53,66% (IMT= 18,5-25); diagnose DM+HT (20,73%) dan DM Tipe II (18,29%). Penggunaan obat anti-diabetes tertinggi adalah obat glulisin sebesar 589 DDD/100 hari rawat inap dan DU90% sebesar 21,39%. Kesimpulan: Obat anti-diabetes yang paling banyak digunakan di RS Bokor Turen adalah glulisin.

Kata kunci: Obat Anti-diabetes Oral dan Insulin; Diabetes Melitus; DDD; DU 90%

INTRODUCTION

Diabetes Mellitus (DM) is a chronic metabolic disorder. Risk factors, damage to insulin secretion, and sensitivity lead to increased blood glucose and changes in fat and protein metabolism (DiPiro, 2020). The lack of insulin function contributes to the development of microvascular, macrovascular, and neuropathy as a chronic consequence of DM (Almasdy *et al.*, 2015).

The World Health Organization (WHO) in 2023 states that in 2014, 8.5% of adults aged 18 years and over have diabetes. In 2019, diabetes was the direct cause of 1.5 million deaths, and 48% of all deaths from diabetes occurred before the age of 70. Furthermore, 460,000 deaths from kidney disease are caused by diabetes, and elevated blood glucose causes about 20% of deaths from cardiovascular disease (WHO, 2023). DM is more common in low- and moderate-income countries and countries undergoing major economic and demographic transformations. DM is a significant global health problem and requires proper prevention and management measures. DM in Indonesia is currently a serious problem. Most DM sufferers are type 2 DM groups. Therefore, it is essential to evaluate the use of the drug as a basis for selection to ensure that the drug is used appropriately, safely, and efficiently (Pitasari, Andayani, and Wijayanti, 2022).

There is a need to develop ATC/DDD systems to acquire knowledge about drug use worldwide to achieve equitable drug availability and prudent drug use, especially in developing countries. The primary purpose of the ATC/DDD system is to facilitate research on drug use and improve the overall quality of drug use (Tahar et al., 2020). A retrospective study using a *cross-sectional study*, which evaluated drug use patterns and costs associated with Type 2 DM in Saudi Arabia, showed that biguanide (metformin) was most widely prescribed as a monotherapy drug followed by a *fixed-dose combination*. The effectiveness of monotherapy drugs decreases with the duration of treatment; in these cases, combination drugs are prescribed. The most commonly prescribed combination drug is a biguanide with sulfonylurea/biguanide with thiazolidinedione, according to guidelines by the *American Diabetes Association* (ADA). The combination of sitagliptin and metformin is most widely preferred and widely prescribed in *fixed-dose combination therapy*, followed by vildagliptin and metformin (Ali et al., 2022). Combination drugs are used when a single pill cannot achieve the desired blood glucose level in diabetic patients (Okoro, Nmeka, and Erah, 2018).

Evaluation of the use of anti-diabetic drugs with ATC/DDD and DU90% methods can provide insight into the dominant drug use patterns and the extent to which these drugs follow existing treatment recommendations and guidelines. Therefore, this study evaluated oral anti-diabetic drugs and insulin using ATC/DDD and DU90% in DM patients with and without comorbidities at the Bokor Turen Salvation Army Hospital, a hospital with type D accreditation.

MATERIAL AND METHODS

Research Methods

The study design used a *cross-sectional study* with retrospective data collection through patient medical records. Quantitative evaluation of the use of anti-diabetic drugs using ATC / DDD and DU 90%

Commented [MOU1]: Explain further, why this research carried out at this hospital. Author can explain the prevalence Dm in this hospital, or the other reason

techniques. This research has gone through the health research ethics committee with no E.5.a/145/KEPKUMM/V/2023.

Population and Sample

This population consists of patients hospitalized with a diagnosis of DM and without comorbidities at the Bokor Turen Salvation Army Hospital. The study sample included patients hospitalized with a diagnosis of DM and without comorbidities from January to December 2022 and who met the inclusion and exclusion criteria. Inclusion criteria include patients aged ≥ 17 years with a diagnosis of DM and without comorbidities. Exclusion criteria were patients whose medical record data was incomplete.

Sampling Techniques

The sampling technique was total sampling, where the number of samples was equal to a population of 164 patients and met the criteria for inclusion and exclusion of the study.

Data Analysis

The calculation in evaluating the use of ATC / DDD method anti-diabetic drugs in the hospitalization of the Bokor Turen Salvation Army Hospital uses the formula:

$$\frac{\text{DDD}}{100} \text{ bed - days} = \frac{\text{total antibiotics (gram)}}{\text{DDD WHO (gram)}} \times \frac{100}{\text{LOS}}$$

DU 90% is used to identify the amount of drug used as much as 90% of the total use of prescribed medicines and compare it with the amount of residual drug use (RI, 2017). The efficiency of drug use must be observed if the amount of drug use in 10% is more. The DU value of 90% is known after calculating DDD / 100 days of hospitalization per year.

RESULTS

a. Demographic Characteristics of Diabetes Mellitus Patients

The demographic characteristics of Diabetes Mellitus patients at the Bokor Turen Salvation Army Hospital from January to December 2022 were 238 patients in age, gender, BMI, diagnosis, comorbidities, drug names, administration intervals, and duration of administration. This study's sample size was 164 patients. Judging from the most significant number, in the age characteristics of 117 patients aged 46-65, as many as 120 were female, and as many as 88 patients had a regular Body Mass Index (BMI) of 18.5-25 (Table 1).

Commented [MOU2]: Give information that this sample meets the inclusion criteria

Table 1. Data on the Characteristics of Diabetes Mellitus Patients

Characteristic of Patient	Parameter	Total N= (164)	%
Age	17-45	12	7,32
	46-65	117	71,34
	66-95	35	21,34
Mean \pm SD	58,78 \pm 2,52		
Gender	Female	120	73,2
	Male	44	26,8
BMI	Underweight <18,5	6	3,66
	Healthy weight 18,5-25	88	53,66
	Overweight >25	70	42,68
Mean \pm SD	24,63 \pm 1,08		

The description of DM and or without comorbidities at the Bokor Turen Salvation Army Hospital showed that the most patients with a diagnosis of DM and Hypertension (HT) were 34 patients (20.73%), followed by patients with a diagnosis of Type 2 DM as many as 28 patients (18.29%). At the same time, DM patients with other comorbidities have fewer than ten patients (Table 2).

Table 2. Description of Patients with Diabetes Mellitus and or Without Comorbidities

Diagnosis with comorbidities	Number of Patients (n= 164)	%
DM (HT)	34	20,73
DM Type 2	28	18,29
DM (HT, CAD)	6	3,66
DM (anemia)	5	3,05
DM (CAD)	5	3,05
DM (CKD)	5	3,05
DM (CVA)	5	3,05
DM (COPD)	5	3,05
DM (CVA, HT)	4	2,44
DM (anemia, thalassemia)	3	1,83
DM (dyspepsia)	3	1,83
DM (GERD, HT)	3	1,83
DM (HT, anemia)	3	1,83
DM (asthma)	2	1,22
DM (DKD)	2	1,22
DM (DKD, HT)	2	1,22
DM (HT, CKD)	2	1,22
DM (HT, HF)	2	1,22
DM (HT, Vertigo)	2	1,22
DM (STEMI)	2	1,22

CAD= coronary artery disease, CKD= chronic kidney disease, CVA= Cerebrovascular Accident, GERD= Gastroesophageal Reflux Disease, DKD= Diabetic Kidney Disease, HF= Heart Failure

Table 2. Description of Patients with Diabetes Mellitus and or Without Comorbidities (continued)

Diagnosis with comorbidities	Number of Patients (n= 164)	%
DM (Vertigo)	2	1,22
DM (GERD)	1	0,61
DM (Anemia, PAD)	1	0,61
DM (AKI)	1	0,61
DM (Anemia, CKD)	1	0,61
DM (Anemia, Dyspepsia)	1	0,61
DM (Angina, Thalassemia)	1	0,61
DM (CAD, COPD)	1	0,61
DM (CAD, CVA, HT)	1	0,61
DM (CAD, HT)	1	0,61
DM (CAD, STEMI)	1	0,61
DM (CAD, TB)	1	0,61
DM (CKD, HT, TB)	1	0,61
DM (CKD, STEMI)	1	0,61
DM (CKD, Thalassemia)	1	0,61
DM (COPD, Dyspepsia)	1	0,61
DM (COPD, HT)	1	0,61
DM (CVA, susp. COVID-19)	1	0,61
DM (Gastritis)	1	0,61
DM (GERD, HF)	1	0,61
DM (Hepatitis)	1	0,61
DM (HF)	1	0,61
DM (HF, CAD, COPD)	1	0,61
DM (HF, Hepatitis)	1	0,61
DM (HT, HF, dyspepsia)	1	0,61
DM (HT, CA infark)	1	0,61
DM (HT, CAD, GERD)	1	0,61
DM (HT, CAD, Parkinson)	1	0,61
DM (HT, COPD, CVA)	1	0,61
DM (HT, COPD, Dyspepsia)	1	0,61
DM (HT, Dyspepsia)	1	0,61
DM (HT, HF, CAD)	1	0,61
DM (HT, STEMI)	1	0,61
DM (myalgia)	1	0,61
DM (PAD)	1	0,61
DM (PAD, Vertigo)	1	0,61
DM Type 1 (HT)	1	0,61

PAD= Peripheral Artery Disease, AKI= Acute Kidney Injury, COPD= Coronary Obstructive Pulmonary Disease, STEMI= ST-Elevation Myocardial Infarction

Based on the DM treatment profile seen in the 12 months of 2022, it shows that the use of anti-diabetic drugs is the most in the use of glargine drugs with a dose strength of 100U/ml in as many as 71 patients. This was followed by glulisine drugs with a dose strength of 100U / ml for as many as 57 patients. The third highest use was aspart 100U / ml, as many as 54 patients. Concerning the administration interval of most anti-diabetic drugs, they are given every 24 hours a day, with the duration of most drug administration being 4-6 days (Table 3).

Table 3. Profile of Anti-diabetic Drug Use in Year 2022

Characteristic of Patient	Parameter	Number of Patients (N= 293)	%
Type of drug	Acarbose 50 mg	5	1,7
	Glulisine 100 IU/ml	57	19,5
	Glargine 100 IU/ml	71	24,2
	Glibenclamide 5 mg	3	1
	Gliquadone 30 mg	18	6,1
	Glimepiride 2 mg	18	6,1
	Glimepiride 3 mg	18	6,1
	Glimepiride 4 mg	18	6,1
	Lispro 100 IU/ml	7	2,4
	Detemir 100 IU/ml	38	13
	Metformin 500 mg	20	6,8
	Aspart 100 IU/ml	54	18,4
	Pioglitazone 30mg	1	0,3
	Degludec + Aspart 100 IU/ml	1	0,3
Interval of administration	24 hours	157	53,6
	12 hours	21	7,2
	8 hours	114	38,9
	6 hours	1	0,3
Duration of administration	1-3 days	127	43,3
	4-6 days	166	56,7

are often used is glulisine drugs with a dose strength of 100U / ml of 589.02 DDD / 100 days of hospitalization, which means that during 100 days of treatment in the hospital around 589 diabetic patients received glulisine amounting to 40 IU every day. The total number of patients hospitalized and using anti-diabetic drugs from January to December 2022 was 568 days. The second most significant use of anti-diabetic medications is aspart with a dose strength of 100U / ml of 519.72 DDD / 100 days of hospitalization, which means that during 100 days of hospitalization in the hospital around 520 DM patients received aspart drugs of 40 IU (Table 4).

Commented [MOU3]: correct the format

Table 4. Analysis of the Use of Anti-diabetic Drugs using the ATC / DDD Method

No.	ATC code	Type of Drugs	DDD (WHO)	Length of Stay (LOS)	DDD/100 bed-days
1	A10AB06	Glulisine 100 IU/ml	40 IU	568 days	589,02

2	A10AB05	Aspart 100 IU/ml	40 IU	519,72
		Glimepirid 2 mg		474,28
3	A10BB12	Glimepirid 3 mg	2 mg	426,85
		Glimepirid 4 mg		221,38
4	A10AE04	Glargine 100 IU/ml	40 IU	426,85
5	A10AE05	Detemir 100 IU/ml	40 IU	221,38
6	A10BA02	Metformin 500 mg	2 g	198,80
7	A10BB08	Gliquadone 30 mg	60 mg	129,22
8	A10AB04	Lispro 100 IU/ml	40 IU	85,20
9	A10BF01	Acarbose 50 mg	0,3 g	43,55
10	A10BB01	Glibenclamide 5 mg	10 mg	31,24
11	A10BG03	Pioglitazone 30 mg	30 mg	22,72
12	A10AD06	Degludec + aspart 100 IU/ml	40 IU	11,36

Commented [MOU4]:

Commented [MOU5]:

The 90% DU in this study was used to evaluate the use of anti-diabetic drugs in the top 90% of services in the population. There were 12 anti-diabetic assessed drugs in 90% DU and showed that glulycin, aspart, glimepiride 2 mg, glimepiride 3 mg, glimepiride 4 mg, glargine 100U/ml, detemir 100U/ml, and metformin 500 mg were anti-diabetic drugs that accounted for 90% of the highest anti-diabetic drug use in the DM patient population at the Bokor Turen Salvation Army Hospital (Table 5).

Commented [MOU6]: ??

Commented [MOU7]: Milligrams and please correct all this abbreviations in this article, in the table also

Commented [MOU8]:

Commented [MOU9]:

Commented [MOU10]:

Table 5. Profile of Anti-diabetic Drug Use with DU Method 90%

ATC code	Type of Drugs	DDD/100 bed-days	Percentage	Cumulative	DU 90% segment
A10AB06	Glulisine 100 IU/ml	589,02	21,39%	21,39%	
A10AB05	Aspart 100 IU/ml	519,72	18,88%	40,27%	
	Glimepiride 2 mg				
A10BB12	Glimepiride 3 mg	474,28	17,23%	57,49%	DU90%
	Glimepiride 4 mg				
A10AE04	Glargine 100 IU/ml	426,85	15,50%	73,00%	
A10AE05	Detemir 100 IU/ml	221,38	8,04%	81,04%	
A10BA02	Metformin 500 mg	198,80	7,22%	88,26%	
A10BB08	Gliquadone 30 mg	129,22	4,69%	92,95%	
A10AB04	Lispro 100 IU/ml	85,20	3,09%	96,05%	
A10BF01	Acarbose 50 mg	43,55	1,58%	97,63%	
A10BB01	Glibenclamide 5 mg	31,24	1,13%	98,76%	DU 10%
A10BG03	Pioglitazone 30 mg	22,72	0,83%	99,59%	
A10AD06	Degludec+Aspart 100 IU/ml	11,36	0,41%	100%	

Commented [MOU11]:

DISCUSSION

Insulin glulisine (Apidra) in this study occurred in patients with Type 2 diabetes with comorbidities, such as hypertension, coronary artery disease (CAD), and chronic kidney disease (CKD). On average, DM patients who get apidra drugs are given every 8 hours, and this drug is widely presented to 76% of

female patients with an age range of 58-61 years. This insulin is widely recommended at the Bokor Turen Salvation Army Hospital because it can lower blood sugar levels quickly and has a more negligible risk of hypoglycemia. Following the 2021 PERKENI guidelines, Apidra is a rapid-acting insulin generally used with food. Apidra is designed to decrease glucose levels in the blood rapidly after a meal or when blood glucose levels are high. Apidra is commonly combined with basal insulins such as Lantus (insulin glargine) and Levemir (insulin detemir) (Soelistijo et al., 2021). A study examining the clinical effects of Type 2 DM patients with cardiovascular comorbidities where patients used insulin rapid-acting showed that it could have beneficial effects from insulin glulisine administration associated with death and stroke. Still, there was no difference in coronary heart disease (CHD) or cardiovascular disease (CVD) (Svensson et al., 2017).

Insulin aspart (novorapid) is given to patients with type 2 diabetes with comorbid HT accompanied by CAD, ST-Elevation Myocardial Infarction (STEMI), and Heart Failure (HF). The interval of insulin administration averaged every 8 hours per day and was used by 66% of female patients with an average of 58-61 years. Like glulisine insulin, this insulin can also quickly lower blood sugar levels and has a low risk of hypoglycemia. This insulin is also rapid-acting, which provides therapeutic effectiveness after 15 minutes, with the peak of therapeutic efficacy occurring within 1-2 hours and can last up to 4-6 hours. In the case of the population in Japan, administering insulin aspart to type 2 DM patients can significantly reduce cardiovascular complications within 5 to 10 years, resulting in improved quality of life and lower costs compared to *human insulin* (Pollock et al., 2011).

The third most significant use of anti-diabetic drugs is glimepiride doses of 2mg, 3 mg, and 4 mg. This drug is given to patients with type 2 diabetes with cardiovascular comorbidities and an interval of 24 hours per day and primarily female patients (83%) with an average age of 58 years. Regular glimepiride is combined with insulin or another oral medication such as metformin. In the case study of type 2 DM patients with CVD, the average patient suffering from DM was around 5.7 ± 4.8 years. CVD suffered by type 2 DM patients in the form of hypertension (68.5% of patients); dyslipidemia (47.9% of patients); CAD (25.4% of patients); Transient Ischemic Attack (TIA) in 3.6% of patients; peripheral artery disease (PAD) accounted for 4.8% of patients and heart failure in 2.9% of patients. Type 2 DM patients with various comorbidities receive the drug glimepiride/metformin Fixed Dose Combination (FDC) as a first-line therapy because glimepiride/metformin FDC is more effective than metformin alone for type 2 DM patients with comorbidities. As many as 68.2% of FDC patients achieved blood pressure within optimal limits. Most of the other patients experienced an increase in glycemic parameters and a change in body weight of about 18.4%. 59.2% of patients experienced weight loss (Ray et al., 2022). Glimepiride is a

Commented [MOU12]: Past tenses

Commented [MOU13]: Why is insulin glulisine more often choice than aspart?

Commented [MOU14]:

Commented [MOU15]: Why is metformin not among the top three most widely used antidiabetic drug? Wasn't it the first choice?

223 sulfonylurea class drug that has pharmacological effects to increase insulin production by pancreatic
224 beta cells. The most common side effects are hypoglycemia and weight gain.

225
226 The limitation of this study is that data were taken only retrospectively from patients' medical record
227 data, and no direct observations were made on DM patients. For Bala Keselamatan Bokor Turen
228 Hospital to provide an overview of drug use patterns in patients with diabetes mellitus. The results of
229 this study can later be used as a basis for determining the selection of the use of antidiabetic drugs for
230 patients in accordance with applicable standards for patients in accordance with applicable standards.

232 CONCLUSION

233 Based on the analysis of demographic data and treatment profiles of DM patients, DM patients are
234 dominated by the age group of 46-65 years, with women who use anti-diabetic drugs more. Patients with
235 comorbidities use more anti-diabetic medications than patients with non-comorbid DM. Evaluation of
236 anti-diabetic drugs ATC / DDD method shows that insulin glulisine (Apidra) is most used in DM patients
237 with and without comorbidities. Through the DU method, 90% of glulisine, aspart, glargine, glimepiride,
238 detemir, and metformin drugs were most widely used in DM patients.

240 CONFLICT OF INTEREST

241 All authors declare no conflict of interest.

242 REFERENCES

- 243
244 Ali, M.D. et al. (2022) 'Evaluation of drug utilization pattern and cost associated with diabetes
245 mellitus Type two management in Saudi Arabia', *Brazilian Journal of Pharmaceutical Sciences*,
246 58, p. E20681. Available at: <https://doi.org/10.1590/s2175-97902022e20681>.
- 247 Almasdy, D. et al. (2015) 'Evaluation of the Use of Anti-diabetic Drugs in Type-2 Diabetes Mellitus
248 Patients in a Government Hospital of Padang City – West Sumatra', *Journal of Pharmaceutical
249 & Clinical Science*, 2(1), p. 104. Available at: <https://doi.org/10.29208/jsfk.2015.2.1.58>.
- 250 DiPiro, J.T. (ed.) (2020) *Pharmacotherapy: a pathophysiologic approach*. Eleventh edition. New York:
251 McGraw Hill Medical.
- 252 Okoro, R.N., Nmeka, C. and Erah, P.O. (2018) 'Utilization study of antidiabetes medicines at a tertiary
253 care hospital in Nigeria', *Future Journal of Pharmaceutical Sciences*, 4(2), pp. 109–115.
254 Available at: <https://doi.org/10.1016/j.fjps.2017.11.004>.

Commented [MOU16]: Add discussion for the other drugs that include DU90 % : glargine, detemir and metformin. Please compare the result of this research with the previous research.

Commented [MOU17]: The number of references not enough, please add more.

- 255 Pitasari, N.W.N., Andayani, T.M. and Wijayanti, T. (2022) 'Evaluation of the Use of Antidiabetic Drugs
256 in Patients of Back-Referral Program at the Demak District Pharmacy', *Journal of Management
257 and Pharmacy Practice*, 12(2), p. 125. Available at: <https://doi.org/10.22146/jmpf.73841>.
- 258 Pollock, R.F. et al. (2011) 'The cost-effectiveness of rapid-acting insulin aspart compared with human
259 insulin in type 2 diabetes patients: an analysis from the Japanese third-party payer perspective',
260 *Journal of Medical Economics*, 14(1), pp. 36–46. Available at:
261 <https://doi.org/10.3111/13696998.2010.541045>.
- 262 Ray, S. et al. (2022) 'Usage Pattern of Glimepiride/Metformin Fixed-dose Combination in Type 2
263 Diabetes Patients with CVD or at Risk of CVD: An Experience in Indian Setting', *Asian Journal
264 of Diabetology*, 23(2), pp. 13–19.
- 265 RI, K. (2017) 'Technical Guidelines for Evaluation of Drug Use in Health Facilities'.
- 266 Soelistijo, S.A. dkk. (2021) 'Guidelines for the management and prevention of adult type 2 diabetes
267 mellitus in INDONESIA - 2021', *PB. PERKENI*, p. 119.
- 268 Svensson, A.-M. et al. (2017) 'Clinical effects, cardiovascular and renal outcomes associated with rapid-
269 acting insulin analogs among individuals with type 2 diabetes: a nation-wide observational
270 cohort study', *Clinical Diabetes and Endocrinology*, 3(1), p. 5. Available at:
271 <https://doi.org/10.1186/s40842-017-0043-2>.
- 272 Tahar, N.dkk. (2020) 'EVALUATION OF THE USE OF ORAL ANTI-DIABETIC DRUGS IN TYPE
273 2 DIABETES MELLITUS PATIENTS USING ATC/DDD AND DU 90% METHODS', *The
274 2nd Alauddin Pharmaceutical Conference and Expo (ALPHA-C) 2020* [Preprint]. Available at:
275 <https://doi.org/10.24252/kesehatan.v1i1.18380>.
- 276 WHO (2023) *Diabetes*. Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes>
277 (Accessed: September 21, 2023).

9. Keputusan Editor “*Revisions Required*” dan *Submit* revisi artikel tahap ketiga 15 Februari 2024

The screenshot displays the author dashboard for Jurnal Farmasi Galenika. The interface includes a top navigation bar with the journal name, a task counter (0), and user options (English, View Site, godeliva13). The main content area is divided into three sections: Reviewer's Attachments, Revisions, and Review Discussions.

Reviewer's Attachments

Attachment	Date
58259-1, 16624-Article Text-58244-1-4-20240131.docx	February 9, 2024

Revisions

Revision	Date	Article Text
58277-1 Article Text, Revisi anti-diabetic drug.docx	February 15, 2024	Article Text

Review Discussions

Name	From	Last Reply	Replies	Closed
No Items				

The URL at the bottom of the page is: [https://bestjournal.untad.ac.id/index.php/Galenika/\\$?call6\\$\\$/tab/author:dashboard/author:dashboard-review-round-tab/fetch-review-round-info?submissionId=16624&stageId=3&reviewRoundId=11918](https://bestjournal.untad.ac.id/index.php/Galenika/$?call6$$/tab/author:dashboard/author:dashboard-review-round-tab/fetch-review-round-info?submissionId=16624&stageId=3&reviewRoundId=11918)

The screenshot shows a notification email titled "[JFG] Editor Decision" dated 2024-02-15 08:08 AM. The email is addressed to Godeliva Adriani Hendra and contains the following text:

We have reached a decision regarding your submission to Jurnal Farmasi Galenika (Galenika Journal of Pharmacy) (e-Journal), "Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients".

Our decision is: Revisions Required

Please note that you have to submit the revised version of your manuscript as soon as possible with the respond to reviewers comments by answering their comments point by point in a separated file (rebuttal letter).


We will not process the manuscript if the manuscript was not revised as reviewer suggestion or if there is no a rebuttal letter

Thanks and regards

Amelia Rumi
Universitas Tadulako
amelia.rumi@gmail.com

The background of the email shows a sidebar with a list of notifications, including several entries from [JFG] Editor.

10. Keputusan Editor “*Revisions Required*” dan Submit revisi artikel tahap keempat 16 Februari 2024



Lampiran Reviewer Tahap Keempat
16 Februari 2024

Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients

ABSTRACT

Background: Diabetes Mellitus (DM), which results from insulin resistance, is closely linked to long-term damage to pancreatic beta cells, organ dysfunction, and organ failure, particularly in the eyes, kidneys, nerves, heart, and blood vessels. The development of the ATC/DDD system is necessary to increase global drug knowledge, ensure equitable drug availability, and promote responsible drug use. **Objectives:** This study aimed to identify data on patient characteristics and oral anti-diabetic drugs and insulin using the ATC/DDD and DU90% methods. **Material and Methods:** This study was a cross-sectional evaluation of oral anti-diabetic medications and insulin using the ATC/DDD method and DU90% at the Bala Keselamatan Bokor Turen Hospital from January to December 2022. The inclusion criteria were patients diagnosed with type 1 or type 2 diabetes mellitus, with or without comorbidities, who were treated with oral anti-diabetic medications and a combination of insulin and oral medications. In the present study, the incomplete medical record data served as the exclusion criteria. The sample for this investigation consisted of 238 patients selected using the complete sampling technique. **Results:** Most patients who used oral anti-diabetic medications and insulin were between the ages of 46 and 65 (71.34%), were female (73.2%), had standard body mass index (BMIs between 18.5 and 25), and had diagnoses of DM + HT (20.73%) and Type II DM (18.29%). Glulisin was the most used anti-diabetic drug, accounting for 589 DDD/100 days of hospitalization, and DU90% was 21.39%. **Conclusion:** Glulisin is the most frequently prescribed anti-diabetic medication at Bokor Turen Hospital.

Keywords: Antidiabetic Oral and Insulin; Diabetes Mellitus; ATC/DDD; DU 90%

ABSTRAK

Latar Belakang: Resistensi insulin yang berkembang menjadi Diabetes Melitus (DM) erat hubungannya dengan terjadinya kerusakan sel beta pankreas dalam jangka panjang, disfungsi organ, dan kegagalan organ terutama pada mata, ginjal, saraf, jantung, dan pembuluh darah. Pengembangan sistem ATC/DDD diperlukan untuk meningkatkan pengetahuan tentang penggunaan obat di seluruh dunia, memastikan ketersediaan obat secara merata, dan mendorong penggunaan obat yang bijak. Tujuan penelitian ini adalah mengidentifikasi data karakteristik pasien serta obat anti-diabetes oral dan insulin menggunakan metode ATC/DDD dan DU90%. Bahan dan Metode: Rancangan penelitian ini menggunakan *cross-sectional study* yang mengevaluasi obat anti-diabetes oral dan insulin dengan metode ATC/DDD dan DU90% di RS Bala Keselamatan Bokor Turen bulan Januari hingga Desember 2022. Kriteria inklusi terdiri dari pasien DM tipe 1 dan tipe 2 dengan dan atau tanpa komorbid, menggunakan obat anti-diabetes oral dan insulin/kombinasi keduanya. Sedangkan, kriteria eksklusinya berupa data rekam medis tidak lengkap. Populasi penelitian ini sebanyak 238 pasien menggunakan teknik total sampling. Hasil: Data karakteristik pasien pengguna obat anti-diabetes oral dan insulin terbanyak berusia 46 hingga 65 tahun (71,34%); jenis kelamin perempuan (73,2%); Indeks Masa Tubuh normal sebesar 53,66% (IMT= 18,5-25); diagnose DM+HT (20,73%) dan DM Tipe II (18,29%). Penggunaan obat anti-diabetes tertinggi adalah obat glulisin sebesar 589 DDD/100 hari rawat inap dan DU90% sebesar 21,39%. Kesimpulan: Obat anti-diabetes yang paling banyak digunakan di RS Bokor Turen adalah glulisin.

Kata kunci: Obat Anti-diabetes Oral dan Insulin; Diabetes Melitus; DDD; DU 90%

INTRODUCTION

Diabetes Mellitus (DM) is a chronic metabolic disorder. Risk factors, damage to insulin secretion, and sensitivity lead to increased blood glucose and changes in fat and protein metabolism (DiPiro, 2020). The lack of insulin function contributes to the development of microvascular, macrovascular, and neuropathy as a chronic consequence of DM (Almasdy *et al.*, 2015).

The *World Health Organization* (WHO) in 2023 states that in 2014, 8.5% of adults aged 18 years and over have diabetes. In 2019, diabetes was the direct cause of 1.5 million deaths, and 48% of all deaths from diabetes occurred before the age of 70. Furthermore, 460,000 deaths from kidney disease are caused by diabetes, and elevated blood glucose causes about 20% of deaths from cardiovascular disease (WHO, 2023). DM is more common in low- and moderate-income countries and countries undergoing major economic and demographic transformations. DM is a significant global health problem and requires proper prevention and management measures. DM in Indonesia is currently a serious problem. Most DM sufferers are type 2 DM groups. Therefore, it is essential to evaluate the use of the drug as a basis for selection to ensure that the drug is used appropriately, safely, and efficiently (Pitasari, Andayani, and Wijayanti, 2022).

There is a need to develop ATC/DDD systems to acquire knowledge about drug use worldwide to achieve equitable drug availability and prudent drug use, especially in developing countries. The primary purpose of the ATC/DDD system is to facilitate research on drug use and improve the overall quality of drug use (Tahar *et al.*, 2020). A retrospective study using a *cross-sectional study*, which evaluated drug use patterns and costs associated with Type 2 DM in Saudi Arabia, showed that biguanide (metformin) was most widely prescribed as a monotherapy drug followed by a *fixed-dose combination*. The effectiveness of monotherapy drugs decreases with the duration of treatment; in these cases, combination drugs are prescribed. The most commonly prescribed combination drug is a biguanide with sulfonylurea/biguanide with thiazolidinedione, according to guidelines by the *American Diabetes Association* (ADA). The combination of sitagliptin and metformin is most widely preferred and widely prescribed in *fixed-dose combination therapy*, followed by vildagliptin and metformin (Ali *et al.*, 2022). Combination drugs are used when a single pill cannot achieve the desired blood glucose level in diabetic patients (Okoro, Nmeka, and Erah, 2018).

Evaluation of the use of anti-diabetic drugs with ATC/DDD and DU90% methods can provide insight into the dominant drug use patterns and the extent to which these drugs follow existing treatment recommendations and guidelines. Therefore, this study evaluated oral anti-diabetic drugs and insulin using ATC/DDD and DU90% in DM patients with and without comorbidities at the Bokor Turen Salvation Army Hospital, a hospital with type D accreditation. Previous research in the form of case studies analyzing the problem of non-compliance of Diabetes Mellitus patients at Bokor Turen Hospital (Chrisnawati, 2020), so the researcher wants to continue the study on evaluating the use of oral drugs and insulin in the same hospital.

MATERIAL AND METHODS

Research Methods

The study design used a *cross-sectional study* with retrospective data collection through patient medical records. Quantitative evaluation of the use of anti-diabetic drugs using ATC / DDD and DU 90% techniques. This research has gone through the health research ethics committee with no E.5.a/145/KEPKUMM/V/2023.

Population and Sample

This population consists of patients hospitalized with a diagnosis of DM and without comorbidities at the Bokor Turen Salvation Army Hospital. The study sample included patients hospitalized with a diagnosis of DM and without comorbidities from January to December 2022 and who met the inclusion and exclusion criteria. Inclusion criteria include patients aged ≥ 17 years with a diagnosis of DM and without comorbidities. Exclusion criteria were patients whose medical record data was incomplete.

Sampling Techniques

The sampling technique was total sampling, where the number of samples was equal to a population of 164 patients and met the criteria for inclusion and exclusion of the study.

Data Analysis

The calculation in evaluating the use of ATC / DDD method anti-diabetic drugs in the hospitalization of the Bokor Turen Salvation Army Hospital uses the formula:

$$\frac{\text{DDD}}{100} \text{ bed - days} = \frac{\text{total antibiotics (gram)}}{\text{DDD WHO (gram)}} \times \frac{100}{\text{LOS}}$$

DU 90% is used to identify the amount of drug used as much as 90% of the total use of prescribed medicines and compare it with the amount of residual drug use (RI, 2017). The efficiency of drug use must be observed if the amount of drug use in 10% is more. The DU value of 90% is known after calculating DDD / 100 days of hospitalization per year.

RESULTS

a. Demographic Characteristics of Diabetes Mellitus Patients

The demographic characteristics of Diabetes Mellitus patients at the Bokor Turen Salvation Army Hospital from January to December 2022 were 238. This study's sample size was 164 patients in age, gender, BMI, diagnosis, comorbidities, drug names, administration intervals, and duration of

administration. Judging from the most significant number, in the age characteristics of 117 patients aged 46-65, as many as 120 were female, and as many as 88 patients had a regular Body Mass Index (BMI) of 18.5-25 (Table 1).

Table 1. Data on the Characteristics of Diabetes Mellitus Patients

Characteristic of Patient	Parameter	n (%)
Age	17-45	12 (7,32)
	46-65	117 (71,34)
	66-95	35 (21,34)
Mean \pm SD	58,78 \pm 2,52	
Gender	Female	120 (73,2)
	Male	44 (26,8)
BMI	Underweight <18,5	6 (3,66)
	Healthy weight 18,5-25	88 (53,66)
	Overweight >25	70 (42,68)
Mean \pm SD	24,63 \pm 1,08	

The description of DM and or without comorbidities at the Bokor Turen Salvation Army Hospital showed that the most patients with a diagnosis of DM and Hypertension (HT) were 34 patients (20.73%), followed by patients with a diagnosis of Type 2 DM as many as 28 patients (18.29%). At the same time, DM patients with other comorbidities have fewer than ten patients (Table 2).

Table 2. Description of Patients with Diabetes Mellitus and or Without Comorbidities

Diagnosis with comorbidities	n (%)
DM (HT)	34 (20,73)
DM Type 2	28 (18,29)
DM (HT, CAD)	6 (3,66)
DM (anemia)	5 (3,05)
DM (CAD)	5 (3,05)
DM (CKD)	5 (3,05)
DM (CVA)	5 (3,05)
DM (COPD)	5 (3,05)
DM (CVA, HT)	4 (2,44)
DM (anemia, thalassemia)	3 (1,83)
DM (dyspepsia)	3 (1,83)
DM (GERD, HT)	3 (1,83)
DM (HT, anemia)	3 (1,83)
DM (asthma)	2 (1,22)
DM (DKD)	2 (1,22)
DM (DKD, HT)	2 (1,22)
DM (HT, CKD)	2 (1,22)
DM (HT, HF)	2 (1,22)
DM (HT, Vertigo)	2 (1,22)

CAD= coronary artery disease, CKD= chronic kidney disease, CVA= Cerebrovascular Accident, GERD= Gastroesophageal Reflux Disease, DKD= Diabetic Kidney Disease, HF= Heart Failure

134 **Table 2.** Description of Patients with Diabetes Mellitus and or Without Comorbidities (continued)

Diagnosis with comorbidities	n (%)
DM (STEMI)	2 (1,22)
DM (Vertigo)	2 (1,22)
DM (GERD)	1 (0,61)
DM (Anemia, PAD)	1 (0,61)
DM (AKI)	1 (0,61)
DM (Anemia, CKD)	1 (0,61)
DM (Anemia, Dyspepsia)	1 (0,61)
DM (Angina, Thalassemia)	1 (0,61)
DM (CAD, COPD)	1 (0,61)
DM (CAD, CVA, HT)	1 (0,61)
DM (CAD, HT)	1 (0,61)
DM (CAD, STEMI)	1 (0,61)
DM (CAD, TB)	1 (0,61)
DM (CKD, HT, TB)	1 (0,61)
DM (CKD, STEMI)	1 (0,61)
DM (CKD, Thallasemia)	1 (0,61)
DM (COPD, Dyspepsia)	1 (0,61)
DM (COPD, HT)	1 (0,61)
DM (CVA, susp. COVID-19)	1 (0,61)
DM (Gastritis)	1 (0,61)
DM (GERD, HF)	1 (0,61)
DM (Hepatitis)	1 (0,61)
DM (HF)	1 (0,61)
DM (HF, CAD, COPD)	1 (0,61)
DM (HF, Hepatitis)	1 (0,61)
DM (HT, HF, dyspepsia)	1 (0,61)
DM (HT, CA infarct)	1 (0,61)
DM (HT, CAD, GERD)	1 (0,61)
DM (HT, CAD, Parkinson)	1 (0,61)
DM (HT, COPD, CVA)	1 (0,61)
DM (HT, COPD, Dyspepsia)	1 (0,61)
DM (HT, Dyspepsia)	1 (0,61)
DM (HT, HF, CAD)	1 (0,61)
DM (HT, STEMI)	1 (0,61)
DM (myalgia)	1 (0,61)
DM (PAD)	1 (0,61)
DM (PAD, Vertigo)	1 (0,61)
DM Type 1 (HT)	1 (0,61)

135 PAD= Peripheral Artery Disease, AKI= Acute Kidney Injury, COPD= Coronary
 136 Obstructive Pulmonary Disease, STEMI= ST-Elevation Myocardial Infarction
 137

138 The DM treatment profile seen in the 12 months of 2022 shows that anti-diabetic drugs are the most
 139 used in insulin glulisine, with a dose strength of 100U/ml in as many as 71 patients. This was followed
 140 by insulin glulisine with a dose strength of 100U / ml for as many as 57 patients. The third highest use
 141 was insulin aspart 100U / ml, as many as 54 patients. Concerning the administration interval of most
 142 anti-diabetic drugs, they are given every 24 hours a day, with the duration of most drug administration
 143 being 4-6 days (Table 3).

Table 3. Profile of Anti-diabetic Drug Use in Year 2022

Characteristic of Patient	Parameter	n (%)
Type of drug	Acarbose 50 milligrams	5 (1,7)
	Insulin Glulisine 100 IU/ml	57 (19,5)
	Insulin Glargine 100 IU/ml	71 (24,2)
	Glibenclamide 5 milligrams	3 (1)
	Gliquidone 30 milligrams	18 (6,1)
	Glimepiride 2 milligrams	18 (6,1)
	Glimepiride 3 milligrams	18 (6,1)
	Glimepiride 4 milligrams	18 (6,1)
	Insulin Lispro 100 IU/ml	7 (2,4)
	Insulin Detemir 100 IU/ml	38 (13)
	Metformin 500 milligrams	20 (6,8)
	Insulin Aspart 100 IU/ml	54 (18,4)
	Pioglitazone 30 milligrams	1 (0,3)
	Insulin Degludec + insulin Aspart 100 IU/ml	1 (0,3)
Interval of administration	24 hours	157 (53,6)
	12 hours	21 (7,2)
	8 hours	114 (38,9)
	6 hours	1 (0,3)
Duration of administration	1-3 days	127 (43,3)
	4-6 days	166 (56,7)

b. Evaluation of the Use of Anti-diabetic Drugs with ATC/DDD and DU 90% Methods

The results of the evaluation of the use of anti-diabetic drugs using the ATC / DDD method showed that the use of antidiabetics that are often used is insulin glulisine with a dose strength of 100U / ml of 589.02 DDD / 100 days of hospitalization, which means that during 100 days of treatment in the hospital around 589 diabetic patients received insulin glulisine amounting to 40 IU every day. The total number of patients hospitalized and using anti-diabetic drugs from January to December 2022 was 568 days. The second most significant use of anti-diabetic medications was insulin aspart with a dose strength of 100U / ml of 519.72 DDD / 100 days of hospitalization, which means that during 100 days of hospitalization in the hospital, around 520 DM patients received insulin aspart of 40 IU (Table 4).

176

Table 4. Analysis of the Use of Anti-diabetic Drugs using the ATC / DDD Method

No.	ATC code	Type of Drugs	DDD (WHO)	Length of Stay (LOS)	DDD/100 bed-days
1	A10AB06	Insulin Glulisine 100 IU/ml	40 IU	568 days	589,02
2	A10AB05	Insulin Aspart 100 IU/ml	40 IU		519,72
3	A10BB12	Glimepiride 2 milligrams	2 milligrams		474,28
		Glimepiride 3 milligrams			426,85
		Glimepiride 4 milligrams			221,38
4	A10AE04	Insulin Glargine 100 IU/ml	40 IU		426,85
5	A10AE05	Insulin Detemir 100 IU/ml	40 IU		221,38
6	A10BA02	Metformin 500 milligrams	2 grams		198,80
7	A10BB08	Gliquidone 30 milligrams	60 milligrams		129,22
8	A10AB04	Insulin Lispro 100 IU/ml	40 IU		85,20
9	A10BF01	Acarbose 50 milligrams	0,3 grams		43,55
10	A10BB01	Glibenclamide 5 milligrams	10 milligrams		31,24
11	A10BG03	Pioglitazone HCl 30 milligrams	30 milligrams		22,72
12	A10AD06	Insulin Degludec + insulin Aspart 100 IU/ml	40 IU	11,36	

177

178 The 90% DU in this study was used to evaluate the use of anti-diabetic drugs in the top 90% of services
 179 in the population. There were 12 anti-diabetic assessed drugs in 90% DU, and showed that insulin
 180 glulisine, insulin aspart, glimepiride 2 milligrams, glimepiride 3 milligrams, glimepiride 4 milligrams,
 181 insulin glargine 100U/ml, insulin detemir 100U/ml, and metformin 500 milligrams were anti-diabetic
 182 drugs that accounted for 90% of the highest anti-diabetic drug use in the DM patient population at the
 183 Bokor Turen Salvation Army Hospital (Table 5).

184

Table 5. Profile of Anti-diabetic Drug Use with DU Method 90%

ATC code	Type of Drugs	DDD/100 bed-days	Percentage	Cumulative	DU 90% segment
A10AB06	Insulin Glulisine 100 IU/ml	589,02	21,39%	21,39%	DU90%
A10AB05	Insulin Aspart 100 IU/ml	519,72	18,88%	40,27%	
A10BB12	Glimepiride 2 milligrams	474,28	17,23%	57,49%	
	Glimepiride 3 milligrams				
	Glimepiride 4 milligrams				
A10AE04	Insulin Glargine 100 IU/ml	426,85	15,50%	73,00%	
A10AE05	Insulin Detemir 100 IU/ml	221,38	8,04%	81,04%	
A10BA02	Metformin 500 milligrams	198,80	7,22%	88,26%	DU 10%
A10BB08	Gliquidone 30 milligrams	129,22	4,69%	92,95%	
A10AB04	Insulin Lispro 100 IU/ml	85,20	3,09%	96,05%	
A10BF01	Acarbose 50 milligrams	43,55	1,58%	97,63%	
A10BB01	Glibenclamide 5 milligrams	31,24	1,13%	98,76%	
A10BG03	Pioglitazone 30 milligrams	22,72	0,83%	99,59%	

A10AD06	Insulin Degludec + Insulin Aspart 100 IU/ml	11,36	0,41%	100%
---------	---	-------	-------	------

DISCUSSION

Insulin glulisine (Apidra) in this study occurred in patients with Type 2 diabetes with comorbidities, such as hypertension, coronary artery disease (CAD), and chronic kidney disease (CKD). On average, DM patients who get insulin Apidra are given every 8 hours, and this drug is widely presented to 76% of female patients with an age range of 58-61 years. This insulin is widely recommended at the Bokor Turen Salvation Army Hospital because it can lower blood sugar levels quickly and has a more negligible risk of hypoglycemia. Following the 2021 PERKENI guidelines, insulin Apidra is a rapid-acting insulin generally used with food. Insulin Apidra is designed to rapidly decrease glucose levels in the blood after a meal or when blood glucose levels are high. Insulin Apidra is commonly combined with basal insulins such as Lantus (insulin glargine) and Levemir (insulin detemir) (Soelistijo et al., 2021). A study examining the clinical effects of Type 2 DM patients with cardiovascular comorbidities where patients used insulin rapid-acting showed that it could have beneficial effects from insulin glulisine administration associated with death and stroke. Still, there was no difference in coronary heart disease (CHD) or cardiovascular disease (CVD) (Svensson et al., 2017).

Insulin aspart (novorapid) was given to patients with type 2 diabetes with comorbid HT accompanied by CAD, ST-Elevation Myocardial Infarction (STEMI), and Heart Failure (HF). The interval of insulin administration averaged every 8 hours per day and was used by 66% of female patients with an average of 58-61 years. Like insulin glulisine, insulin aspart can also quickly lower blood sugar levels and has a low risk of hypoglycemia. Insulin aspart is also rapid-acting, which provides therapeutic effectiveness after 15 minutes, with the peak of therapeutic efficacy occurring within 1-2 hours and can last up to 4-6 hours. In the case of the population in Japan, administering insulin aspart to type 2 DM patients can significantly reduce cardiovascular complications within 5 to 10 years, resulting in improved quality of life and lower costs compared to *human insulin* (Pollock et al., 2011).

The third most significant use of anti-diabetic drugs is glimepiride doses of 2 milligrams, 3 milligrams, and 4 milligrams. This drug is given to patients with type 2 diabetes with cardiovascular comorbidities and an interval of 24 hours per day and primarily female patients (83%) with an average age of 58 years. Regular glimepiride is combined with insulin or another oral medication such as metformin. Similarly, metformin was used in female patients (81%), with an average age of 58-61 years, a 12-hour drug administration interval and patients suffering from DM with cardiovascular comorbidities. Metformin is usually combined with glimepiride or glibenclamide. In the case study of type 2 DM patients with

CVD, the average patient suffering from DM was around 5.7 ± 4.8 years. CVD suffered by type 2 DM patients in the form of hypertension (68.5% of patients); dyslipidemia (47.9% of patients); CAD (25.4% of patients); Transient Ischemic Attack (TIA) in 3.6% of patients; peripheral artery disease (PAD) accounted for 4.8% of patients and heart failure in 2.9% of patients. Type 2 DM patients with various comorbidities receive the drug glimepiride/metformin Fixed Dose Combination (FDC) as a first-line therapy because glimepiride/metformin FDC is more effective than metformin alone for type 2 DM patients with comorbidities. As many as 68.2% of FDC patients achieved blood pressure within optimal limits. Most of the other patients experienced an increase in glycemic parameters and a change in body weight of about 18.4%. 59.2% of patients experienced weight loss (Ray et al., 2022). Glimepiride is a sulfonylurea class drug that has pharmacological effects to increase insulin production by pancreatic beta cells. The most common side effects are hypoglycemia and weight gain.

Insulin glargine was used in patients with type II DM with cardiovascular comorbidities. The dosage strength used was 100 IU/ml, with a 24-hour drug administration interval. Insulin glargine was mostly used by female patients (71%), and the average age was 58 years. According to a review article, the use of insulin glargine as a basal insulin analog lowers triglycerides causes less weight gain, causes less hypoglycemia when compared to intermediate-acting insulin and has a neutral effect on blood pressure (Joseph & Donner, 2015). According to The Outcome Reduction with Initial Glargine Intervention (ORIGIN trial), the results of the glargine trial showed no increase in cardiovascular risk (Gerstein et al., 2012).

The next anti-diabetic drug in the DU 90% is insulin detemir. In the market, this insulin was known as insulin Levemir (trade name). In this study, insulin detemir was mostly used by 69% of female patients with type II DM with cardiovascular comorbidities. The average age of the patients was 58 years, and the time interval of insulin administration was 24 hours. A systematic review study shows insulin detemir can improve glycaemic control with lower cardiovascular risk and no weight gain in patients with type II DM (Czech et al., 2015).

The limitation of this study is that data were taken only retrospectively from patients' medical record data, and no direct observations were made on DM patients. For Bala Keselamatan Bokor Turen Hospital to provide an overview of drug use patterns in patients with diabetes mellitus. The results of this study can later be used as a basis for determining the selection of anti-diabetic drugs for patients by applicable standards.

CONCLUSION

Based on the analysis of demographic data and treatment profiles of DM patients, DM patients are dominated by the age group of 46-65 years, with women who use anti-diabetic drugs more. Patients with comorbidities use more anti-diabetic medications than patients with non-comorbid DM. Evaluation of anti-diabetic drugs ATC / DDD method shows that insulin glulisine (Apidra) is most used in DM patients with and without comorbidities. Through the DU method, 90% of glulisine, aspart, glargine, glimepiride, detemir, and metformin drugs were most widely used in DM patients.

CONFLICT OF INTEREST


All authors declare no conflict of interest.

REFERENCES

- Ali, M.D., Ahmad A., Banu N., Patel M., Gohsn S.A., I Z.E. (2022) 'Evaluation of drug utilization pattern and cost associated with diabetes mellitus Type two management in Saudi Arabia', *Brazilian Journal of Pharmaceutical Sciences*, 58, p. E20681. Available at: <https://doi.org/10.1590/s2175-97902022e20681>.
- Almasdy, D., Sari D.P., Suhatri S., Darwin D., Kurniasih N. (2015) 'Evaluation of the Use of Anti-diabetic Drugs in Type-2 Diabetes Mellitus Patients in a Government Hospital of Padang City – West Sumatra', *Journal of Pharmaceutical & Clinical Science*, 2(1), p. 104. Available at: <https://doi.org/10.29208/jsfk.2015.2.1.58>.
- Chrisnawati, H. (30 January 2024). Citing Internet sources URL <https://repository.stikespantiwaluya.ac.id/id/eprint/70/>.
- Czech, M., Rdzanek, E., Pawęska, J., Adamowicz-Sidor, O., Niewada, M., & Jakubczyk, M. (2015). Drug-related risk of severe hypoglycaemia in observational studies: A systematic review and meta-analysis. *BMC Endocrine Disorders*, 15(1). <https://doi.org/10.1186/s12902-015-0052-z>
- DiPiro, J.T. (ed.) (2020) *Pharmacotherapy: a pathophysiologic approach*. Eleventh edition. New York: McGraw Hill Medical.
- Gerstein, H., Bosch, J., Dagenais, G. R., Diaz, R., Jung, H., Maggioni, A. P., ... Probstfield, J. (2012). Basal Insulin and Cardiovascular and Other Outcomes in Dysglycemia. *New England Journal of Medicine*, 367(4), 319–328. <https://doi.org/10.1056/NEJMoa1203858>
- Okoro, R.N., Nmeka, C. and Erah, P.O. (2018) 'Utilization study of antidiabetes medicines at a tertiary care hospital in Nigeria', *Future Journal of Pharmaceutical Sciences*, 4(2), pp. 109–115. Available at: <https://doi.org/10.1016/j.fjps.2017.11.004>.
- Joseph, J. J., & Donner, T. W. (2015). Long-term insulin glargine therapy in type 2 diabetes mellitus: A focus on cardiovascular outcomes. *Vascular Health and Risk Management*, 11, 107–116. <https://doi.org/10.2147/VHRM.S50286>

- 288 Pitasari, N.W.N., Andayani, T.M. and Wijayanti, T. (2022) 'Evaluation of the Use of Antidiabetic Drugs
289 in Patients of Back-Referral Program at the Demak District Pharmacy', *Journal of Management
290 and Pharmacy Practice*, 12(2), p. 125. Available at: <https://doi.org/10.22146/jmpf.73841>.
- 291 Pollock, R.F., Valentine W.J., Pilgaard T., Nishimura H. (2011) 'The cost-effectiveness of rapid-acting
292 insulin aspart compared with human insulin in type 2 diabetes patients: an analysis from the
293 Japanese third-party payer perspective', *Journal of Medical Economics*, 14(1), pp. 36–46.
294 Available at: <https://doi.org/10.3111/13696998.2010.541045>.
- 295 Ray, S. et al. (2022) 'Usage Pattern of Glimepiride/Metformin Fixed-dose Combination in Type 2
296 Diabetes Patients with CVD or at Risk of CVD: An Experience in Indian Setting', *Asian Journal
297 of Diabetology*, 23(2), pp. 13–19.
- 298 RI, K. (2017) 'Technical Guidelines for Evaluation of Drug Use in Health Facilities'.
- 299 Soelistijo, S.A. dkk. (2021) 'Guidelines for the management and prevention of adult type 2 diabetes
300 mellitus in INDONESIA - 2021', *PB. PERKENI*, p. 119.
- 301 Svensson, A.-M. et al. (2017) 'Clinical effects, cardiovascular and renal outcomes associated with rapid-
302 acting insulin analogs among individuals with type 2 diabetes: a nation-wide observational
303 cohort study', *Clinical Diabetes and Endocrinology*, 3(1), p. 5. Available at:
304 <https://doi.org/10.1186/s40842-017-0043-2>.
- 305 Tahar, N.dkk. (2020) 'EVALUATION OF THE USE OF ORAL ANTI-DIABETIC DRUGS IN TYPE
306 2 DIABETES MELLITUS PATIENTS USING ATC/DDD AND DU 90% METHODS', *The
307 2nd Alauddin Pharmaceutical Conference and Expo (ALPHA-C) 2020* [Preprint]. Available at:
308 <https://doi.org/10.24252/kesehatan.v1i1.18380>.
- 309 WHO (2023) *Diabetes*. Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes>
310 (Accessed: September 21, 2023).

11. Keputusan Editor “*Revisions Required*” 20 Februari 2024



12. Lampiran Revisi Artikel Tahap Kelima 21 Februari 2024

Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients

ABSTRACT

Background: Diabetes Mellitus (DM), which results from insulin resistance, is closely linked to long-term damage to pancreatic beta cells, organ dysfunction, and organ failure, particularly in the eyes, kidneys, nerves, heart, and blood vessels. The development of the ATC/DDD system is necessary to increase global drug knowledge, ensure equitable drug availability, and promote responsible drug use. **Objectives:** This study aimed to identify data on patient characteristics and oral anti-diabetic drugs and insulin using the ATC/DDD and DU90% methods. **Material and Methods:** This study was a cross-sectional evaluation of oral anti-diabetic medications and insulin using the ATC/DDD method and DU90% at the Bala Keselamatan Bokor Turen Hospital from January to December 2022. The inclusion criteria were patients diagnosed with type 1 or type 2 diabetes mellitus, with or without comorbidities, who were treated with oral anti-diabetic medications and a combination of insulin and oral medications. In the present study, the incomplete medical record data served as the exclusion criteria. The sample for this investigation consisted of 238 patients selected using the complete sampling technique. **Results:** Most patients who used oral anti-diabetic medications and insulin were between the ages of 46 and 65 (71.34%), were female (73.2%), had standard body mass index (BMIs between 18.5 and 25), and had diagnoses of DM + HT (20.73%) and Type II DM (18.29%). Glulisin was the most used anti-diabetic drug, accounting for 589 DDD/100 days of hospitalization, and DU90% was 21.39%. **Conclusion:** Glulisin is the most frequently prescribed anti-diabetic medication at Bokor Turen Hospital.

Keywords: Antidiabetic Oral and Insulin; Diabetes Mellitus; ATC/DDD; DU 90%

ABSTRAK

Latar Belakang: Resistensi insulin yang berkembang menjadi Diabetes Melitus (DM) erat hubungannya dengan terjadinya kerusakan sel beta pankreas dalam jangka panjang, disfungsi organ, dan kegagalan organ terutama pada mata, ginjal, saraf, jantung, dan pembuluh darah. Pengembangan sistem ATC/DDD diperlukan untuk meningkatkan pengetahuan tentang penggunaan obat di seluruh dunia, memastikan ketersediaan obat secara merata, dan mendorong penggunaan obat yang bijak. Tujuan penelitian ini adalah mengidentifikasi data karakteristik pasien serta obat anti-diabetes oral dan insulin menggunakan metode ATC/DDD dan DU90%. Bahan dan Metode: Rancangan penelitian ini menggunakan *cross-sectional study* yang mengevaluasi obat anti-diabetes oral dan insulin dengan metode ATC/DDD dan DU90% di RS Bala Keselamatan Bokor Turen bulan Januari hingga Desember 2022. Kriteria inklusi terdiri dari pasien DM tipe 1 dan tipe 2 dengan dan atau tanpa komorbid, menggunakan obat anti-diabetes oral dan insulin/kombinasi keduanya. Sedangkan, kriteria eksklusinya berupa data rekam medis tidak lengkap. Populasi penelitian ini sebanyak 238 pasien menggunakan teknik total sampling. Hasil: Data karakteristik pasien pengguna obat anti-diabetes oral dan insulin terbanyak berusia 46 hingga 65 tahun (71,34%); jenis kelamin perempuan (73,2%); Indeks Masa Tubuh normal sebesar 53,66% (IMT= 18,5-25); diagnose DM+HT (20,73%) dan DM Tipe II (18,29%). Penggunaan obat anti-diabetes tertinggi adalah obat glulisin sebesar 589 DDD/100 hari rawat inap dan DU90% sebesar 21,39%. Kesimpulan: Obat anti-diabetes yang paling banyak digunakan di RS Bokor Turen adalah glulisin.

Kata kunci: Obat Anti-diabetes Oral dan Insulin; Diabetes Melitus; DDD; DU 90%

INTRODUCTION

Diabetes Mellitus (DM) is a chronic metabolic disorder. Risk factors, damage to insulin secretion, and sensitivity lead to increased blood glucose and changes in fat and protein metabolism (DiPiro, 2020). The lack of insulin function contributes to the development of microvascular, macrovascular, and neuropathy as a chronic consequence of DM (Almasdy *et al.*, 2015).

The *World Health Organization* (WHO) in 2023 states that in 2014, 8.5% of adults aged 18 years and over have diabetes. In 2019, diabetes was the direct cause of 1.5 million deaths, and 48% of all deaths from diabetes occurred before the age of 70. Furthermore, 460,000 deaths from kidney disease are caused by diabetes, and elevated blood glucose causes about 20% of deaths from cardiovascular disease (WHO, 2023). DM is more common in low- and moderate-income countries and countries undergoing major economic and demographic transformations. DM is a significant global health problem and requires proper prevention and management measures. DM in Indonesia is currently a serious problem. Most DM sufferers are type 2 DM groups. Therefore, it is essential to evaluate the use of the drug as a basis for selection to ensure that the drug is used appropriately, safely, and efficiently (Pitasari, Andayani, and Wijayanti, 2022).

There is a need to develop ATC/DDD systems to acquire knowledge about drug use worldwide to achieve equitable drug availability and prudent drug use, especially in developing countries. The primary purpose of the ATC/DDD system is to facilitate research on drug use and improve the overall quality of drug use (Tahar *et al.*, 2020). A retrospective study using a *cross-sectional study*, which evaluated drug use patterns and costs associated with Type 2 DM in Saudi Arabia, showed that biguanide (metformin) was most widely prescribed as a monotherapy drug followed by a *fixed-dose combination*. The effectiveness of monotherapy drugs decreases with the duration of treatment; in these cases, combination drugs are prescribed. The most commonly prescribed combination drug is a biguanide with sulfonylurea/biguanide with thiazolidinedione, according to guidelines by the *American Diabetes Association* (ADA). The combination of sitagliptin and metformin is most widely preferred and widely prescribed in *fixed-dose combination therapy*, followed by vildagliptin and metformin (Ali *et al.*, 2022). Combination drugs are used when a single pill cannot achieve the desired blood glucose level in diabetic patients (Okoro, Nmeka, and Erah, 2018).

Evaluation of the use of anti-diabetic drugs with ATC/DDD and DU90% methods can provide insight into the dominant drug use patterns and the extent to which these drugs follow existing treatment recommendations and guidelines. Therefore, this study evaluated oral anti-diabetic drugs and insulin using ATC/DDD and DU90% in DM patients with and without comorbidities at the Bokor Turen Salvation Army Hospital, a hospital with type D accreditation. Previous research in the form of case studies analyzing the problem of non-compliance of Diabetes Mellitus patients at Bokor Turen Hospital (Chrisnawati, 2020), so the researcher wants to continue the study on evaluating the use of oral drugs and insulin in the same hospital.

MATERIAL AND METHODS

Research Methods

The study design used a *cross-sectional study* with retrospective data collection through patient medical records. Quantitative evaluation of the use of anti-diabetic drugs using ATC / DDD and DU 90% techniques. This research has gone through the health research ethics committee with no E.5.a/145/KEPKUMM/V/2023.

Population and Sample

This population consists of patients hospitalized with a diagnosis of DM and without comorbidities at the Bokor Turen Salvation Army Hospital. The study sample included patients hospitalized with a diagnosis of DM and without comorbidities from January to December 2022 and who met the inclusion and exclusion criteria. Inclusion criteria include patients aged ≥ 17 years with a diagnosis of DM and without comorbidities. Exclusion criteria were patients whose medical record data was incomplete.

Sampling Techniques

The sampling technique was total sampling, where the number of samples was equal to a population of 164 patients and met the criteria for inclusion and exclusion of the study.

Data Analysis

The calculation in evaluating the use of ATC / DDD method anti-diabetic drugs in the hospitalization of the Bokor Turen Salvation Army Hospital uses the formula:

$$\frac{\text{DDD}}{100} \text{ bed - days} = \frac{\text{total antibiotics (gram)}}{\text{DDD WHO (gram)}} \times \frac{100}{\text{LOS}}$$

DU 90% is used to identify the amount of drug used as much as 90% of the total use of prescribed medicines and compare it with the amount of residual drug use (RI, 2017). The efficiency of drug use must be observed if the amount of drug use in 10% is more. The DU value of 90% is known after calculating DDD / 100 days of hospitalization per year.

RESULTS

a. Demographic Characteristics of Diabetes Mellitus Patients

The demographic characteristics of Diabetes Mellitus patients at the Bokor Turen Salvation Army Hospital from January to December 2022 were 238. This study's sample size was 164 patients in age, gender, BMI, diagnosis, comorbidities, drug names, administration intervals, and duration of

administration. Judging from the most significant number, in the age characteristics of 117 patients aged 46-65, as many as 120 were female, and as many as 88 patients had a regular Body Mass Index (BMI) of 18.5-25 (Table 1).

Table 1. Data on the Characteristics of Diabetes Mellitus Patients

Characteristic of Patient	Parameter	n (%)
Age	17-45	12 (7,32)
	46-65	117 (71,34)
	66-95	35 (21,34)
Mean \pm SD	58,78 \pm 2,52	
Gender	Female	120 (73,2)
	Male	44 (26,8)
BMI	Underweight <18,5	6 (3,66)
	Healthy weight 18,5-25	88 (53,66)
	Overweight >25	70 (42,68)
Mean \pm SD	24,63 \pm 1,08	

The description of DM and or without comorbidities at the Bokor Turen Salvation Army Hospital showed that the most patients with a diagnosis of DM and Hypertension (HT) were 34 patients (20.73%), followed by patients with a diagnosis of Type 2 DM as many as 28 patients (18.29%). At the same time, DM patients with other comorbidities have fewer than ten patients (Table 2).

Table 2. Description of Patients with Diabetes Mellitus and or Without Comorbidities

Diagnosis with comorbidities	n (%)
DM (HT)	34 (20,73)
DM Type 2	28 (18,29)
DM (HT, CAD)	6 (3,66)
DM (anemia)	5 (3,05)
DM (CAD)	5 (3,05)
DM (CKD)	5 (3,05)
DM (CVA)	5 (3,05)
DM (COPD)	5 (3,05)
DM (CVA, HT)	4 (2,44)
DM (anemia, thalassemia)	3 (1,83)
DM (dyspepsia)	3 (1,83)
DM (GERD, HT)	3 (1,83)
DM (HT, anemia)	3 (1,83)
DM (asthma)	2 (1,22)
DM (DKD)	2 (1,22)
DM (DKD, HT)	2 (1,22)
DM (HT, CKD)	2 (1,22)
DM (HT, HF)	2 (1,22)
DM (HT, Vertigo)	2 (1,22)

CAD= coronary artery disease, CKD= chronic kidney disease, CVA= Cerebrovascular Accident, GERD= Gastroesophageal Reflux Disease, DKD= Diabetic Kidney Disease, HF= Heart Failure

134 **Table 2.** Description of Patients with Diabetes Mellitus and or Without Comorbidities (continued)

Diagnosis with comorbidities	n (%)
DM (STEMI)	2 (1,22)
DM (Vertigo)	2 (1,22)
DM (GERD)	1 (0,61)
DM (Anemia, PAD)	1 (0,61)
DM (AKI)	1 (0,61)
DM (Anemia, CKD)	1 (0,61)
DM (Anemia, Dyspepsia)	1 (0,61)
DM (Angina, Thalassemia)	1 (0,61)
DM (CAD, COPD)	1 (0,61)
DM (CAD, CVA, HT)	1 (0,61)
DM (CAD, HT)	1 (0,61)
DM (CAD, STEMI)	1 (0,61)
DM (CAD, TB)	1 (0,61)
DM (CKD, HT, TB)	1 (0,61)
DM (CKD, STEMI)	1 (0,61)
DM (CKD, Thallasemia)	1 (0,61)
DM (COPD, Dyspepsia)	1 (0,61)
DM (COPD, HT)	1 (0,61)
DM (CVA, susp. COVID-19)	1 (0,61)
DM (Gastritis)	1 (0,61)
DM (GERD, HF)	1 (0,61)
DM (Hepatitis)	1 (0,61)
DM (HF)	1 (0,61)
DM (HF, CAD, COPD)	1 (0,61)
DM (HF, Hepatitis)	1 (0,61)
DM (HT, HF, dyspepsia)	1 (0,61)
DM (HT, CA infarct)	1 (0,61)
DM (HT, CAD, GERD)	1 (0,61)
DM (HT, CAD, Parkinson)	1 (0,61)
DM (HT, COPD, CVA)	1 (0,61)
DM (HT, COPD, Dyspepsia)	1 (0,61)
DM (HT, Dyspepsia)	1 (0,61)
DM (HT, HF, CAD)	1 (0,61)
DM (HT, STEMI)	1 (0,61)
DM (myalgia)	1 (0,61)
DM (PAD)	1 (0,61)
DM (PAD, Vertigo)	1 (0,61)
DM Type 1 (HT)	1 (0,61)

135 PAD= Peripheral Artery Disease, AKI= Acute Kidney Injury, COPD= Coronary
 136 Obstructive Pulmonary Disease, STEMI= ST-Elevation Myocardial Infarction
 137

138 The DM treatment profile seen in the 12 months of 2022 shows that anti-diabetic drugs are the most
 139 used in insulin glulisine, with a dose strength of 100U/ml in as many as 71 patients. This was followed
 140 by insulin glulisine with a dose strength of 100U / ml for as many as 57 patients. The third highest use
 141 was insulin aspart 100U / ml, as many as 54 patients. Concerning the administration interval of most
 142 anti-diabetic drugs, they are given every 24 hours a day, with the duration of most drug administration
 143 being 4-6 days (Table 3).

Table 3. Profile of Anti-diabetic Drug Use in Year 2022

Characteristic of Patient	Parameter	n (%)
Type of drug	Acarbose 50 milligrams	5 (1,7)
	Insulin Glulisine 100 IU/ml	57 (19,5)
	Insulin Glargine 100 IU/ml	71 (24,2)
	Glibenclamide 5 milligrams	3 (1)
	Gliquidone 30 milligrams	18 (6,1)
	Glimepiride 2 milligrams	18 (6,1)
	Glimepiride 3 milligrams	18 (6,1)
	Glimepiride 4 milligrams	18 (6,1)
	Insulin Lispro 100 IU/ml	7 (2,4)
	Insulin Detemir 100 IU/ml	38 (13)
	Metformin 500 milligrams	20 (6,8)
	Insulin Aspart 100 IU/ml	54 (18,4)
	Pioglitazone 30 milligrams	1 (0,3)
	Insulin Degludec + insulin Aspart 100 IU/ml	1 (0,3)
Interval of administration	24 hours	157 (53,6)
	12 hours	21 (7,2)
	8 hours	114 (38,9)
	6 hours	1 (0,3)
Duration of administration	1-3 days	127 (43,3)
	4-6 days	166 (56,7)

b. Evaluation of the Use of Anti-diabetic Drugs with ATC/DDD and DU 90% Methods

The results of the evaluation of the use of anti-diabetic drugs using the ATC / DDD method showed that the use of antidiabetics that are often used is insulin glulisine with a dose strength of 100U / ml of 589.02 DDD / 100 days of hospitalization, which means that during 100 days of treatment in the hospital around 589 diabetic patients received insulin glulisine amounting to 40 IU every day. The total number of patients hospitalized and using anti-diabetic drugs from January to December 2022 was 568 days. The second most significant use of anti-diabetic medications was insulin aspart with a dose strength of 100U / ml of 519.72 DDD / 100 days of hospitalization, which means that during 100 days of hospitalization in the hospital, around 520 DM patients received insulin aspart of 40 IU (Table 4).

176

Table 4. Analysis of the Use of Anti-diabetic Drugs using the ATC / DDD Method

No.	ATC code	Type of Drugs	DDD (WHO)	Length of Stay (LOS)	DDD/100 bed-days
1	A10AB06	Insulin Glulisine 100 IU/ml	40 IU	568 days	589,02
2	A10AB05	Insulin Aspart 100 IU/ml	40 IU		519,72
3	A10BB12	Glimepiride 2 milligrams	2 milligrams		474,28
		Glimepiride 3 milligrams			426,85
		Glimepiride 4 milligrams			221,38
4	A10AE04	Insulin Glargine 100 IU/ml	40 IU		426,85
5	A10AE05	Insulin Detemir 100 IU/ml	40 IU		221,38
6	A10BA02	Metformin 500 milligrams	2 grams		198,80
7	A10BB08	Gliquidone 30 milligrams	60 milligrams		129,22
8	A10AB04	Insulin Lispro 100 IU/ml	40 IU		85,20
9	A10BF01	Acarbose 50 milligrams	0,3 grams		43,55
10	A10BB01	Glibenclamide 5 milligrams	10 milligrams		31,24
11	A10BG03	Pioglitazone HCl 30 milligrams	30 milligrams		22,72
12	A10AD06	Insulin Degludec + insulin Aspart 100 IU/ml	40 IU	11,36	

177

178 The 90% DU in this study was used to evaluate the use of anti-diabetic drugs in the top 90% of services
 179 in the population. There were 12 anti-diabetic assessed drugs in 90% DU, and showed that insulin
 180 glulisine, insulin aspart, glimepiride 2 milligrams, glimepiride 3 milligrams, glimepiride 4 milligrams,
 181 insulin glargine 100U/ml, insulin detemir 100U/ml, and metformin 500 milligrams were anti-diabetic
 182 drugs that accounted for 90% of the highest anti-diabetic drug use in the DM patient population at the
 183 Bokor Turen Salvation Army Hospital (Table 5).

184

Table 5. Profile of Anti-diabetic Drug Use with DU Method 90%

ATC code	Type of Drugs	DDD/100 bed-days	Percentage	Cumulative	DU 90% segment
A10AB06	Insulin Glulisine 100 IU/ml	589,02	21,39%	21,39%	DU90%
A10AB05	Insulin Aspart 100 IU/ml	519,72	18,88%	40,27%	
A10BB12	Glimepiride 2 milligrams	474,28	17,23%	57,49%	
	Glimepiride 3 milligrams				
	Glimepiride 4 milligrams				
A10AE04	Insulin Glargine 100 IU/ml	426,85	15,50%	73,00%	
A10AE05	Insulin Detemir 100 IU/ml	221,38	8,04%	81,04%	
A10BA02	Metformin 500 milligrams	198,80	7,22%	88,26%	DU 10%
A10BB08	Gliquidone 30 milligrams	129,22	4,69%	92,95%	
A10AB04	Insulin Lispro 100 IU/ml	85,20	3,09%	96,05%	
A10BF01	Acarbose 50 milligrams	43,55	1,58%	97,63%	
A10BB01	Glibenclamide 5 milligrams	31,24	1,13%	98,76%	
A10BG03	Pioglitazone 30 milligrams	22,72	0,83%	99,59%	

A10AD06	Insulin Degludec + Insulin Aspart 100 IU/ml	11,36	0,41%	100%
---------	---	-------	-------	------

DISCUSSION

Insulin glulisine (Apidra) in this study occurred in patients with Type 2 diabetes with comorbidities, such as hypertension, coronary artery disease (CAD), and chronic kidney disease (CKD). On average, DM patients who get insulin Apidra are given every 8 hours, and this drug is widely presented to 76% of female patients with an age range of 58-61 years. This insulin is widely recommended at the Bokor Turen Salvation Army Hospital because it can lower blood sugar levels quickly and has a more negligible risk of hypoglycemia. Following the 2021 PERKENI guidelines, insulin Apidra is a rapid-acting insulin generally used with food. Insulin Apidra is designed to rapidly decrease glucose levels in the blood after a meal or when blood glucose levels are high. Insulin Apidra is commonly combined with basal insulins such as Lantus (insulin glargine) and Levemir (insulin detemir) (Soelistijo et al., 2021). A study examining the clinical effects of Type 2 DM patients with cardiovascular comorbidities where patients used insulin rapid-acting showed that it could have beneficial effects from insulin glulisine administration associated with death and stroke. Still, there was no difference in coronary heart disease (CHD) or cardiovascular disease (CVD) (Svensson et al., 2017).

Insulin aspart (novorapid) was given to patients with type 2 diabetes with comorbid HT accompanied by CAD, ST-Elevation Myocardial Infarction (STEMI), and Heart Failure (HF). The interval of insulin administration averaged every 8 hours per day and was used by 66% of female patients with an average of 58-61 years. Like insulin glulisine, insulin aspart can also quickly lower blood sugar levels and has a low risk of hypoglycemia. Insulin aspart is also rapid-acting, which provides therapeutic effectiveness after 15 minutes, with the peak of therapeutic efficacy occurring within 1-2 hours and can last up to 4-6 hours. In the case of the population in Japan, administering insulin aspart to type 2 DM patients can significantly reduce cardiovascular complications within 5 to 10 years, resulting in improved quality of life and lower costs compared to *human insulin* (Pollock et al., 2011).

The third most significant use of anti-diabetic drugs is glimepiride doses of 2 milligrams, 3 milligrams, and 4 milligrams. This drug is given to patients with type 2 diabetes with cardiovascular comorbidities and an interval of 24 hours per day and primarily female patients (83%) with an average age of 58 years. Regular glimepiride is combined with insulin or another oral medication such as metformin. Similarly, metformin was used in female patients (81%), with an average age of 58-61 years, a 12-hour drug administration interval and patients suffering from DM with cardiovascular comorbidities. Metformin is usually combined with glimepiride or glibenclamide. In the case study of type 2 DM patients with

CVD, the average patient suffering from DM was around 5.7 ± 4.8 years. CVD suffered by type 2 DM patients in the form of hypertension (68.5% of patients); dyslipidemia (47.9% of patients); CAD (25.4% of patients); Transient Ischemic Attack (TIA) in 3.6% of patients; peripheral artery disease (PAD) accounted for 4.8% of patients and heart failure in 2.9% of patients. Type 2 DM patients with various comorbidities receive the drug glimepiride/metformin Fixed Dose Combination (FDC) as a first-line therapy because glimepiride/metformin FDC is more effective than metformin alone for type 2 DM patients with comorbidities. As many as 68.2% of FDC patients achieved blood pressure within optimal limits. Most of the other patients experienced an increase in glycemic parameters and a change in body weight of about 18.4%. 59.2% of patients experienced weight loss (Ray et al., 2022). Glimepiride is a sulfonylurea class drug that has pharmacological effects to increase insulin production by pancreatic beta cells. The most common side effects are hypoglycemia and weight gain.

Insulin glargine was used in patients with type II DM with cardiovascular comorbidities. The dosage strength used was 100 IU/ml, with a 24-hour drug administration interval. Insulin glargine was mostly used by female patients (71%), and the average age was 58 years. According to a review article, the use of insulin glargine as a basal insulin analog lowers triglycerides causes less weight gain, causes less hypoglycemia when compared to intermediate-acting insulin and has a neutral effect on blood pressure (Joseph & Donner, 2015). According to The Outcome Reduction with Initial Glargine Intervention (ORIGIN trial), the results of the glargine trial showed no increase in cardiovascular risk (Gerstein et al., 2012).

The next anti-diabetic drug in the DU 90% is insulin detemir. In the market, this insulin was known as insulin Levemir (trade name). In this study, insulin detemir was mostly used by 69% of female patients with type II DM with cardiovascular comorbidities. The average age of the patients was 58 years, and the time interval of insulin administration was 24 hours. A systematic review study shows insulin detemir can improve glycaemic control with lower cardiovascular risk and no weight gain in patients with type II DM (Czech et al., 2015).

The limitation of this study is that data were taken only retrospectively from patients' medical record data, and no direct observations were made on DM patients. For Bala Keselamatan Bokor Turen Hospital to provide an overview of drug use patterns in patients with diabetes mellitus. The results of this study can later be used as a basis for determining the selection of anti-diabetic drugs for patients by applicable standards.

CONCLUSION

Based on the analysis of demographic data and treatment profiles of DM patients, DM patients are dominated by the age group of 46-65 years, with women who use anti-diabetic drugs more. Patients with comorbidities use more anti-diabetic medications than patients with non-comorbid DM. Evaluation of anti-diabetic drugs ATC / DDD method shows that insulin glulisine (Apidra) is most used in DM patients with and without comorbidities. Through the DU method, 90% of glulisine, aspart, glargine, glimepiride, detemir, and metformin drugs were most widely used in DM patients.

CONFLICT OF INTEREST

All authors declare no conflict of interest.

REFERENCES

- Ali, M.D., Ahmad A., Banu N., Patel M., Gohsn S.A., I Z.E. (2022) 'Evaluation of drug utilization pattern and cost associated with diabetes mellitus Type two management in Saudi Arabia', *Brazilian Journal of Pharmaceutical Sciences*, 58, p. E20681. Available at: <https://doi.org/10.1590/s2175-97902022e20681>.
- Almasdy, D., Sari D.P., Suhatri S., Darwin D., Kurniasih N. (2015) 'Evaluation of the Use of Anti-diabetic Drugs in Type-2 Diabetes Mellitus Patients in a Government Hospital of Padang City – West Sumatra', *Journal of Pharmaceutical & Clinical Science*, 2(1), p. 104. Available at: <https://doi.org/10.29208/jsfk.2015.2.1.58>.
- Chrisnawati, H. (30 January 2024). Citing Internet sources URL <https://repository.stikespantiwaluya.ac.id/id/eprint/70/>.
- Czech, M., Rdzanek, E., Pawęska, J., Adamowicz-Sidor, O., Niewada, M., & Jakubczyk, M. (2015). Drug-related risk of severe hypoglycaemia in observational studies: A systematic review and meta-analysis. *BMC Endocrine Disorders*, 15(1). <https://doi.org/10.1186/s12902-015-0052-z>
- DiPiro, J.T. (ed.) (2020) *Pharmacotherapy: a pathophysiologic approach*. Eleventh edition. New York: McGraw Hill Medical.
- Gerstein, H., Bosch, J., Dagenais, G. R., Diaz, R., Jung, H., Maggioni, A. P., ... Probstfield, J. (2012). Basal Insulin and Cardiovascular and Other Outcomes in Dysglycemia. *New England Journal of Medicine*, 367(4), 319–328. <https://doi.org/10.1056/NEJMoa1203858>
- Okoro, R.N., Nmeka, C. and Erah, P.O. (2018) 'Utilization study of antidiabetes medicines at a tertiary care hospital in Nigeria', *Future Journal of Pharmaceutical Sciences*, 4(2), pp. 109–115. Available at: <https://doi.org/10.1016/j.fjps.2017.11.004>.
- Joseph, J. J., & Donner, T. W. (2015). Long-term insulin glargine therapy in type 2 diabetes mellitus: A focus on cardiovascular outcomes. *Vascular Health and Risk Management*, 11, 107–116. <https://doi.org/10.2147/VHRM.S50286>

- 288 Pitasari, N.W.N., Andayani, T.M. and Wijayanti, T. (2022) 'Evaluation of the Use of Antidiabetic Drugs
289 in Patients of Back-Referral Program at the Demak District Pharmacy', *Journal of Management
290 and Pharmacy Practice*, 12(2), p. 125. Available at: <https://doi.org/10.22146/jmpf.73841>.
- 291 Pollock, R.F., Valentine W.J., Pilgaard T., Nishimura H. (2011) 'The cost-effectiveness of rapid-acting
292 insulin aspart compared with human insulin in type 2 diabetes patients: an analysis from the
293 Japanese third-party payer perspective', *Journal of Medical Economics*, 14(1), pp. 36–46.
294 Available at: <https://doi.org/10.3111/13696998.2010.541045>.
- 295 Ray, S., Ganguly K., Rao M.S., Sinha S., Kumar Y.S., Manchanda S.C., Sharma R.K., Mehta A.,
296 Sawhney JPS., Prasad A., Talathi M. (2022) 'Usage Pattern of Glimepiride/Metformin Fixed-
297 dose Combination in Type 2 Diabetes Patients with CVD or at Risk of CVD: An Experience in
298 Indian Setting', *Asian Journal of Diabetology*, 23(2), pp. 13–19.
- 299 RI, K. (2017) 'Technical Guidelines for Evaluation of Drug Use in Health Facilities'.
- 300 Soelistijo, S.A., Suastika K., Lindarto D., Decroli E., Permana H., Sucipto KW., Kusnadi Y., Budiman.,
301 Ikhsan R., Sasiarini L., Sanusi H., Nugroho HS KH., Susanto H. (2021) 'Guidelines for the
302 management and prevention of adult type 2 diabetes mellitus in INDONESIA - 2021', *PB.
303 PERKENI*, p. 119.
- 304 Svensson, A.-M. Miftaraj M., Franzen S., Eliasson B. (2017) 'Clinical effects, cardiovascular and renal
305 outcomes associated with rapid-acting insulin analogs among individuals with type 2 diabetes:
306 a nation-wide observational cohort study', *Clinical Diabetes and Endocrinology*, 3(1), p. 5.
307 Available at: <https://doi.org/10.1186/s40842-017-0043-2>.
- 308 Tahar, N., Febriyanti A.P., Wahyuddin M., Hasti S.A. (2020) 'EVALUATION OF THE USE OF ORAL
309 ANTI-DIABETIC DRUGS IN TYPE 2 DIABETES MELLITUS PATIENTS USING
310 ATC/DDD AND DU 90% METHODS', *The 2nd Alauddin Pharmaceutical Conference and
311 Expo (ALPHA-C) 2020* [Preprint]. Available at: <https://doi.org/10.24252/kesehatan.v1i1.18380>.
- 312 WHO (2023) *Diabetes*. Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes>
313 (Accessed: September 21, 2023).

13. Copyediting 25 Februari 2024

Jurnal Farmasi Galenika (Galenika Journal of Pharma...)

Tasks 0

English View Site godeliva13

Submission Library View Metadata

Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients
Godeliva Adriani Hendra, Dhanang Prawira Nugraha, Tuswatul Anggi Krisdia

Submission Review Copyediting Production

Copyediting Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
Author Addition in the Article	godeliva13 2024-02-25 01:21 PM	-	0	<input type="checkbox"/>
Publikasi Artikel	saifulm11 2024-03-01 05:41 PM	godeliva13 2024-03-02 01:25 PM	1	<input type="checkbox"/>

Copyedited

Q Search

58360-1	saifulm11, Copyedit_Godeliva.docx	March 1, 2024	Article Text
---------	-----------------------------------	---------------	--------------

https://bestjournal.untad.ac.id/index.php/Galenika/author-dashboards/author-dashboards-tab/fetch-tab?submissionid=16624&stageids=4

14. Copyright transfer agreement 2 Maret 2024

Jurnal Farmasi Galenika (Galenika Journal of Pharma...)

Tasks 0

English View Site godeliva13

Submission Library View Metadata

Evaluation of Anti-diabetic Drugs using ATC/DDD and DU90% Methods in Diabetes Mellitus Patients
Godeliva Adriani Hendra, Dhanang Prawira Nugraha, Tuswatul Anggi Krisdia

Submission Review Copyediting Production

Copyediting Discussions

Add discussion

Name	From	Last Reply	Replies	Closed
Author Addition in the Article	godeliva13 2024-02-25 01:21 PM	-	0	<input type="checkbox"/>
Publikasi Artikel	saifulm11 2024-03-01 05:41 PM	godeliva13 2024-03-02 01:25 PM	1	<input type="checkbox"/>

Copyedited

Q Search

58360-1	saifulm11, Copyedit_Godeliva.docx	March 1, 2024	Article Text
---------	-----------------------------------	---------------	--------------

https://bestjournal.untad.ac.id/index.php/Galenika/author-dashboards/author-dashboards-tab/fetch-tab?submissionid=16624&stageids=4