

International Journal of Food Properties

Country [United States](#) -  [SJR Ranking of United States](#)

Subject Area and Category [Agricultural and Biological Sciences](#)
[Food Science](#)

Publisher [Marcel Dekker Inc.](#)

Publication type Journals

ISSN 15322386, 10942912





Coverage 1998-ongoing





42

H Index

Scope The International Journal of Food Properties publishes original research papers devoted to all scientific and applied aspects of food properties. The emphasis is on measurement methods, development of standards, and data on food properties, predictions, and applications. The International Journal of Food Properties brings together the widely scattered research in the area of food properties and provides an international forum for scientists and technologists for rapid dissemination of their research results, ideas, and knowledge. Other features include review articles, book reviews, letters to the editor, conference papers, news, and commercial advertisements.

-  [Homepage](#)
- [How to publish in this journal](#)
- [Contact](#)
-  [Join the conversation about this journal](#)





International Journal of Food Properties

← Show this widget in your own website

Q2 Food Science
best quartile

SJR 2018 0.49

powered by scimagojr.com

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

Advertisement



Hire the best resea
in **any discipline**

International Journal of Food Properties

Publisher: Taylor & Francis

Journal description

Providing a multidisciplinary, international forum for the rapid dissemination of the latest knowledge, this indispensable new journal offers original research papers devoted to all aspects of food properties from terminology, prediction models, and standardized materials and equipment development to measurement techniques, applications, and future directions. The only periodical focusing entirely on food properties, this stimulating journal considers acoustical, colorimetric, electrical, functional, mass transfer, mass-volume-area-related, mechanical, medical, morphometric, optical, radiative, respiratory, rheological, sensory, surface, thermodynamic, textural, and thermal properties; microbial death-growth-related parameters; and physicochemical constants, in areas such as food product development process and quality control objective texture measurement shelf life prediction process and equipment design marketing and distribution nutritional improvement and much more!

RG Journal Impact: 0.95 *

*This value is calculated using ResearchGate data and is based on average citation counts from work published in this journal. The data used in the calculation may not be exhaustive.


RG Journal impact history

2019	Available summer 2020
2018	0.95
2017	0.90
2016	1.45
2015	1.51
2014	1.41
2013	0.95
2012	1.13
2011	1.35
2010	1.08

2009	0.72
2008	0.86
2007	0.58
2002	1.09
2001	0.25
2000	0.33

RG Journal impact over time

RG Journal impact



Additional details


Cited half-life	4.90
Immediacy index	0.25
Eigenfactor	0.00
Article influence	0.20
Website	http://www.informaworld.com/openurl?genre=journal&issn=10942912
Website description	International Journal of Food Properties website
Other titles	International journal of food properties (Online)
ISSN	1532-2386
OCLC	45274078
Material type	Document, Periodical, Internet resource
Document type	Internet Resource, Computer File, Journal / Magazine / Newspaper

This article was downloaded by: [Leenawaty Limantara]

On: 09 July 2015, At: 17:37

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: 5 Howick Place, London, SW1P 1WG



International Journal of Food Properties

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/ljfp20>

Stability of Palm Carotenes in an Organic Solvent and in a Food Emulsion System

Renny Indrawati^a, Agnieszka Chomiuk^a, Indriatmoko^a, Marcelinus A. S. Adhiwibawa^a, Donald Siahaan^b, Tatas H. P. Brotosudarmo^a & Leenawaty Limantara^a

^a Ma Chung Research Center for Photosynthetic Pigments, Universitas Ma Chung, Malang, East Java, Indonesia

^b Indonesian Oil Palm Research Institute, Medan, North Sumatera, Indonesia

Accepted author version posted online: 09 Apr 2015.



CrossMark

[Click for updates](#)

To cite this article: Renny Indrawati, Agnieszka Chomiuk, Indriatmoko, Marcelinus A. S. Adhiwibawa, Donald Siahaan, Tatas H. P. Brotosudarmo & Leenawaty Limantara (2015) Stability of Palm Carotenes in an Organic Solvent and in a Food Emulsion System, *International Journal of Food Properties*, 18:11, 2539-2548, DOI: [10.1080/10942912.2014.999374](https://doi.org/10.1080/10942912.2014.999374)

To link to this article: <http://dx.doi.org/10.1080/10942912.2014.999374>






PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Stability of Palm Carotenes in an Organic Solvent and in a Food Emulsion System

Renny Indrawati¹ , Agnieszka Chomiuk¹, Indriatmoko¹ ,
Marcelinus A. S. Adhiwibawa¹ , Donald Siahaan², Tatas H. P. Brotosudarmo¹ , and
Leenawaty Limantara¹ 

¹*Ma Chung Research Center for Photosynthetic Pigments, Universitas Ma Chung, Malang, East Java, Indonesia*

²*Indonesian Oil Palm Research Institute, Medan, North Sumatera, Indonesia*

Palm carotene has potential as an application for a natural food colorant with bioactivity as pro-vitamin A and an antioxidant. However, during processing and storage, the palm carotene encounters excessive treatments. In this study, the stability of palm carotenes was observed in organic solvent and a food emulsion system in order to learn its molecular behavior, as well as to evaluate the color stability due to exposure to light and temperature. Spectroscopy and chromatography measurements proved the formation of *cis* isomers and even colorless compounds after the treatment of its acetonic solution at 90°C (11 h) or under excessive illumination (11,470 lux). An application of 0.050–0.500% concentrate of palm carotene into emulsion system gave diverse shades of yellowness, which remained without any obvious color difference ($\Delta E \leq 2$) after 28 days at 4°C (dark) or 8 days at 30°C under average illumination of most display racks at stores (2500 lux).

Keywords: Carotenoid, Color, Emulsification, Pigment, Food stability.

INTRODUCTION

The production of palm oil has undeniably been an economic importance as one of the major export commodities for Indonesia. The Indonesian Directorate General of Estate reported that the increased export volume of palm oil has reached to an average of 13% per year during 2003–2012, and it is accompanied by a 22.24% increase per year on its export value.^[1–3] The statistical data published by FAO has acknowledged Indonesia as the largest producer of the palm oil with a production of 23,672,000 tonnes in 2012, surpassing Malaysia.^[4] The productivity of the plantation continuously grows and was 24,431,640 tonnes by the year 2013.^[5]

The fruit of oil palm tree has also been known to contain high amount of β -carotene which has important biological function as a vitamin A precursor. In Indonesia, *Elaeis guineensis* and *Elaeis*

Received 15 July 2014; accepted 14 December 2014.

Address correspondence to Leenawaty Limantara, Ma Chung Research Center for Photosynthetic Pigments, Universitas Ma Chung, Villa Puncak Tidar N-1, Malang 65151, East Java, Indonesia. E-mail: leenawaty.limantara@machung.ac.id

Color versions of one or more of the figures in the article can be found online at www.tandfonline.com/ljfp.

oleifera are the most cultivated oil palm trees. The mesocarp of the oil palm fruit from *Elaeis guineensis* contains 500–700 ppm carotenoids and even it may reach to 4000 ppm in *Elaeis oleifera*. The α - and β -carotene constitute up to 91% of the total carotenoids (including *cis* and *trans* isomers), while the other carotenoids are small amounts of acyclic carotenes (lycopene, phytofluene, phytoene, neurosporene), cyclic carotenes, as well as lutein.^[6] Nevertheless, prior to produce the commercial cooking oil, the palm oil has to be processed through stages of bleaching and refining. Hence, the final product of cooking oil in the market has a color of light golden yellow and contains only about 27 ppm of carotenoids.^[7]

Due to the plentiful amount of carotenoids, the red palm oil is then used as source of pro-vitamin A. There have been several studies on its effectiveness in elevating vitamin A status of pregnant and lactating women, as well as school-age children.^[8–10] In addition, the consumption of palm oil has also been related to reduction of oxidative tissue damages and prevention of some degenerative diseases.^[11–13] Among many constituents, palm carotenoids and tocopherols were supposed to give main contribution in those activities.

Recently, the present research group has developed a scaled-up process of the carotenoids extraction from red palm esters in order to produce the concentrate.^[14] The carotenes concentrate has a deep yellowish red color and contains less oil fraction. Therefore, it has a potential application for a natural food colorant that has bioactivities such as, pro-vitamin A and antioxidants. The toxicological study has been carried out without any significant effect on vital organs of mice.^[15] However, the stability of the palm carotenes upon their application has not been evaluated. The objective of present study was to evaluate the thermal and photo-stability of palm carotenes when it was dissolved in organic solvent and when it was incorporated in a food emulsion. This study will give reference for further product developments as well as proper storages of palm carotenes.

MATERIALS AND METHODS

The carotenes concentrate of palm oil was obtained from The Indonesian Oil Palm Research Institute (North Sumatra, Indonesia). It was prepared by a modified solvolytic micellization method. The concentrate consisted of up to 80% of ester and was stored in a refrigerator to prevent any degradation before it was used.

Thermal and Photo-Stability in an Organic Solvent

In the thermal stability assay, carotenes concentrate was first dissolved in acetone and adjusted to give OD = 1.0 cm⁻¹ at its maximum absorption (λ_{max} = 447 nm). The solution was placed on a water bath at 90°C for 11 h. The spectral data was recorded using UV-1700 Spectrophotometer (Shimadzu, Japan) subsequently after 1, 2, 3, 6, and 11 h of heating treatment. The photostability assay of palm carotene was determined by treating the equivalent starting solution under Volpi Lamp at 11,470 lux with intense observations using MultiSpec-1501 Spectrophotometer (Shimadzu, Japan).

Chromatography Analysis

The carotenoid composition in the concentrate was determined by high performance liquid chromatography (HPLC) equipped with photo diode array detector (Shimadzu, Japan) using gradient elution as previously described.^[16] The peak area was calculated using the multi-chromatogram approach^[17] at 355–510 nm region.

Emulsion Preparation and Storage Conditions

Oil-in-water emulsion (mayonnaise like) was made by mixing 50% (w/w) canola oil, 4% (w/w) soya lecithin (PT Panadia Corp., Indonesia), 1.25% (w/w) xanthan gum (Qingdao ICD Foreign Trade Co., Ltd., China), and the concentrate of palm carotenes. The water was sterilized prior to emulsification, while the lecithin and xanthan gum were subjected to dry sterilization using ultraviolet lamp. No preservative was added to this emulsion system. The amount of palm carotene was varied as follows: 0.000 (control), 0.050, 0.125, 0.250, and 0.500%, resulting in different shades of yellowness. A commercial hand mixer (Zauberstab M100S, Germany) was used to homogenize the components with the following conditions: 7000 rpm for 1 min and then continued mixing at 10,000 rpm for 2 min. Then, the whole series of emulsions with different concentration of palm carotenes were subjected to three different storage conditions: refrigerated at 4°C (dark) for 28 days, illuminated with 2500 lux monochromatic light at 30°C for 8 days, as well as in a dark oven at 60°C for 8 days. The light intensity of illumination was chosen according to average light intensity of most display racks at food stores.

Color Measurement

ColorFlex[®] EZ No.45/0 (HunterLab, USA) was used to measure emulsion color according to color system of Commission Internationale de l'Eclairage (CIELAB system). It measures L^* for lightness, a^* for redness, and b^* for yellowness. In order to have a valid calculation, the color reading was taken three times and the average value was used. A white tile standard (L^* 92.93, a^* 0.92, b^* 1.48; HunterLab, USA) was used to calibrate the instrument. Change of colors was calculated as ΔE (Eq. 1) from the Hunter L^* , a^* , and b^* values,^[18–20] in which higher values indicate a greater color difference;

$$\Delta E = \sqrt{\frac{1}{3} \left[\left(\frac{L^* - L_0^*}{L_0^*} \right)^2 + \left(\frac{a^* - a_0^*}{a_0^*} \right)^2 + \left(\frac{b^* - b_0^*}{b_0^*} \right)^2 \right]} \quad (1)$$

The subscript “0” refers to the color reading of emulsion prior to storage.

Statistical Analysis

Data were analyzed statistically by one-way analysis of variance (ANOVA) and Scheffé test using SPSS Statistics program, version 16.0. Statistically significance data was accepted at $p < 0.05$ levels.

RESULTS AND DISCUSSION

Pigment Composition

The result of chromatographic analysis of the carotenes concentrate is shown in Fig. 1. The chromatogram was monitored at 450 nm, recognized as maximum absorption of most carotenes. Two dominant peaks were found at 58.21 (peak 1) and 60.10 min (peak 2) which are identified as α - and β -carotene, respectively. Peak identifications were made by retention times and by UV-Vis spectra. Spectral identification was confirmed in *n*-hexane and acetone, as depicted in Fig. 2. The absorption spectrum of α -carotene (β, ϵ -carotene) in *n*-hexane is characterized with three maxima at 422, 442, and 469 nm, whereas β -carotene (β, β -carotene) exhibits a peak-pattern at 424, 447, and 472 nm.^[21,22] Compared to β -carotene, the alpha structural isomer undergoes a hypsochromic shift of 2–5 nm, being similar to that found in geometrical isomerization.^[23] The ration of peak area between α - and β -carotene was 0.804, which provisionally represents their composition in the concentrate. This proportion is comparable to those previously reported for palm oil, red palm oil,




FIGURE 1 HPLC isogram of palm carotenes (a) and the chromatographic profile monitored at 450 nm (b), showing the presence of two major pigments, i.e., α -carotene (peak 1, tR 58.21 min) and β -carotene (peak 2, tR 60.10 min). The photo diode array spectra are portrayed (c) to support the identification.




FIGURE 2 UV-Visible absorption spectra of pigment fractions found as peak 1 at tR 58.21 min (a) and peak 2 at tR 60.10 min (b), collected after chromatography, measured in different solvents: acetone and *n*-hexane.

as well as their olein fraction, having the ratio ranges from 0.627 to 1.327.^[24] According to their biological functions, both α - and β -carotene have shown antioxidant and pro-vitamin A activities. One molecule of β -carotene will be enzymatically cleaved into two molecules of vitamin A (retinol), while cleavage of the α -carotene will generate retinol and α -retinol, having only half the activity than that of the β -carotene.^[25]

Thermal and Photo Stability in an Organic Solvent

Thermal and photo stabilities of carotenes concentrate were evaluated by monitoring the absorption spectra in UV-visible region. The maximum absorption at 447 nm has decreased to 52.35% after 11 h of incubation in a water bath at 90°C. The decrease was also observed to 56.76% after 60 min of irradiation at 11,470 lux. Figure 3a shows that about one-third of the total decrease has taken place in the first hour of heating and afterward it decreased exponentially within the next 10 h. On the other




FIGURE 3 Absorption spectra of carotenes concentrate in acetone during 11 h of heating at 90°C (a) and 60 min of irradiation at 11470 lux (b), as well as the different absorption spectra for thermal (c) and irradiation (d) treatments.

TABLE 1
Degradation percentage of pigment fractions in palm carotenes upon thermal and irradiation treatment, based on peak area data obtained from HPLC analysis

Treatment	Degradation percentage (%)	
	Peak 1	Peak 2
Thermal (90°C, 11 h)	48.48	53.39
Irradiation (11,470 lux, 1 h)	67.24	60.89

hand, exposing palm carotene under irradiation has caused the degradation at constant rate (Fig. 3b). It has been known that excessive exposure of light and/or heat leads to geometrical isomerizations of carotenes and even to a chemical breakdown that cause color change.^[26] In this regard, Fig. 3c shows that an absorption band emerges at the near-ultraviolet region, which is corresponding to the absorption band of the degradation products of carotenes. However, similar situation is not recorded in Fig. 3d that shows the spectral evolution through exposure of light. The new arising band at around 362 nm in Fig. 3a and 3c can be expected as the *cis* band of α - and β -carotene in the *cis* configuration.^[23] Chen and Huang previously identified the degradation and isomerization of all-*trans*- β -carotene in an organic solvent during heating and illumination treatments. Thermo-isomerization leads to the formation of 13-*cis*- β -carotene in largest amount, followed by 9-*cis*- and 15-*cis*- β -carotene, which occurs mostly during the first hour of heating at 70°C.^[27] In the case of an excessive illumination, all-*trans*- β -carotene undergoes geometrical isomerization very quickly in the early phase of photo-degradation, then the *cis* isomers break into colorless compounds. This report is similar with the present results. Unfortunately, the evolution of the absorption band of the isomers is beyond the current resolution of instrument. Table 1 provides degradation percentage of each α - (peak 1) and β -carotene (peak 2) based on peak area obtained from HPLC analysis. The thermal stability of β -carotene was supposed to be slightly lower than that of α -carotene, while the photo stability of α -carotene was suspected to be lesser than that of β -carotene. The degradation pathway of α -carotene as it is affected by high temperature and light still needs to be further studied.

Thermal and Photo Stability in Food Emulsion

The use of palm carotenes in oil-in-water emulsion gave different shades of yellowness at various concentrations (Fig. 4). The L^* , a^* , and b^* values obtained from color measurement are highly correlated with the carotenoid content as well as its chemical structure.^[28] The control emulsion without any colorant gave L^* 88.68, a^* -0.38, and b^* 14.48. When the colorant was added, the lightness parameter (L^* values) was reduced (darker). The chromatic hue a^* (redness) and b^* (yellowness) were greatly varied from a^* 5.827 - b^* 46.04 until a^* 20.403 - b^* 89.48, when 0.050 up to 0.500% carotenes concentrate was incorporated, respectively. It shows the efficacy of this material as a strong colorant. Three different storage conditions were applied based on the possible application of this concentrate, such as ice cream, creamy yogurt, mayonnaise, salad dressings, peanut butter, as well as condiments. Dairy products are usually stored at low temperatures and in the dark for preservation purpose, while the oil-based products are often kept on display racks of the store or kitchen in which there is light exposure at room temperature. Furthermore, dark storage at fairly high temperature (up to 60°C) might be applied inside the container during long distance transportation. A significant statistical shift ($p < 0.05$) of the L^* , a^* , and b^* values over time was observed during storage period in these three different conditions, revealing the existence of color instability. Figure 5 shows the evolution of yellowness (b^* parameter) during certain period of different storage conditions. An adequate linear relation between the b^* parameter and time was



FIGURE 4 Oil-in-water emulsions contain the concentrate of palm carotenes with varied concentrations.



FIGURE 5 Evolution of the yellowness b^* value of oil-in-water emulsion containing palm carotenes during different storage conditions, i.e., dark fridge (T = 40C; 28 days) (a), illumination at 2500 lux (T = 300C; 8 days) (b), and dark oven (T = 600C; 8 days) (c).

TABLE 2
Color differences between the initial condition of food emulsion and those at the end of storage period, when the palm carotenes was applied as natural colorant

Storage condition	Carotene concentrate (% w/w)	ΔE^*
$T = 4^\circ\text{C}$; dark fridge; 28 days	0.000	1.11 ± 0.17
	0.050	0.62 ± 0.24
	0.125	0.82 ± 0.17
	0.250	0.40 ± 0.06
	0.500	1.04 ± 0.18
$T = 30^\circ\text{C}$; illumination at 2500 lux; 8 days	0.000	1.64 ± 0.09
	0.050	2.08 ± 0.11
	0.125	1.82 ± 0.27
	0.250	1.99 ± 0.30
	0.500	1.31 ± 0.21
$T = 60^\circ\text{C}$; dark oven; 8 days	0.000	1.90 ± 0.24
	0.050	7.11 ± 0.22
	0.125	7.52 ± 0.21
	0.250	8.10 ± 0.32
	0.500	7.88 ± 0.14

Data were expressed as the mean \pm SEM;

* ΔE : color difference.

found in low temperature and illuminated storage. During illumination, a negative linear relation between yellowness intensity and time was detected in 0.050 and 0.500% application. Opaque properties of oil-in-water emulsion might help to reduce light penetration, and hence, prevent the degradation pigments.^[29] On the other hand, high-temperature storage for a long time caused lowered redness and yellowness hues in all concentrations with strong negative linear correlation toward time of observation (Fig. 5). The percentage of degradation is retarded on higher concentration. The reduction of absorption intensity due to degradation of *trans* carotene and formation of *cis* isomer become the reason of this discoloration. Color differences were also determined by calculating ΔE to scale the invisibility of any color change (Table 2). The total color difference, ΔE , which is a combination of the parameters (L^* , a^* , and b^* values), is colorimetric parameters extensively used to characterize the variation of colors in foods during processing. The validated value compromises $\Delta E < 1$ for a normally invisible difference, $1 < \Delta E < 2$ for a very small difference that is only obvious to a trained eye, $2 < \Delta E < 3.5$ for medium difference that is obvious to an untrained eye, and $\Delta E > 3.5$ for an obvious difference.^[30] Hence, it can be assumed that at the end of the study, most color emulsions that were stored in a dark fridge as well as under illumination at room temperature did not show a noticeable difference, but those in a dark oven began to give apparent change on the fourth day of storage (data not shown). This prediction was consistent with the characteristic of faster b^* value reduction found during dark oven storage. Overall, there is no statistical significance was found for ΔE among the series of concentration in the same storage condition, suggesting that the concentration of palm carotenes on food emulsion might not influence its color stability.

CONCLUSION

Palm carotenes concentrated from palm oil contained both α - and β -carotene in a nearly equivalent amount. The carotenes concentrate is applicable as a natural yellow colorant, but high-temperature

processing and strong illumination should be minimized due to its instability. The investigation of its molecular behavior by means of chromatography and spectroscopy analysis revealed the formation of degradation products during heating (90°C, 11 h) or illumination with high intensity (11,470 lux, 1 h), i.e., the *cis* isomers and even colorless compounds. In an emulsion system, the color of palm carotenes could remain without any obvious changes during 28 days storage in a dark fridge or 8 days with illumination of 2500 lux at room temperature. However, storage at 60°C for 8 days unveiled apparent color fading. Though it was shown that higher concentration of this colorant has lower percentage of degradation, there is no statistical significance was found for its color difference (ΔE), supposing that the application of concentrate at 0.050–0.500% might not influence its color stability. Further study is now in progress in order to learn the kinetics of color change by using varied light intensities and temperatures prior to application in commercial products.

FUNDING

This research project was supported by Indonesian Directorate General of Higher Education, through the National Strategic research grant (No. 0525/E5.2/PL/2013 and No. 0263/E5/2014).

ORCID

Renny Indrawati  <http://orcid.org/0000-0003-0682-4126>
 Indriatmoko  <http://orcid.org/0000-0001-9537-5398>
 Marcelinus A.S. Adhiwibawa  <http://orcid.org/0000-0002-2191-7411>
 Tatas H.P. Brotosudarmo  <http://orcid.org/0000-0002-8219-3293>
 Leenawaty Limantara  <http://orcid.org/0000-0002-7622-6220>

REFERENCES

1. Nuryanti, S. Nilai Strategis Industri Sawit. Analisis Kebijakan Pertanian **2008**, 6(4), 378–392.
2. Directorate General of Estates. *Production, Area, and Productivity Estate Crops in Indonesia*; Ministry of Agriculture Republic of Indonesia: Jakarta, Indonesia, 2013.
3. Hadi, S.; Tety, E. Analisis Daya Saing Ekspor Minyak Sawit Indonesia Dan Malaysia Di Pasar Internasional. PEKBIS (Jurnal Pendidikan Ekonomi Dan Bisnis) **2012**, 4(3), 180–191.
4. FAOSTAT. *Crops Production*; Food and Agriculture Organization of the United Nations: Quebec, Canada, 2013.
5. Directorate General of Estates. *Productivity Estate Crops 2008–2013*; Ministry of Agriculture Republic of Indonesia: Jakarta, Indonesia, 2014.
6. Ping, B.T.Y. Palm Carotene Concentrates from Crude Palm Oil Using Vacuum Liquid Chromatography on Silica Gel. *Journal of Oil Palm Research* **2007**, 19, 421–427.
7. Hariyadi, P. Lima Alasan Mengapa SNI Minyak Goreng Perlu Revisi (Five reasons why the national standard of frying oil needs to be revised). *Food Review Indonesia*: Bogor, Indonesia, 2013; 26–27.
8. Rice, A.L.; Burns, J.B. Moving from Efficacy to Effectiveness: Red Palm Oil's Role in Preventing Vitamin A Deficiency. *Journal of the American College of Nutrition* **2010**, 29(Sup.3), 302S–313S.
9. Lietz, G.; Mulokozi, G.; Henry, J.C.; Tomkins, A.M. Xanthophyll and Hydrocarbon Carotenoid Patterns Differ in Plasma and Breast Milk of Women Supplemented with Red Palm Oil During Pregnancy and Lactation. *The Journal of Nutrition* **2006**, 136(7), 1821–1827.
10. Zeba, A.N.; Prével, Y.M.; Somé, I.T.; Delisle, H.F. The Positive Impact of Red Palm Oil in School Meals on Vitamin A Status: Study in Burkina Faso. *Nutrition Journal* **2006**, 5(1), 17.
11. Oguntibeju, O.; Esterhuyse, A.; Truter, E. Red Palm Oil: Nutritional, Physiological, and Therapeutic Roles in Improving Human Wellbeing and Quality of Life. *British Journal of Biomedical Science* **2008**, 66(4), 216–222.

12. Oguntibeju, O.; Esterhuysen, A.; Truter, E. Possible Role of Red Palm Oil Supplementation in Reducing Oxidative Stress in HIV/AIDS and TB Patients: A Review. *Journal of Medicinal Plants Research* **2010**, *4*(3), 188–196.
13. Budin, S.B.; Othman, F.; Louis, S.R.; Bakar, M.A.; Das, S.; Mohamed, J. The Effects of Palm Oil Tocotrienol-Rich Fraction Supplementation on Biochemical Parameters, Oxidative Stress, and the Vascular Wall of Streptozotocin-Induced Diabetic Rats. *Clinics* **2009**, *64*(3), 235–244.
14. Rivani, M.P.; Panjaitan, F.R.; Siahaan, D.; Herawan, T.; Hasibuan, H.A. *Scaled Up of Palm Carotenoids Extraction Via Esters Route*, International Oil Palm Conference “Transforming Oil Palm Industry,” Yogyakarta, Indris A.S., L.M., Frank Schuchardt, Chanprasert, Walter Ajambang, J. Ch. Jacquemard, Ed. Indonesian Oil Palm Research Institute: Yogyakarta, 2010; 354–359.
15. Hasibuan, H.A.; Siahaan, D.; Sumantri, I.B. *Acute Toxicity of Palm Carotenoids Concentrate in Mice*, International Oil Palm Conference “Transforming Oil Palm Industry,” Yogyakarta, Indris A.S., L.M., Frank Schuchardt, Chanprasert, Walter Ajambang, J. Ch. Jacquemard, Ed. Indonesian Oil Palm Research Institute: Yogyakarta, 2010; 360–368.
16. Limantara, L.; Heriyanto. Studi Komposisi Pigmen dan Kandungan Fukosantin Rumput Laut Coklat dari Perairan Madura dengan Kromatografi Cair Kinerja Tinggi (Study of pigment composition and fucoxanthin content of brown seaweed in Madura Waters by means of high performance liquid chromatography). *Ilmu Kelautan* **2010**, *15*(1), 23–32.
17. Indrawati, R.; Ozols, M.; Indriatmoko; Heriyanto; Brotosudarmo, T.H.P.; Limantara, L. Re-Evaluation on Multi Chromatogram Approach of 3D-Chromatographic Data. *The Journal of Biomaterial Chemistry* **2013**, *1*(1), 10–14.
18. Gupta, R.K.; Kumar, P.; Sharma, A.; Patil, R.T. Color Kinetics of Aonla Shreds with Amalgamated Blanching During Drying. *International Journal of Food Properties* **2011**, *14*(6), 1232–1240.
19. Zielinska, M.; Markowski, M. Color Characteristics of Carrots: Effect of Drying and Rehydration. *International Journal of Food Properties* **2012**, *15*(2), 450–466.
20. Wrolstad, R.E.; Smith, D.E. Color Analysis. In *Food Analysis*; Nielsen, S.S. Ed.; Springer: New York, NY, **2010**; 573–586.
21. Britton, G.; Liaaen-Jensen, S.; Pfander, H. *Carotenoid. Volume 1B: Spectroscopy*; Birkhäuser Verlag: Basel, Switzerland, 1995.
22. Wright, S.; Jeffrey, S.; Mantoura, R. *Phytoplankton Pigments in Oceanography: Guidelines to Modern Methods*. UNESCO Publishing: Paris, France, 2005.
23. Britton, G.; Liaaen-Jensen, S.; Pfander, H. *Carotenoids: Handbook*. Springer: Basel, Switzerland, 2004.
24. Nagendran, B.; Unnithan, U.; Choo, Y.; Sundram, K. Characteristics of Red Palm Oil, A Carotene and Vitamin E Rich Refined Oil for Food Uses. *Food and Nutrition Bulletin* **2000**, *21*(2), 189–194.
25. Tanumihardjo, S.A.; Howe, J.A. Twice the Amount of α -Carotene Isolated from Carrots is as Effective as β -Carotene in Maintaining the Vitamin A Status of Mongolian Gerbils. *The Journal of Nutrition* **2005**, *135*(11), 2622–2626.
26. Zepka, L.Q.; Borsarelli, C.D.; da Silva, M.A.A.P.; Mercadante, A.Z. Thermal Degradation Kinetics of Carotenoids in a Cashew Apple Juice Model and its Impact on the System Color. *Journal of Agricultural and Food Chemistry* **2009**, *57*(17), 7841–7845.
27. Chen, B.; Huang, J. Degradation and Isomerization of Chlorophyll α and β -Carotene as Affected by Various Heating and Illumination Treatments. *Food Chemistry* **1998**, *62*(3), 299–307.
28. Meléndez-Martínez, A.J.; Britton, G.; Vicario, I.M.; Heredia, F.J. Relationship between the Colour and the Chemical Structure of Carotenoid Pigments. *Food Chemistry* **2007**, *101*(3), 1145–1150.
29. Wang, Z.; Zhang, M.; Wu, Q. Effects of Temperature, pH, and Sunlight Exposure on the Color Stability of Strawberry Juice During Processing and Storage. *LWT-Food Science and Technology* **2015**, *60*(2), 1174–1178.
30. Habekost, M. Which Color Differencing Equation Should be Used? *International Circular of Graphic Education and Research* **2013**, *6*, 20–33.