

Reconstitution Approach to Tune Spectral Features of Light Harvesting Complexes for Improved Light Absorption and Energy Transfer

by Tatas Hardo Panintingjati Brotosudarmo

Submission date: 13-May-2019 11:20AM (UTC+0700)

Submission ID: 1129477992

File name: 17._Energy_Procedia-Reconstitution.pdf (574.56K)

Word count: 5958

Character count: 34123



Conference and Exhibition Indonesia Renewable Energy & Energy Conservation
[Indonesia EBTKE CONEX 2013]

26

Reconstitution Approach to Tune Spectral Features of Light Harvesting Complexes for Improved Light Absorption and Energy Transfer

Heriyanto^{a,b}, Maciej Michalik^b, Tatas H.P. Brotosudarmo^{a,*},
Leenawaty Limantara^a, Leszek Fiedor^{b,†}

9

^aMa Chung Research Center for Photosynthetic Pigments, Ma Chung University, Villa Puncak Tidar N-01, Malang 65151, Indonesia
^bFaculty of Biochemistry, Biophysics and Biotechnology, Jagiellonian University, ul. Gronostajowa 7, 30-387 Krakow, Poland

Abstract

Light harvesting complexes developed by living organisms render themselves as an excellent system for understanding basic physical and chemical processes behind the conversion of sunlight energy. Although light harvesting complexes are pretty robust, biochemical reconstitution and genetic modifications have proven the flexibility to tailor their absorption spectra and energy transfer. Importantly, the refolding of the protein and the exchanging of the pigment in micellar media results in very similar pigment arrangement within the native complexes. Here, we show reconstitution approaches with different pigments that have been carried out in PCP, LHCII, and LHI complexes. Monitoring on the spectral changes and energy transfer has also been described.

© 2014 The Authors. Published by Elsevier Ltd.

Selection and peer-review under responsibility of the Scientific Committee of Indonesia EBTKE Conex 2013

Keywords: reconstitution methods, pigment-protein, light harvesting complex, light absorption, energy transfer

BCHL	Bacteriochlorophyll
β-OG	n-octyl beta-D-glucopyranoside
B780	Monomeric LH1 subunit
B820	Dimeric LH1 Subunit

25

* T.H.P. Brotosudarmo; Tel.: +62-341-550-171; fax: +62-341-550-172; *E-mail address:* tatas.brotosudarmo@machung.ac.id

† L. Fiedor; Tel.: +48—6646358; *E-mail:* leszek.fiedor@uj.edu.pl

24

1876-6102 © 2014 The Authors. Published by Elsevier Ltd.

Selection and peer-review under responsibility of the Scientific Committee of Indonesia EBTKE Conex 2013

[doi:10.1016/j.egypro.2014.01.204](https://doi.org/10.1016/j.egypro.2014.01.204)

23

B870	Carotenoid-less LH1 complex
B880	Fully reconstituted LH1 complex
CD	Circular dichroism
CHL	Chlorophyll
CRT	Carotenoid
DEAE	Diethylaminoethanol
DGDG	Digalactosyldiacylglycerol
iB873	Intermediate complex in the assembly of LH1 complex
LDAO	N,N-dimethyldodecylamine N-oxide
LDS-PAGE	Lithium dodecylsulfate polyacrylamide gel electrophoresis
LH	Light harvesting
LHC	Light harvesting complex
NEU	Neurosporene
NIR	Near infrared
Ni-BCHL	Ni-substituted bacteriochlorophyll a
N-PCP	Apoprotein of N-terminal domain from PCP
PCP	Peridinin chlorophyll protein
RB	Rhodobacter
RC	Reaction center
RSP	Rhodospirillum
SDS	Sodium dodecylsulfate
SPH	Spheroidene

1. Introduction

Sun is one of the most attractive sources of alternative energy. The total energy of sunlight that irradiates the surface of the Earth is believed to provide sufficiently the requirement of a clean and green energy of human population [1]. It will become profitable when a few per cent of this energy can be collected and transformed into some usable energy. Photovoltaic is one of the current technologies that are able to transform sunlight into electricity. The photovoltaic effect was discovered as early as in the first half of the 19th century by Becquerel. Its physical principle, i.e. a light-driven charge separation, resembles the one followed by natural photosynthetic system [2].

Enormous progress in determining the structure of photosynthetic pigment-protein complexes has been carried out using x-ray crystallography to understand the exact position, orientation of the pigments with the resolution ≤ 2.5 Å, well as their binding within the protein matrix [3–5]. In conjunction with spectroscopic data, it reveals how the light harvesting complexes (LHCs) are functioning to capture the energy efficiently and transfer it to the reaction centers (RCs) and how the charge separation occurs in the latter [6]. The structures of LHCs show large variation in terms of the pigment composition depending on the available light spectrum in certain ecological niche in which that photosynthetic organism lives. In the higher plants, chlorophyll (Chl) *a* plays important role in the light harvesting (LH) and other pigments, i.e. Chl *b* and carotenoids (Crts) act as accessory LH pigments. In dinoflagellate, most of the available light spectrum is in the blue-green region of the seawater column. Dinoflagellate have evolved LHCs in which most of the light capturing capacity is provided by Crts, which absorb maximally in the blue-green region of the spectrum. Similar situation occurs in the purple photosynthetic bacteria, which are found in the anaerobic layers of ponds pre-fixed by Chl-based photosynthetic organisms. Thus they use bacteriochlorophyll (BChl) for absorbing the remaining near-infrared (NIR) region of the light spectrum and Crts for absorbing the green light.

LH is the major function of Crt beside its photo-protecting function in photosynthesis process [7]. Crt acts as an accessory pigment to capture light in 400–550 nm spectral region and to transfer the energy to B(Chl). Crt serves as efficient energy donor to B(Chl) in the LH process due to a close contact of Crt with B(Chl) molecules in photosynthetic antenna. Moreover, the fluorescence from the first (S_1) and second (S_2) singlet excited states of Crt

has a good spectral overlap with Qy and Qx bands of B(Chl) which favors energy transfer. Therefore, the efficiency of intra-complex Crt to B(Chl) singlet energy transfer might be affected by densely packed pigments and the Crt structure in LHCs [8]. In dinoflagellate, peridinin is the main Crt and LH pigment in peridinin-chlorophyll-protein (PCP) has a high efficiency of energy transfer to Chl *a*, which approaches to 100%. 80-100% of the overall energy transfer efficiency from Crts to Chls in LHCII might depend on the exact Crt composition in green plant antenna proteins [9–11]. In purple photosynthetic bacteria, the structure of a particular Crt affects its ability to perform energy transfer to BChl *a*. The efficiencies of Crt to BChl energy transfer have been reported to be 80-100% for LH2 of *Rhodobacter sphaeroides* G1C containing mainly neurosporene (Neu) and LH2 of *Rb. sphaeroides* 2.4.1 containing mainly spheroidene (Sph) [12–15], 35-75% for LH2 of *Rhodospirillum acidophila* containing mainly rhodopin glucoside [14–17] and 30% for LH1 of *Rsp. rubrum* containing mainly spirilloxanthin [18].

The utilization of LHCs for applications in different environments triggers ways to modify its structure. Reconstitution of LHCs has been developed to exchange pigments with other types of pigments that have different absorption properties. This approach may be useful in increasing the capacity of LH, e.g. by expanding the range of absorption and enhancing the efficiency of intra-complex energy transfer. The main purpose of this review is to present exemplary reconstitutions and their progress prior to enhance harnessing capacity of solar energy. In this review, we focus on the works that have been done with the PCP from dinoflagellate, the major antenna complexes from purple photosynthetic bacteria (LH1) and from higher plants (LHCII).

2. Reconstitution in peridinin-chlorophyll protein (PCP)

Instead of collecting light mainly by Chls or BChls, the main LH pigment of PCP is peridinin, which absorbs light between 350 and 550 nm. The structure of PCP monomer from *Amphidinium carterae* consists of two Chl *a* molecules and eight peridinin molecules organized in two homologous clusters, which are embedded in protein matrix [4]. The center-to-center distance between Chl *a* is 17.4 Å. The resulting weak coupling between the two Chls and the high Crt-Chl ratio of 4:1 indicate a particular strategy for LH. Upon absorption of light by the peridinin, the energy is then transferred to Chl *a* with a quantum efficiency close to 64% [19, 20]. The energy transfer between the two Chl *a* can be described in terms of Förster energy transfer, with a transfer rate of 0.082 ps⁻¹ [21].

PCP can be reconstituted with Chl *a*, Chl *b*, Chl *d*, acetyl-Chl *a*, and BChl *a* [22–25]. In this way the absorption in the region of NIR can be varied. Importantly, the reconstitution results in very similar pigment arrangement as with the native PCP [25]. PCP can also be reconstituted with Chl mixtures, e.g. Chl *a* and Chl *b* (N-PCP-Chl *a/b*); Chl *a* and acetyl-Chl *a* (N-PCP-Chl *a/acetyl-Chl a*); or Chl *b* and acetyl-Chl *a* (N-PCP-Chl *b/acetyl-Chl a*) [24]. This reconstitution experiment requires apoprotein of N-terminal domain from PCP (N-PCP). It is usually composed of 150 amino acid residues and is topologically a “half-mer” of the monomeric native PCP protomer from *A. carterae* [22]. For the use of reconstitution N-PCP can be prepared by heterologous overexpression in bacteria [22, 26]. The reconstitution procedure was accomplished by mixing apoprotein and ethanol solution of pigments in 1:1.5 molar ratio followed by 48 h of incubation at 4°C [24, 26]. A monomer of PCP from *A. carterae* has one pair of lipid digalactosyldiacylglycerol (DGDG), which binds tightly to pigments [4]. The isolated pigments from native PCP from *A. carterae* already contain DGDG [25]. The use of pigments from different source can be suggested with an addition of DGDG to constitute “high-grade” of PCP for crystallization purpose [25].

The reconstitution procedure yields dimeric aggregates that correspond topologically to native monomer of PCP. The success of reconstitution can be monitored by spectroscopy methods, e.g. absorption, circular dichroism (CD), and fluorescence. Fig. 1 shows the results of reconstitution with different types of the Chl molecule. In these absorption spectra the incorporated Chls give Qy band centered at 670 nm (Chl *a*), 648 nm (Chl *b*), and 651 and 668 nm (Chl *a/b*). The visible region (400-600 nm) of the absorption spectra of PCP reconstituted with Chl *a* is similar to those of native PCP. Reconstitution with Chl *b* yields more pronounced Soret band in the visible region due to the larger extinction coefficient of Chl *b* as compared to Chl *a* [27]. The fluorescence spectrum of N-PCP-Chl *a* shows a single line at 673 nm originating from the S₁-S₀ transition of Chl *a* that has been generated by energy transfer from peridinin through excitation at 532 nm. In N-PCP-Chl *b*, the fluorescence emission is blue shifted to 652 nm. It was calculated that the fluorescence quantum yield of Chl *a* and *b* in N-PCP were 24% and 11%, respectively [28].




Figure 1. The absorption (a) and fluorescence emission (b) spectra of native PCP (black line with shaded area) and the reconstituted complexes: N-PCP Chl *a* (dashed line), N-PCP Chl *b* (dotted line), and N-PCP Chl *a/b* (dash-dot line). The fluorescence emission spectra were taken with excitation wavelength at 532 nm.

The absorption spectra of the reconstituted PCP with Chl mixtures show superficially a mix of the spectra of reconstituted complexes with single type of Chl. The fluorescence spectrum of N-PCP reconstituted with both Chl *a* and *b* is also characterized by two emission lines corresponding to the two Chl pigments (Fig. 1B). The presence of reconstituted PCP containing Chl *a* in the one half and Chl *b* in the other half of dimeric aggregate can be monitored by single molecule spectroscopy [21,9]. The detailed experiment using single molecule spectroscopy can also show Förster energy transfer between Chl *a* and *b* in a monomer. The downhill energy transfer from Chl *b* to Chl *a* took about 31 ps, while the uphill energy transfer took 47 ps. It is also shown that energy transfer from peridinin to Chl *a* is almost three times more efficient than that from peridinin to Chl *b* [28].

3. Reconstitution in core light-harvesting complex II

An *in vitro* assembly of native and recombinant LHCIIb protein with its native chromophores has first been reported by Plumley and Schmidt [30] and Paulsen et al. [31], respectively. These methods rely on the self-assembly of sodium dodecylsulfate (SDS) denatured LHCIIb apoprotein when mixed with the total pigment extract of spinach leaves, followed by several freeze-thaw cycles. The minimum components required for LHCIIb folding were Chl *a/b* and xanthophylls, but not diacylglycerides.

A similar reconstitution technique was employed to assess the structure-function relationships in LHCII formed from its apoprotein overexpressed in bacterial hosts. The deletion mutagenesis helped to elucidate the minimum amino acid requirements for LHCII assembly [32, 33]. Since the atomic model of LHCII provided by Kühlbrandt et al. [34] did not distinguish between Chl *a* and Chl *b*, the mutations of Chl-binding amino acid residues were expected to address the question of assignment of specific Chl species to their binding sites in the antenna [35, 36]. Surprisingly, partially inconsistent results have been obtained, leading the authors to the conclusion that some binding sites in LHCIIb are non-specific. This observation has been confirmed by Kleima et al. [37], when a wild-type LHCIIb was reconstituted with different Chl *a/b* ratios, ranging 1 Chl *a* per 35 Chl *b* molecules in LHCII trimer. Analysis of the results showed a large degree of promiscuity among most Chl *a* binding sites, which can be utilized to introduce Chls other than the native ones.

During over 25 years period, the protocol for LHCII reconstitution has been substantially improved. It relies on spontaneous refolding of SDS denatured LHCIIb in the presence of pigment dissolved in ethanol. More recently [38], the detergent exchange was achieved via mixed micelles formed in the presence of *n*-octyl beta-D-glucopyranoside (β -OG), and SDS is precipitated out at 4°C with potassium ions. After centrifugation, the supernatant yields monomeric LHCII and pigments solubilized in the micelles. The latter can be efficiently separated from the complexes via partially denaturing polyacrylamide gel electrophoresis in the presence of lithium

dodecylsulfate (LDS-PAGE) [30–33, 36] or a combination of chromatographic and density gradient centrifugation methods [35, 37–39]. The assessment of preparation quality is based on the electronic absorption, CD and fluorescence emission measurements, which can be performed in excised green gel bands or after complex elution from the gel. Spectral properties of native and reconstituted LHCII, as presented in Fig. 2, are fairly similar [31, 38, 40]. The differences seen between two types of complexes, particularly in the CD spectra, can be explained by specific interactions between chromophores of individual LHCII monomers in the native antenna. In turn, the discrepancy in the Chl absorption intensities originates from altered Chl *a/b* ratio in the reconstituted LHCII (not shown), caused probably by promiscuity of Chl *a* binding sites [37].

LHCII reconstitution served to study the kinetics of antenna folding, thermodynamics of cofactor binding, and excitation transfer dynamics, but in contrast to LH1, very few reports can be found on pigment exchange in LHCII. Some modification of spectral properties has been introduced, e.g. by means of covalent attachment of rhodamine [11] to LHCIIb apoprotein, which did not interfere with complex assembly [11] functioning [39]. The efficient excitation energy transfer from the rhodamine molecules to Chls resulted in an increase in the absorption cross section and overall LH efficiency, but such an approach can not be considered as pigment exchange. Thus, LHCII and LH2 remain the antenna complexes whose pigment content and spectral features have not been modified by means of reconstitution. This fact is noteworthy, especially considering a large number of reports on other aspects of LHCII reconstitution, in contrast to LH2. The methods for pigment exchange in this complex are yet to be worked out.




Figure 2. Electronic absorption and circular dichroism spectra of native trimeric (dotted line) and reconstituted monomeric (solid line) forms of LHCII. The samples were obtained according to [38], purified by LDS-PAGE method and eluted from the gel prior to the measurement.

4. Reconstitution in core light-harvesting I complex

The LH1 complex is the simplest type of photosynthetic antenna complexes which can now be isolated and is well characterized. In addition, several methods exist for its *in vitro* reconstitution from the individual components, which also provide means for pigment exchange [41, 42]. Reconstitution of LH1 proved a highly successful approach to understand the structure-function relationships in this antenna [43]. According to the structural modeling and low resolution crystallographic data [44–46], the reaction center complex (RC) fits into the inner space of the LH1 complex. The two complexes closely interact to ensure a high efficiency of inter-complex energy transfer. LH1 is composed of 15–16 heterodimeric subunits of highly hydrophobic α - and β -polypeptides, which host a circular array of 30–32 strongly interacting BChl molecules and 15–16 Crt molecules. Both BChl and Crts are involved in the LH process. BChls are in contact with Crts via van der Waals interaction with the long alcohol residues esterifying the C-173 propionic acid side group [47].

The LH1 reconstitution was initially introduced by Paul A. Loach and the co-workers, taking advantage of the fact that LH1, in micellar media and in the absence of Crts, reversibly dissociates into dimeric B820 subunits and further to its monomeric B780 subunits. Prior to dissociation, Crts must be removed from LH1, either at the stage of

chromatophores of wild-type photosynthetic bacteria, e.g. by extraction with a hydrocarbon solvent, or using carotenoidless strains of bacteria. The Crt-depleted LH1 has the maximum absorption wavelength (λ_{max}) at around 870 nm, and is termed B870. On the basis of biochemical and spectroscopic properties, the composition of B820 subunit is assigned to $\alpha\beta$ -BChl. Based on these properties, Paul A. Loach and the co-workers have used the native and modified BChls and the polypeptides for reconstituting the B820 subunits, B870 and LH1 complexes to reveal the structural requirements for their formation [48, 49]. These reconstitutions were performed on analytical scale and no attempts were made to isolate the reconstitution products.

The LH1 reconstitution approach has recently been further elaborated by Leszek Fiedor and the co-workers. The protocols developed provided means to obtain large quantities of the reconstituted pigment-protein complex. To this end, three different reconstitution strategies have been developed: (1) LH1 formation from organic solvent-extracted LH1 components, (2) LH1 formation from detergent-dissociated Crt-depleted antenna and (3) LH1 formation via oligomerization of the B820 subunit.

4.1 Reconstitution from organic solvent-extracted LH1 components

The components of LH1 from freeze-dried membrane of an LH1-only mutant of *Rb. sphaeroides* (DD13 strain) were extracted with a mixture of chloroform and methanol containing ammonium acetate [50, 51]. The extract, which contains pigments and polypeptides of the LH1 complex, is dried under vacuum. The resulted solid form was suspended in Tris-HCl buffer, followed by solubilization at high concentrations of β -OG (20% or 3.4%). The progress of reconstitution can be monitored as an increase in the absorption at 875 nm. The reconstituted LH1 complex is nearly identical to the native one in terms of absorption, fluorescence and CD spectra as well as energy transfer efficiency from Crt to BChl. Ion-exchange chromatography on a diethylaminoethanol (DEAE)-Sephacrose column was used for purification of this reconstituted LH1 complex. Generally, the higher recovery of the reconstituted LH1 complex, up to 40%, is achieved in reconstitution with the lower detergent concentration (3.4%). This method allows for insertion of Ni-substituted analog of BChl (Ni-BChl) a to the LH1 complex, when this pigment is added at the stage of organic solvent extract.

4.2 Reconstitution from Crt-depleted antenna

A convenient way for reconstituting LH1 complex either from the Crt-depleted antenna of *Rsp. rubrum* or the Crt-less antenna of *Rsp. rubrum* G9, which facilitates Crt insertion, has been introduced by Fiedor et al. [42]. This reconstitution method can be applied for the insertion of modified BChl into LH1. The Crt-depleted antenna is prepared either from the freeze-dried chromatophores of *Rsp. rubrum* S1 after removal of Crts by benzene extraction or of the Crt-less strain *Rsp. rubrum* G9 [52]. The Crt-less antenna is solubilized in 0.03% N,N-dimethyldodecylamine N-oxide (LDAO) to achieve its dissociation into a mixture of the B780 and B820 subunits. Upon the addition of acetone, the equilibrium between the LH1 subunits shifts towards B870, due to an increase in dielectric permittivity of the medium [53]. The fully reconstituted LH1 complex (B880) can be formed in the presence of Crt added as acetone solution. The position of the Q_y band of BChl a in this complex is red shifted by 10-12 nm compared to that in the Crt-less LH1. A similar shift is observed in the absorption of Crts. Most likely, excitonic coupling between BChl a molecules in LH1 is enhanced due to the Crts. Indeed, thermodynamic analysis of LH1 formation indicates strong interactions of Crts with the LH1 subunits [53]. Preparative purification of the reconstituted LH1-Crt complexes was performed by ion-exchange chromatography. If necessary, a sucrose density gradient centrifugation step can be added to remove contaminating Crt aggregates [54]. The yields of the purified complexes with respect to the Crt-depleted antenna vary between 40% and 75%, depending on the stability of the complex [42, 54]. In order to introduce a modified BChl into LH1, e.g. Ni-BChl a, the addition of the pigment in acetone to the reconstitution mixture is done before the addition of Crt [55].

The spectral features of the Crt-less antenna and the LH1 complexes reconstituted with two Crts, Neu and Sph (Fig. 3), are in agreement with those reported previously [42, 53]. The Q_y band of BChl of the Crt-less antenna has λ_{max} at 870 nm, while this band of the reconstituted LH1-Crt complexes is typically red shifted to 880 nm due to the correct assembly of LH1 complex induced by Crt. Absorption spectra of the LH1 complexes reconstituted with Crts show that Crt absorption can cover the energy gap in blue-green region of BChl absorption spectrum to enhance the

efficiency of photon capturing. CD spectra of LH1 complex reconstituted with Crt show a specific feature of Crt absorption due to Crt being placed in the chiral environment of LH1 polypeptides. These results indicate the correct incorporation of Crt into the complex, which is also confirmed by a high efficiency of the intra-complex (Crt to BChl) singlet energy transfer. The efficiency of this energy transfer decreases gradually from 80% to 30% along with increasing the number of conjugated double bonds (n) from 9 to 13. It has been shown that as the n in Crt increases above 11 not all low-lying singlet states of Crt are available as channels for energy transfer to BChl a [52, 54].




Figure 3. Electronic absorption and circular dichroism spectra of Crt-less antenna (B870, solid line) in 0.4% β -OG and the LH1 complexes reconstituted with neurosporene (LH1-Neu; dashed line) and spheroidene (LH1-Sph; dotted line) in 0.03% and 0.045% LDAO, respectively.

4.3 Reconstitution from the B820 subunit

The B820 subunit was prepared from freeze-dried chromatophores of the Crt-less antenna of *Rsp. rubrum* by solubilization in 3.4% β -OG and purified on a DEAE-Sepharose column. The yield of pure B820 reaches 70-80% with respect to the initial amount of the chromatophores. LH1 complex can be reconstituted with the incorporation of Crt into B820 subunits in 1% β -OG. The progress of LH1 formation can be monitored as described above. This approach facilitated the observation of a new short-lived intermediate in the assembly of LH1 termed iB873, which can be converted to the LH1 complex if more Crt is applied [56]. Also this reconstitution method allows the replacement of the native BChl *a* with the modified pigments, such as Ni-BChl *a*, simply by adding its solution (in acetone) into the B820 subunits in 1% β -OG, before the oligomerization of the subunits is induced [55].

5. Summary

The fully reconstituted PCP requires peridinin as the main Crt and apoprotein of N-terminal domain as well as Chl. PCP reconstitution with the modified Chl *a*, i.e. its analogs, can enhance the light absorption in the NIR region. Although energy transfer from peridinin to Chl *a* is more efficient compared to that one from peridinin to its analogs, i.e. Chl *b*. Importantly, the energy transfer from peridinin to Chl *a* is close to 100% upon the absorption of light by peridinin. Three reconstitution strategies were developed for reconstituting LH1 in large quantities by use of organic solvent-extracted LH1 components, detergent-dissociated Crt-depleted antenna, and oligomerization of B820 subunit. These reconstitution approaches manipulate independently its components, especially Crts, for enhancing the light absorption by LH1 complex in the blue-green region which is not covered by BChl, and in the NIR region. Therefore, Crt acts as one of the structural requirements in LH1 complex. Intra-complex energy transfer from Crt to BChl can be varied between 30% to 80%, and depends on the n of Crts incorporated to the reconstituted LH1 complexes. By using these LH1 complexes reconstituted with Crts, antioxidant and photo-protective functions of Crt can be studied. Crts may significantly enhance the stability of photosynthetic complexes against oxidation and their protective effect depends on the type of the oxidant [57]. Photo-protective function of Crt, which quenches the

lowest triplet state of BChl *a* and dissipates the transferred triplet energy as heat, is more efficient in the LH1 rather than in the LH2 complex [58]. The generation of the Crt triplet state increases accordingly with the number of conjugated double bonds of incorporated Crt [52].

Acknowledgements

THPB acknowledges financial support from the Indonesian Ministry of Research and Technology through National Innovation System Research Grant (RT-2013-166, No:187/M/Kp/XI/2012) and International Collaboration Research Grant from Directorate General for Higher Education (DIKTI No. 088/SP2H/PL/DIT.LITABMAS/V/2013). This work was in part supported by a research grant from the Foundation for Polish Science (TEAM/2010-5/3 to L.F.)

References

- [1] MacKay DJC. Solar. In: Sustainable energy—without the hot air. Cambridge: UIT Cambridge; 2009. p. 38-49.
- [2] Landsberg PT, Markvart T. The car not factor in solar-cell theory. Solid State Electron 1998; 42:657-59.
- [3] Koepke J, Hu X, Muenke C, Schulten K, Michel H. The crystal structure of the light-harvesting complex II (B800–850) from *Rhodospirillum rubrum*. Structure 1996; 4:581-97.
- [4] Hofmann E, Wrench PM, Sharples FP, Hiller RG, Welte W, Diederichs K. Structural basis of light harvesting by carotenoids: peridinin-chlorophyll-protein from *Amphidinium carterae*. Science 1996; 272:1788-91.
- [5] Standfuss J, van Scheltinga ACT, Lamborghini M, Kühlbrandt W. Mechanisms of photoprotection and nonphotochemical quenching in pea light-harvesting complex at 2.5 Å resolution. EMBO J 2005; 24:919-28.
- [6] Frank HA, Cogdell RJ. Carotenoids in photosynthesis. Photochem Photobiol 1996; 63:257-64.
- [7] Govindjee. Carotenoids in photosynthesis: An historical perspective. In: Frank HA, Young AJ, Britton G, Cogdell RJ, editors. The photochemistry of carotenoids. Dordrecht: Kluwer Academic Publishers; 1999. p. 1-19.
- [8] Zigmantas D, Hiller RG, Sundstrom V, Polivka T. Carotenoid to chlorophyll energy transfer in the peridinin-chlorophyll-a-protein complex involves an intramolecular charge transfer state. Proc Natl Acad Sci USA 2002; 99:16760-5.
- [9] Holt N, Kennis J, Dall'Osto L, Bassi R, Fleming GR. Carotenoid to chlorophyll energy transfer in light harvesting complex II from *Arabidopsis thaliana* probed by femtosecond fluorescence upconversion. Chem Phys Lett 2003; 379:305-13.
- [10] Peterman EJ, Gradinaru CC, Calkoen F, Borst JC, van Grondelle R, van Amerongen H. Xanthophylls in light-harvesting complex II of higher plants: light harvesting and triplet quenching. Biochemistry 1997; 36:12208-15.
- [11] Connelly J, Müller M, Bassi R, Croce R, Holzwarth AR. Femtosecond transient absorption study of carotenoid to chlorophyll energy transfer in the light-harvesting complex II of photosystem II. Biochemistry 1997; 36:281-7.
- [12] Cogdell RJ, Hipkins MF, MacDonald W, Truscott TG. Energy transfer between the carotenoid and the bacteriochlorophyll within the B-800-850 light-harvesting pigment-protein complex of *Rhodospseudomonas sphaeroides*. Biochim Biophys Acta 1981; 634:191-202.
- [13] van Grondelle R, Kramer HJM, Rijgersberg CP. Energy transfer in the B800-850-carotenoid light-harvesting complex of various mutants of *Rhodospseudomonas sphaeroides* and of *Rhodospseudomonas capsulata*. Biochim Biophys Acta 1982; 682:208-15.
- [14] Angerhofer A, Bomhäuser F, Gall A, Cogdell RJ. Optical and optically detected magnetic resonance investigation on purple photosynthetic bacterial antenna complexes. Chem Phys 1995; 194:259-74.
- [15] Rondonuwu FS, Yokoyama K, Fujii R, Koyama Y, Cogdell RJ, Watanabe Y. The role of the 1^1B_u state in carotenoid-to-bacteriochlorophyll singlet-energy transfer in the LH2 antenna complexes from *Rhodobacter sphaeroides* G1C, *Rhodobacter sphaeroides* 2.4.1, *Rhodospirillum rubrum* and *Rhodospseudomonas acidophila*. Chem Phys Lett 2004; 390:314-22.
- [16] Angerhofer A, Cogdell RJ, Hipkins MF. A spectral characterisation of the light-harvesting pigment-protein complexes from *Rhodospseudomonas acidophila*. Biochim Biophys Acta 1986; 848:333-41.
- [17] Chadwick BW, Zhang C, Cogdell RJ, Frank HA. The effects of lithium dodecyl sulfate and sodium borohydride on the absorption spectrum of the B800-850 light-harvesting complex from *Rhodospseudomonas acidophila* 7750. Biochim Biophys Acta 1987; 893:444-51.
- [18] Rademaker H, Hoff AJ, van Grondelle R. Carotenoid triplet yields in normal and deuterated *Rhodospirillum rubrum*. Biochim Biophys Acta 1980; 592:240-57.
- [19] Krueger BP, Lampoura SS, van Stokkum IHM, Papagiannakis E, Salverda JM, Gradinaru CC, Rutkauskas D, Hiller RG, van Grondelle R. Energy transfer in the peridinin chlorophyll-a protein of *Amphidinium carterae* studied by polarized transient absorption and target analysis. Biophys J 2001; 80:2843-55.
- [20] Linden PA, Zimmermann J, Brixner T, Holt NE, Vaswani HM, Hiller RG, Fleming GR. Transient absorption study of peridinin and peridinin-chlorophyll *a*-protein after two-photon excitation. J Phys Chem B 2004; 108:10340-5.
- [21] Kleima FJ, Hofmann E, Gobets B, van Stokkum IHM, van Grondelle R, Diederichs K, van Amerongen H. Förster excitation energy transfer in peridinin-chlorophyll-a-protein. Biophys J 2000; 78: 344-53.
- [22] Miller DJ, Catmull J, Puskeiler R, Tweedale H, Sharples FP, Hiller RG. Reconstitution of the peridinin-chlorophyll *a* protein (PCP): evidence for functional flexibility in chlorophyll binding. Photosynth Res 2005; 86:229-40.

- [23] Brotsudarmo THP, Hofmann E, Hiller RG, Wörmke S, Mackowski S, Zumbusch A, Bräuchle C, Scheer H. Peridinin-chlorophyll-protein reconstituted with chlorophyll mixtures: Preparation, bulk and single molecule spectroscopy. *FEBS Lett* 2006; 580:5257-62.
- [24] Brotsudarmo THP, Mackowski S, Hofmann E, Hiller RG, Bräuchle C, Scheer H. Relative binding affinities of chlorophylls in peridinin-chlorophyll-protein reconstituted with heterochlorophyllous mixtures. *Photosynth Res* 2008; 95:247-52.
- [25] Schulte T, Hiller RG, Hofmann E. X-ray structures of the peridinin–chlorophyll-protein reconstituted with different chlorophylls. *FEBS Lett* 2010; 584:973-8.
- [26] Schulte T, Sharples FP, Hiller RG, Hofmann E. X-ray structure of the high-salt form of the peridinin-chlorophyll *a*-protein from the dinoflagellate *Amphidinium carterae*: modulation of the spectral properties of pigments by the protein environment. *Biochemistry* 2009; 48:4466-75.
- [27] Kobayashi M, Akiyama M, Kano H, Kise H. Spectroscopy and structure determination. In: Grimm B, Porra RJ, Rüdiger W, Scheer H, editors. *Chlorophylls and bacteriochlorophylls: Biochemistry, biophysics, functions and applications*. Dordrecht: Springer; 2006. p. 79-94.
- [28] Mackowski S, Wörmke S, Brotsudarmo THP, Scheer H, Bräuchle C. Fluorescence spectroscopy of reconstituted peridinin-chlorophyll-protein complexes. *Photosynth Res* 2008; 95:253-60.
- [29] Wörmke S, Mackowski S, Brotsudarmo THP, Jung C, Zumbusch A, Ehrl M, Scheer H, Hofmann E, Hiller RG, Bräuchle C. Monitoring fluorescence of individual chromophores in peridinin–chlorophyll–protein complex using single molecule spectroscopy. *Biochim Biophys Acta* 2007; 1767:956-64.
- [30] Plumley FG, Schmidt GW. Reconstitution of chlorophyll *a/b* light-harvesting complexes: Xanthophyll-dependent assembly and energy transfer. *Proc Natl Acad Sci USA* 1987; 84:146-50.
- [31] Paulsen H, Ruml U, Rudiger W. Reconstitution of pigment-containing complexes from light-harvesting chlorophyll *a/b*-binding protein overexpressed in *Escherichia coli*. *Planta* 1990; 181:204-11.
- [32] Cammarata K, Schmidt G. In vitro reconstitution of a light-harvesting gene product: Deletion mutagenesis and analyses of pigment binding. *Biochemistry* 1992; 31:2779-89.
- [33] Paulsen H, Hobe S. Pigment-binding properties of mutant light-harvesting chlorophyll-*a/b*-binding protein. *Eur J Biochem* 1992;205:71-6.
- [34] Kuhlbrandt W, Wang DN, Fujiyoshi Y. Atomic model of plant light-harvesting complex by electron crystallography. *Nature* 1994; 367:614-20.
- [35] Remelli R, Varotto C, Sandonà D, Croce R, Bassi R. Chlorophyll binding to monomeric light-harvesting complex. *J Biol Chem* 1999; 274:33510-21.
- [36] Rogl H, Kuhlbrandt W. Mutant trimers of light-harvesting complex II exhibit altered pigment content and spectroscopic features. *Biochemistry* 1999; 38:16214-22.
- [37] Kleima FJ, Hobe S, Calkoen F, Urbanus ML, Peterman EJ, van Grondelle R, Paulsen H, van Amerongen H. Decreasing the chlorophyll *a/b* ratio in reconstituted LHClI: structural and functional consequences. *Biochemistry* 1999; 38:6587-96.
- [38] Rühle W, Paulsen H. Preparation of native and recombinant light-harvesting chlorophyll-*a/b* complex. In: Carpentier R, editor. *Photosynthesis research protocols*. New Jersey: Humana Press Inc; 2011. p. 113-25.
- [39] Gundlach K, Werwie M, Wiegand S, Paulsen H. Filling the “green gap” of the major light-harvesting chlorophyll *a/b* complex by covalent attachment of Rhodamine Red. *Biochim Biophys Acta* 2009; 1787:1499-504.
- [40] Hobe S, Prytulla S, Kuhlbrandt W, Paulsen H. Trimerization and crystallization of reconstituted light-harvesting chlorophyll *a/b* complex. *EMBO J* 1994; 13:3423-9.
- [41] Loach PA, Parkes-Loach PS. Structure-function relationships in core light-harvesting complexes (LHI) as Determined by characterization of the structural subunit and by reconstitution experiments. In: Blankenship RE, Madigan MT, Bauer CE, editors. *Anoxygenic photosynthetic bacteria*. Dordrecht: Kluwer Academic Publishers; 1995. p. 437-71.
- [42] Fiedor L, Akahane J, Koyama Y. Carotenoid-induced cooperative formation of bacterial photosynthetic LHI complex. *Biochemistry* 2004; 43:16487-96.
- [43] Fiedor L, Fiedor J, Pilch M, Susz A, Michalik M. Controlling structural and functional features of photosynthetic antenna. *Acta Phys Pol A* 2012; 122:255-8.
- [44] Karrasch S, Bullough PA, Ghosh R. The 8.5 Å projection map of the light-harvesting complex I from *Rhodospirillum rubrum* reveals a ring composed of 16 subunits. *EMBO J* 1995; 14:631-8.
- [45] Walz T, Jamieson SJ, Bowers CM, Bullough PA, Hunter CN. Projection structures of three photosynthetic complexes from *Rhodobacter sphaeroides*: LH2 at 6 Å, LHI and RC-LHI at 25 Å. *J Mol Biol* 1998; 282:833-45.
- [46] Hu X, Damjanovic A, Ritz T, Schulten K. Architecture and mechanism of the light-harvesting apparatus of purple bacteria. *Proc Natl Acad Sci USA* 1998; 95:5935-41.
- [47] Parkes-Loach PS, Majeed AP, Law CJ, Loach PA. Interactions stabilizing the structure of the core light-harvesting complex (LHI) of photosynthetic bacteria and its subunit (B820). *Biochemistry* 2004; 43:7003-16.
- [48] Meadows K, Iida K, Tsuda K, Recchia P, Heller B, Antonio B, Nango M, Loach PA. Enzymatic and chemical cleavage of the core light-harvesting polypeptides of photosynthetic bacteria: determination of the minimal polypeptide size and structure required for subunit and light-harvesting complex formation. *Biochemistry* 1995; 34:1559-74.
- [49] Davis CM, Parkes-Loach PS, Cook CK, Meadows KA, Bandilla M, Scheer H, Loach PA. Comparison of the structural requirements for bacteriochlorophyll binding in the core light-harvesting complexes of *Rhodospirillum rubrum* and *Rhodobacter sphaeroides* using reconstitution methodology with bacteriochlorophyll analogs. *Biochemistry* 1996; 35:3072-84.
- [50] Fiedor L, Scheer H, Hunter CN, Tschirschwitz F, Voigt B, Ehler J, Nibbering E, Leupold D, Elsaesser T. Introduction of a 60 fs deactivation channel in the photosynthetic antenna LHI by Ni-bacteriopeophytin *a*. *Chem Phys Lett* 2000; 319:145-52.

- [51] Fiedor L, Leupold D, Teuchner K, Voigt B, Hunter CN, Scherz A, Scheer H. Excitation trap approach to analyze size and pigment-pigment coupling: reconstitution of LH1 antenna of *Rhodobacter sphaeroides* with Ni-substituted bacteriochlorophyll. *Biochemistry* 2001; 40:3737-47.
- [52] Akahane J, Rondonuwu FS, Fiedor L, Watanabe Y, Koyama Y. Dependence of singlet-energy transfer on the conjugation length of carotenoids reconstituted into the LH1 complex from *Rhodospirillum rubrum* G9. *Chem Phys Lett* 2004; 393:184-91.
- [53] Fiedor J, Pilch M, Fiedor L. Tuning the thermodynamics of association of transmembrane helices. *J Phys Chem B* 2009; 113:12831-8.
- [54] Nakagawa K, Suzuki S, Fujii R, Gardiner AT, Cogdell R, Nango M, Hashimoto H. Probing the effect of the binding site on the electrostatic behavior of a series of carotenoids reconstituted into the light-harvesting 1 complex from purple photosynthetic bacterium *Rhodospirillum rubrum* detected by stark spectroscopy. *J Phys Chem B* 2008; 112: 9467-75.
- [55] Lambrev PH, Miloslavina Y, van Stokkum IHM, Stahl AD, Michalik M, Susz A, Tworzydło J, Fiedor J, Huhn G, Groot M-L, van Grondelle R, Garab G, Fiedor L. Excitation energy trapping and dissipation by Ni-substituted bacteriochlorophyll *a* in reconstituted LH1 complexes from *Rhodospirillum rubrum*. *J Phys Chem B* 2013; 117:11260-71.
- [56] Fiedor L, Scheer H. Trapping of an assembly intermediate of photosynthetic LH1 antenna beyond B820 subunit. Significance for the assembly of photosynthetic LH1 antenna. *J Biol Chem* 2005; 280:20921-6.
- [57] Fiedor J, Sulikowska A, Orzechowska A, Fiedor J, Burda K. Antioxidant effects of carotenoids in a model pigment-protein complex. *Acta Physica Polonica A* 2012; 59:61-64.
- [58] Kakitani Y, Akahane J, Ishii H, Sogabe H, Nagae H, Koyama Y. Conjugation-length dependence of the T1 lifetimes of carotenoids free in solution and incorporated into the LH2, LH1, RC, and RC-LH1 complexes: possible mechanisms of triplet-energy. *Biochemistry* 2007; 46:2181-97.

Reconstitution Approach to Tune Spectral Features of Light Harvesting Complexes for Improved Light Absorption and Energy Transfer

ORIGINALITY REPORT

22%

SIMILARITY INDEX

12%

INTERNET SOURCES

21%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Brotosudarmo, T.H.P.. "Peridinin-chlorophyll-protein reconstituted with chlorophyll mixtures: Preparation, bulk and single molecule spectroscopy", FEBS Letters, 20061002
Publication 1%
- 2** edoc.hu-berlin.de
Internet Source 1%
- 3** www.pnas.org
Internet Source 1%
- 4** Advances in Photosynthesis and Respiration, 2014.
Publication 1%
- 5** Fiedor, Joanna, Mariusz Pilch, and Leszek Fiedor. "Tuning the Thermodynamics of Association of Transmembrane Helices", The Journal of Physical Chemistry B, 2009.
Publication 1%
- 6** Advances in Photosynthesis and Respiration, 1995. 1%

7	pdfs.semanticscholar.org Internet Source	1%
8	www.jbc.org Internet Source	1%
9	philjournalsci.dost.gov.ph Internet Source	1%
10	Fiedor, Leszek, Junji Akahane, and Yasushi Koyama. "Carotenoid-Induced Cooperative Formation of Bacterial Photosynthetic LH1 Complex [†] ", <i>Biochemistry</i> , 2004. Publication	1%
11	Tihana Mirkovic, Evgeny E. Ostroumov, Jessica M. Anna, Rienk van Grondelle, Govindjee, Gregory D. Scholes. "Light Absorption and Energy Transfer in the Antenna Complexes of Photosynthetic Organisms", <i>Chemical Reviews</i> , 2017 Publication	1%
12	docplayer.net Internet Source	1%
13	Łukasz Orzeł, Bartłomiej Szmyd, Dorota Rutkowska-Żbik, Leszek Fiedor, Rudi van Eldik, Grażyna Stochel. "Fine tuning of copper()–chlorophyll interactions in organic media. Metalation versus oxidation of the macrocycle ", <i>Dalton Transactions</i> , 2015 Publication	1%

- | | | |
|----|--|-----|
| 14 | Advances in Photosynthesis and Respiration, 1996.
Publication | 1% |
| 15 | agro.icm.edu.pl
Internet Source | <1% |
| 16 | L Fiedor, H Scheer, C.N Hunter, F Tschirschwitz, B Voigt, J Ehlert, E Nibbering, D Leupold, T Elsaesser. "Introduction of a 60 fs deactivation channel in the photosynthetic antenna LH1 by Ni-bacteriopheophytin a", Chemical Physics Letters, 2000
Publication | <1% |
| 17 | Junji Akahane, Ferdy S Rondonuwu, Leszek Fiedor, Yasutaka Watanabe, Yasushi Koyama. "Dependence of singlet-energy transfer on the conjugation length of carotenoids reconstituted into the LH1 complex from Rhodospirillum rubrum G9", Chemical Physics Letters, 2004
Publication | <1% |
| 18 | pubs.acs.org
Internet Source | <1% |
| 19 | Marilena Di Valentin, Giancarlo Agostini, Enrico Salvadori, Stefano Ceola et al. "Triplet–triplet energy transfer in Peridinin-Chlorophyll a-protein reconstituted with Chl a and Chl d as revealed by optically detected magnetic resonance and pulse EPR: Comparison with the native PCP complex | <1% |

from *Amphidinium carterae*", *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, 2009

Publication

20

tel.archives-ouvertes.fr

Internet Source

<1 %

21

www.botanik.biologie.uni-muenchen.de

Internet Source

<1 %

22

T. H. P. Brotosudarmo, S. Mackowski, E. Hofmann, R. G. Hiller, C. Bräuchle, H. Scheer. "Relative binding affinities of chlorophylls in peridinin–chlorophyll–protein reconstituted with heterochlorophyllous mixtures", *Photosynthesis Research*, 2007

Publication

<1 %

23

d-nb.info

Internet Source

<1 %

24

reiner-lemoine-institut.de

Internet Source

<1 %

25

Brotosudarmo, Tatas Hardo Panintingjati, Leenawaty Limantara, Heriyanto, and Monika Nur Utami Prihastyanti. "Adaptation of the Photosynthetic Unit of Purple Bacteria to Changes of Light Illumination Intensities", *Procedia Chemistry*, 2015.

Publication

<1 %

26

onlinelibrary.wiley.com

Internet Source

<1 %

- 27 "Carotenoids in Photosynthesis", Springer Nature, 1993
Publication <1%
-
- 28 Roberta Croce, Giusy Canino, Francesca Ros, Roberto Bassi. "Chromophore Organization in the Higher-Plant Photosystem II Antenna Protein CP26", Biochemistry, 2002
Publication <1%
-
- 29 Legall, H.. "Near edge X-ray absorption fine structure spectroscopy (NEXAFS) of pigment-protein complexes: Peridinin-chlorophyll a protein (PCP) of *Amphidinium carterae*", Journal of Biochemical and Biophysical Methods, 20070410
Publication <1%
-
- 30 Hong Cong, Dariusz M. Niedzwiedzki, George N. Gibson, Amy M. LaFountain et al. "Ultrafast Time-Resolved Carotenoid to-Bacteriochlorophyll Energy Transfer in LH2 Complexes from Photosynthetic Bacteria", The Journal of Physical Chemistry B, 2008
Publication <1%
-
- 31 Fiedor, Leszek, Dieter Leupold, Klaus Teuchner, Berndt Voigt, C. Neil Hunter, Avigdor Scherz, and Hugo Scheer. "Excitation Trap Approach to Analyze Size and Pigment-Pigment Coupling: Reconstitution of LH1 Antenna of *Rhodobacter sphaeroides*" <1%

with Ni-Substituted Bacteriochlorophyll[†]",
Biochemistry, 2001.

Publication

32

Dyer, A.. "Cation-exchange in synthetic zeolite L: The exchange of hydronium and ammonium ions by alkali metal and alkaline earth cations", Zeolites, 199304/05

<1%

Publication

33

A.A. Moskalenko. "Effects of Carotenoid Inhibition on the Photosynthetic RC–LH1 Complex in Purple Sulphur Bacterium *Thiorhodospira sibirica*", Photosynthesis Research, 11/2005

<1%

Publication

34

Tomáš Polívka, Roger G. Hiller, Harry A. Frank. "Spectroscopy of the peridinin–chlorophyll-a protein: Insight into light-harvesting strategy of marine algae", Archives of Biochemistry and Biophysics, 2007

<1%

Publication

35

Samuel D. Bockenhauer, W. E. Moerner. "Photo-Induced Conformational Flexibility in Single Solution-Phase Peridinin-Chlorophyll-Proteins", The Journal of Physical Chemistry A, 2013

<1%

Publication

36

mrcpp.machung.ac.id

Internet Source

<1%

37 Robielyn P. Ilagan, Sumie Shima, Alexander Melkozernov, Su Lin et al. " Spectroscopic Properties of the Main-Form and High-Salt Peridinin–Chlorophyll Proteins from ", Biochemistry, 2004

Publication

38 Magdaong, Nikki M., Amy M. LaFountain, Jordan Alan Greco et al. "High Efficiency Light-Harvesting by Carotenoids in the LH2 Complex from Photosynthetic Bacteria: Unique Adaptation to Growth Under Low-Light Conditions", The Journal of Physical Chemistry B

Publication

39 Croce, Roberta, Giusy Canino, Francesca Ros, and Roberto Bassi. "Chromophore Organization in the Higher-Plant Photosystem II Antenna Protein CP26", Biochemistry, 2002.

Publication

40 Janik, Ewa, Joanna Bednarska, Monika Zubik, Karol Sowinski, Rafal Luchowski, Wojciech Grudzinski, and Wieslaw I. Gruszecki. "Is It Beneficial for the Major Photosynthetic Antenna Complex of Plants to Form Trimers?", The Journal of Physical Chemistry B

Publication

41 intl.biophysj.org
Internet Source

<1 %

42

H.A. Frank, R.J. Cogdell. "8.6 Light Capture in Photosynthesis", Elsevier BV, 2012

Publication

<1 %

43

www.biophysj.org

Internet Source

<1 %

44

Advances in Photosynthesis and Respiration, 2008.

Publication

<1 %

45

The Photosynthetic Bacterial Reaction Center, 1988.

Publication

<1 %

46

Richard J. Cogdell, Andrew Gall, Jürgen Köhler. " The architecture and function of the light-harvesting apparatus of purple bacteria: from single molecules to membranes ", Quarterly Reviews of Biophysics, 2006

Publication

<1 %

47

Photosynthesis Energy from the Sun, 2008.

Publication

<1 %

48

H. Paulsen, U. Rümmler, W. Rüdiger. "Reconstitution of pigment-containing complexes from light-harvesting chlorophyll a/b-binding protein overexpressed in *Escherichia coli*", Planta, 1990

Publication

<1 %

49	www.mdpi.com Internet Source	<1 %
50	bmcplantbiol.biomedcentral.com Internet Source	<1 %
51	Submitted to University of Sheffield Student Paper	<1 %
52	www.phototech.eu Internet Source	<1 %
53	Nikodem Czechowski, Piotr Nyga, Mikołaj K. Schmidt, Tatas H. P. Brotosudarmo et al. "Absorption Enhancement in Peridinin–Chlorophyll–Protein Light-Harvesting Complexes Coupled to Semicontinuous Silver Film", Plasmonics, 2011 Publication	<1 %
54	theses.gla.ac.uk Internet Source	<1 %
55	Harvey, Pierre D., Christine Stern, and Roger Guilard. "49 Bio-inspired Molecular Devices Based on Systems Found in Photosynthetic Bacteria", Handbook of Porphyrin Science, 2011. Publication	<1 %
56	carotenoidsociety.org Internet Source	<1 %
57	epdf.tips Internet Source	<1 %

58

Nakagawa, Katsunori, Satoru Suzuki, Ritsuko Fujii, Alastair T. Gardiner, Richard J. Cogdell, Mamoru Nango, and Hideki Hashimoto. "Probing the Effect of the Binding Site on the Electrostatic Behavior of a Series of Carotenoids Reconstituted into the Light-Harvesting 1 Complex from Purple Photosynthetic Bacterium Rhodospirillum rubrum Detected by Stark Spectroscopy", The Journal of Physical Chemistry B, 2008.

Publication

<1 %

59

"Chlorophylls and Bacteriochlorophylls", Springer Nature, 2006

Publication

<1 %

60

"The Photochemistry of Carotenoids", Springer Nature, 2004

Publication

<1 %

61

Advances in Photosynthesis and Respiration, 2010.

Publication

<1 %

62

Current Research in Photosynthesis, 1990.

Publication

<1 %

63

Parak, Wolfgang J., Marek Osinski, Kenji I. Yamamoto, Aneta Prymaczek, and Sebastian Mackowski. "Plasmonics with silver nanowires: plasmons affect the energy transfer", Colloidal Nanoparticles for Biomedical Applications IX, 2014.

Publication

<1 %

Tomáš Polívka, Villy Sundström. "Ultrafast Dynamics of Carotenoid Excited States—From Solution to Natural and Artificial Systems", *Chemical Reviews*, 2004

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On