

# Isolation of Active Compound from Zingiber Purpureum Roxb. Using Bioassay Guided

*by* Rollando S. Farm

---

**Submission date:** 29-Oct-2019 07:20PM (UTC-0700)

**Submission ID:** 1203229908

**File name:** Compound\_from\_Zingiber\_Purpureum\_Roxb.\_Using\_Bioassay\_Guided.pdf (366.58K)

**Word count:** 3425

**Character count:** 16690



10

Rollando Rollando, Maureen Engracia, Laurensia Nina Irawati, Edward Hartono, Eva Monica, Rehmadata Sitepu

Program of Pharmacy, <sup>24</sup> University, MalangVila

\*Corresponding Author: <sup>24</sup> : ro.rollando@

<sup>15</sup> Zingiber purpureum Roxb. was a plant that has the potential to be developed as a chemotherapy agent. Bioassay guided fractionation aims to examine the fractions from <sup>15</sup> of bangle plant rhizome to colon cancer cell type WiDr and Vero cells. The apoptotic test and modulating <sup>15</sup> by <sup>15</sup> isolation process of <sup>15</sup> most active fraction. The pure isolates obtained were identified by structures with structural elucidation using ultraviolet, 1D-NMR, 2D-NMR and liquid chromatography-mass spectrometer. Obtained <sup>15</sup> compound of ethyl acetate fraction number 21. With a value of CC<sub>50</sub> to a WiDr cell of 11.24 ± 3.44 µg/mL and 342. 22 ± 4.98 µg/mL to Vero cells.

<sup>4</sup> **Keywords:** *Zingiber purpureum* Roxb., WiDr, Vero,

## Introduction

Colorectal cancer ranked fourth in all men and women worldwide, with an incidence of 1,400,000 new cases diagnosed in 2012 [1]. IARC states that cancers of the colon and rectum are the most common incidence of cancer with the number 3 (34,000 per year, 15.9%) and the third cause of death (18,000 per year, 10.8%). Chemotherapy was one of the therapies in the treatment of cancer. Chemotherapy agents generally have low selectivity properties because they are antiproliferative to normal selkanker and cell [2]. In addition, some chemotherapy has a narrow therapeutic index, and may lead to multidrug resistance (MDR) and adverse side effects [3].

One of the problems that often arise in the treatment of cancer is the resistance of chemotherapy drugs (drug resistance). Doxorubicin is a chemotherapy agent of various types of cancer of the anthracycline class that has provoked resistance. In addition to causing resistance, doxorubicin may also cause cardiotoxicity in long-term

use [4]. Increased doses of doxorubicin are not an appropriate way to overcome resistance problems because they can cause normal cells to be exposed by cytotoxic drugs that will trigger toxicity and transform normal cells into cancer cells [5].

One of the medicinal plants in Indonesia that potentially developed as a chemopreventive agent in colon cancer is bangle. The bangle plant has been used by society traditionally to cope with several diseases. Empirically, boiled water rhizome bangle used the community as a headache medicine, constipation drugs, abdominal pain, jaundice, as body warmers, slimming, helps the gas out of the digestive tract, asthma, rheumatism, and antipyretics.

The research on scientific proof of the efficacy of the rhizome of bangle plant is still minimal publication and the efficacy of rhizome of bangle plant still many that have not been explored. So in this study, conducted an active fraction search on the ethanolic extract

of rhizome plant bangle that can kill and suppress the growth of cancer cells, especially in colon cancer cell type WiDr.

### Material and Instrument

Methanol, Hexane and ethyl acetate fraction 1 fal oak bark, DMSO 0.1%, cisplatin (Wako), (FBS) % (v/v) (Qualified, Gibco, Invitrogen USA), penicillin-streptomycin 1.5% (v/v) (Gibco, Invitrogen USA and Fungizone 0.5% v/v Gibco), Trypsin-E 0.25% (Gibco, Invitrogen Canada).

MTT prepared with a concentration of dissolved 1 x 10<sup>5</sup>, 0.01 N HCl (Merck, Darmstadt, Germany). PBS containing

Liquid nitrogen, cabinets, micropipet (Eppendorf), balance sheets (Ohaus), Benchmark FACTS caliber Biospectrometry NMR 400MHz, stirrer (Nuova, Thermolyne).

### Procedure

#### Plant Material

(250 g) ethanol maceration method.

filtere

### Fractionation with Column Chromatography

Separation of the active compound GF<sub>204</sub> and the 200 mL n-hexane, ethyl acetate and methanol as motion phase was carried out gradiently with the addition of 1 drop of CH<sub>3</sub>COOH. Each fraction obtained was evaporated at ± 40 °C to obtain a concentrated fraction and then dried in an oven at ± 50 °C. Each fraction tested to the cytotoxic activity to obtain the value of CC<sub>50</sub>.

DMEM each fraction

%.

### Isolation and Purification under Bioassay-Guided Screening

According to cytotoxicity test UV, infrared, LC-MS one (ethyl acetate fraction 21). The fraction (4.7).

## Determination of Cell Cycle and Apoptosis



## Result and Discussion

### Results of Cytotoxic Test

The cytotoxic activity of the fraction was expressed as  $CC_{50}$ . The test was performed by dissolving 44 fractions of fractionation of ethanolic extract with gradient system (n-

hexane, ethyl acetate and methanol), then into it added DMSO solution until obtained the desired concentration. A single cytotoxic effect showed decreased cell viability and morphological changes in colon cancer cells of WiDr. The test results (Fig.1) show that Ethyl Acetate Fraction 21 treatment (Fig. 1b-c) shows the wrinkled cell. This indicates a decrease in living cells when compared to controls (Fig. 1a).

appear to indicate unknown changes were due to cell deaths due to necrotic or apoptotic processes and proliferative inhibition processes. Meanwhile, the  $CC_{50}$  value of the largest was ethyl acetate fraction 21 with concentrations of  $11.24 \pm 3.44$   $\mu\text{g}/\text{mL}$  and doxorubicin was  $13.88 \pm 0.34$   $\mu\text{g}/\text{mL}$ .




Fig.1: Ethyl Acetate Fraction 21 testing was performed using cell controls (a) and Vero cells (d) Effect of Ethyl Acetate Fraction 21 on WiDr colon cancer cells (b-c). Figure (b) at concentrations of 200  $\mu\text{g}/\text{mL}$  and image (c) Vero cells treated with a fraction of concentration of 200  $\mu\text{g}/\text{mL}$ .

**Table 1: Cytotoxic activity on WiDr and Vero**

Number	Experiment	CC <sub>50</sub> on WiDr Cell	CC <sub>50</sub> on Vero Cell
1	Doxorubicin	13,88±0,34 µg/mL	160,67± 76 µg/mL
2	Methanol fraction 1	56,54 ±1,67 µg/mL	240,89 ± 96 µg/mL
3	Methanol fraction 2	70,92 ± 1,92 µg/mL	140,92 ± 1,22 µg/mL
4	Methanol fraction 3	31,22 ±2,12 µg/mL	341,67 ± 2,39 µg/mL
5	Methanol fraction 4	48,88 ± 1,88 µg/mL	131,33 ± 1,96 µg/mL
6	Methanol fraction 5	74,38 ± 1,55 µg/mL	402,43 ± 1,07 µg/mL
7	Methanol fraction 6	99,08 ±0,87 µg/mL	148,56 ± 3,84 µg/mL
8	Methanol fraction 7	58,33 ±1,99 µg/mL	340,44 ± 1,45 µg/mL
9	Ethyl Acetate Fraction 1	41,11 ±2,68 µg/mL	441,44 ± 0,34 µg/mL
10	Ethyl Acetate Fraction 2	66,78 ±1,03 µg/mL	561,90 ± 1,52 µg/mL
11	Ethyl Acetate Fraction 3	36,99 ±0,86 µg/mL	445,95 ± 1,14 µg/mL
12	Ethyl Acetate Fraction 4	98,82 ±3,35 µg/mL	123,98 ± 3,22 µg/mL
13	Ethyl Acetate Fraction 5	95,85 ±3,08 µg/mL	248,07 ± 2,58 µg/mL
14	Ethyl Acetate Fraction 6	88,55 ±2,88 µg/mL	338,77 ± 4,95 µg/mL
15	Ethyl Acetate Fraction 7	13,11 ±2,08 µg/mL	565,44 ± 1,08 µg/mL
16	Ethyl Acetate Fraction 8	35,88 ±1,07 µg/mL	341,69 ± 0,11 µg/mL
17	Ethyl Acetate Fraction 9	32,54 ±1,91 µg/mL	249,22 ± 0,65 µg/mL
18	Ethyl Acetate Fraction 10	16,22 ±0,79 µg/mL	241,53 ± 0,96 µg/mL
19	Ethyl Acetate Fraction 11	99,11 ±2,16 µg/mL	299,19 ± 4,98 µg/mL
20	Ethyl Acetate Fraction 12	88,19 ±5,12 µg/mL	349,93 ± 0,96 µg/mL
21	Ethyl Acetate Fraction 13	14,11 ±4,33 µg/mL	240,11 ± 1,93 µg/mL
22	Ethyl Acetate Fraction 14	88,11 ±3,16 µg/mL	360,24 ± 5,11 µg/mL
23	Ethyl Acetate Fraction 15	16,92 ± 2,12 µg/mL	190,11 ± 8,22 µg/mL
24	Ethyl Acetate Fraction 16	28,11 ±3,30 µg/mL	223,11 ± 2,30 µg/mL
25	Ethyl Acetate Fraction 17	92,22 ±3,93 µg/mL	292,12 ± 2,16 µg/mL
26	Ethyl Acetate Fraction 18	31,21 ±3,33 µg/mL	234,44 ± 4,12 µg/mL
27	Ethyl Acetate Fraction 19	39,92 ±4,22 µg/mL	234,11 ± 3,16 µg/mL
28	Ethyl Acetate Fraction 20	21,22 ±2,22 µg/mL	134,21 ± 3,12 µg/mL
29	Ethyl Acetate Fraction 21	11,24 ±3,44 µg/mL	342,11 ± 4,98 µg/mL
30	Ethyl Acetate Fraction 22	41,98 ±9,92 µg/mL	232,22 ± 1,92 µg/mL
31	Ethyl Acetate Fraction 23	34,22 ±2,33 µg/mL	234,03 ± 3,22 µg/mL
32	Ethyl Acetate Fraction 24	92,23 ±2,22 µg/mL	445,11 ± 2,39 µg/mL
33	Ethyl Acetate Fraction 25	82,11 ±3,12 µg/mL	354,73 ± 1,39 µg/mL
34	Hexane fraction 1	77,92 ± 1,93 µg/mL	551,82 ± 3,11 µg/mL
35	Hexane fraction 2	98,28 ±3,34 µg/mL	458,11 ± 1,28 µg/mL
36	Hexane fraction 3	90,01 ± 1,88 µg/mL	388,29 ± 3,29 µg/mL
37	Hexane fraction 4	34,22 ±1,12 µg/mL	123,21 ± 2,92 µg/mL
38	Hexane fraction 5	22,11 ± 1,62 µg/mL	220,91 ± 0,16 µg/mL
39	Hexane fraction 6	97,14 ± 1,62 µg/mL	240,11 ± 1,12 µg/mL
40	Hexane fraction 7	36,22 ±4,11 µg/mL	233,11 ± 1,92 µg/mL
41	Hexane fraction 8	90,11 ± 1,66 µg/mL	102,11 ± 1,81 µg/mL
42	Hexane fraction 9	83,29 ± 1,98 µg/mL	358,09 ± 0,12 µg/mL
43	Hexane fraction 10	73,12 ± 1,93 µg/mL	241,72 ± 1,12 µg/mL
44	Hexane fraction 11	99,54 ±1,67 µg/mL	229,11 ± 1,88 µg/mL
45	Hexane fraction 12	96,11 ±2,22 µg/mL	221,11 ± 0,11 µg/mL

**Structure Elucidation**

Compound 1, yellow crystal, m.p. 97-99 °C, UV (MeOH) λ<sub>max</sub> nm (log ε): 232 (3.92), 290 (4.10), and 325 nm (3.66). EIMS: m/z (270, M<sup>+</sup>, 100, base peak), 193 (76.1), 166 (80.3), 138 (41.4), 105 (21.2), and 77 (16.4). <sup>1</sup>H NMR (400 MHz in acetone d<sub>6</sub>), δ<sub>H</sub> ppm: 12.17 (1H, br, s, 5-OH), 7.58 (3H, m, H-3',4',5'), 7.41 (2H, m, H-2',6'), 6.09 (1H, d, J=2.2 Hz, H-8), 6.07 (1H, d, J=2.2 Hz, H-6), 5.61 (1H, dd, J=3.8; 12.3 Hz, H-2), 3.35 (3H, s, 7-OCH<sub>3</sub>),

3.24 (1H, dd, J=12.8; 16.4 Hz, H-3<sub>ax</sub>), and 2.81 (1H, dd, J=3.8; 16.4 Hz, H-3<sub>eq</sub>). <sup>13</sup>C NMR (100 MHz in acetone d<sub>6</sub>), δ<sub>C</sub> ppm: 196.3 (C-4), 168.6 (C-7), 164.8 (C-5), 163.3 (C-8a), 140.1 (C-1), 129.6 (C-3',5'), 128.5 (C-4'), 123.9 (C-2',6'), 101.5 (C-4a), 95.09 (C-8), 95.66 (C-6), 79.6 (C-2), 56.1 (7-OCH<sub>3</sub>), and 43.9 (C-3). The elucidation results showed that the compound of the ethyl acetate fraction 21 was 2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one.




Fig. 20 Compound (1) (JMBC)

Observation of [redacted]

Furthermore, [redacted] test of cell cycle. The process of [redacted] cells [redacted] same cell [redacted] as normal cells [6]. [redacted] inhibiting [redacted] cell [redacted] is the modulation of cell cycle that can be observed using flowcytometry method (Rolando, 2000).

Flowcytometry is [redacted] each [redacted] cell [redacted] the number of chromosomes on [redacted] [7]. [redacted] is [redacted] because [redacted] is capable of interacting with DNA [8]. [redacted] profile [redacted] 24 hours [9]. [redacted] percentage of cell cycle distribution in detail is shown in Table 2.

Table 2: Percentage of cell cycle distribution

Sample	G1 (%)	S (%)	G2M (%)	% CV
Control	38,11	12,34	13,34	3,87
Doxorubicin 2,5 μM	42,88	13,11	24,55	2,22
2,3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one 2,5 μM	34,11	23,11	11,23	5,98




Fig. 3: Distribution of cell cycle percentages. (a) Control (b) Doxorubicin (c) [redacted] Compound

[redacted] test results showed doxorubicin causing accumulation in phase S. The [redacted]

[redacted] compound causes cell accumulation in phase [redacted] when compared with cell control. In the [redacted]

in amounted to 23.11% greater than the single doxorubicin of 13.11%. Cell accumulation in the S-phase of combination was increased compared with

cells without treatment (cell control) of 12.34%. Cell accumulation is possible because of cell cycle arrest in that phase.

**Table 3: Percentage of death after treatment**

	Control	Doxorubicin 2.5 µM	2,3- 2.5 µM
Initial apoptosis (%)	2.11	2.33	9.87
End apoptosis (%)	1.63	2.53	1.24
Necrosis (%)	0.86	0.87	1.62
Total	3.94	6.04	13.78

**Observation of apoptotic compounds**

flow cytometer. Induction apoptosis was used to investigate the mechanism of cell death from treatment of the compound on colon cancer cell WiDr incubated for 24 hours [10].

The study is flowcytometry of apoptosis that occurred in the treated cells. Annexin V phospholipid binding are on strongly negatively charged cellular membranes. Cell cell death caused by apoptosis or necrosis can be distinguished by the staining of Propidium Iodide (PI) through intercalation with DNA [11]. The percentage of cell death after treatment of induced 3.

Analysis of the percentage of cell death after (Table 3) showed that

**References**

1. Yuniati Y, Alfanaar R, Rollando R (2018) Structural conformational study of isoflavon derivatives in soybean using semiempirical methods. *Journal of Global Pharma Technology*, 10(03):220-225.
2. Valeriote F, Greshaber CK, Media J, Pietraszkewicz H, Hoffmann J, Pan M (2002) Discovery and development of anticancer agents from plants. *J. Exp. Ther. Oncol.*, 2:228-236.
3. Rollando R (2018) Combination of hedyotis corymbosa L. and tinospora crispa ethanolic extract increase cisplatin cytotoxicity on T47D

control cells showed a 3.94% percentage of cell death. Single-treated cells with doxorubicin showed 6.04% cell death, whereas 2, 3-dihydro-5- hydroxy-7- methoxy-2-phenylchromen-4-one compound resulted in 13.78% cell death. It showed an increase in the percentage of cell death by 7.74% in the 2,3-dihydro- 5-hydroxy- 7-methoxy- 2-phenylchromen-4-one compound compared with doxorubicin.

**Conclusion**

The results showed that isolate 2 was 3-dihydro-5-hydroxy-7-methoxy-2-phenylchromen-4-one having activity cytotoxic in breast cancer cell type T47D with active category with IC<sub>50</sub> 11, 24 ±3,44 µg/mL.

**Acknowledgements**

for on Student Creativity Program 2018.

4. (2018)
5. Bravo-Cellar
6. Amular-Lemarriva A

pentoxifyline increases . Molecular  
29  
6. .  
7. (2018) .  
10(2) -254.  
8. dan Prilianti KR (200) .  
20 . J. Pharm. Sci.  
Community;14:1-14.  
9. (2006) .  
21 . The  
Journal of Antibiotics, 59(4):209-211.  
10. (2018)  
Cytotoxic activity of 2- iminoethyl 2-(2-(1-  
hydroxypentan-2-yl) phenyl) acetate from  
sterculia quadrifida R.Br ethyl acetate  
fraction. Journal of Global Pharma  
Technology, 10(06):204-212.



# Isolation of Active Compound from Zingiber Purpureum Roxb. Using Bioassay Guided

## ORIGINALITY REPORT

15%

SIMILARITY INDEX

32%

INTERNET SOURCES

21%

PUBLICATIONS

23%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://media.neliti.com">media.neliti.com</a> Internet Source	1%
2	<a href="http://europepmc.org">europepmc.org</a> Internet Source	1%
3	S. Bouazzi, H. Jmii, R. El Mokni, K. Faidi, D. Falconieri, A. Piras, H. Jaïdane, S. Porcedda, S. Hammami. "Cytotoxic and antiviral activities of the essential oils from Tunisian Fern, <i>Osmunda regalis</i> ", <i>South African Journal of Botany</i> , 2018 Publication	1%
4	<a href="http://malariajournal.biomedcentral.com">malariajournal.biomedcentral.com</a> Internet Source	1%
5	Submitted to Higher Education Commission Pakistan Student Paper	1%
6	<a href="http://perpustakaan.unitomo.ac.id">perpustakaan.unitomo.ac.id</a> Internet Source	1%
7	Submitted to University of Kufa Student Paper	

1%

8

[simdos.unud.ac.id](http://simdos.unud.ac.id)

Internet Source

1%

9

R. Srinivasan, D. Natarajan, M.S. Shivakumar,  
T. Vinuchakkaravarthy, D. Velmurugan.

"Bioassay guided isolation of mosquito larvicidal  
compound from acetone leaf extract of  
Elaeagnus indica Servett Bull and its in-silico  
study", Industrial Crops and Products, 2015

Publication

1%

10

[journal.uad.ac.id](http://journal.uad.ac.id)

Internet Source

1%

11

Rollando Rollando, Maywan Hariono.

"Antimicrobial, Antioxidant and T47D Cytotoxic  
Activities of Trichaptum sp., A Fungal  
Endophyte from Phyllanthus niruri Linn.: In vitro  
and in silico Studies", Asian Journal of Cell  
Biology, 2016

Publication

1%

12

[jpsr.pharmainfo.in](http://jpsr.pharmainfo.in)

Internet Source

1%

13

[journals.plos.org](http://journals.plos.org)

Internet Source

1%

14

Hua-Bin Li, Feng Chen, Yue Jiang.

1%

"Determination of vitamin B12 in multivitamin tablets and fermentation medium by high-performance liquid chromatography with fluorescence detection", Journal of Chromatography A, 2000

Publication

15

[e-journal.usd.ac.id](http://e-journal.usd.ac.id)

Internet Source

1%

16

Submitted to University of Newcastle

Student Paper

1%

17

[innovareacademics.in](http://innovareacademics.in)

Internet Source

1%

18

Sbai, H., I. Zribi, M. DellaGreca, and R. Haouala. "Bioguided fractionation and isolation of phytotoxic compounds from *Apium graveolens* L. aerial parts (Apiaceae)", South African Journal of Botany, 2016.

Publication

1%

19

Submitted to Texas A&M University, College Station

Student Paper

1%

20

H. Sbai, I. Zribi, M. DellaGreca, R. Haouala. "Bioguided fractionation and isolation of phytotoxic compounds from *Apium graveolens* L. aerial parts (Apiaceae)", South African Journal of Botany, 2017

Publication

1%

---

21	<a href="http://ijbd.ir">ijbd.ir</a> Internet Source	1%
22	<a href="http://www.jpsr.pharmainfo.in">www.jpsr.pharmainfo.in</a> Internet Source	1%
23	<a href="http://digital.csic.es">digital.csic.es</a> Internet Source	1%
24	Leny Yuliati, Juliana, R Indrawati. "Tuning the stability of red color natural pigments in fruit extracts by pH control", Journal of Physics: Conference Series, 2019 Publication	1%
25	Submitted to Udayana University Student Paper	<1%
26	<a href="http://molecular-cancer.biomedcentral.com">molecular-cancer.biomedcentral.com</a> Internet Source	<1%
27	<a href="http://www.tandfonline.com">www.tandfonline.com</a> Internet Source	<1%
28	Ntie-Kang, Fidele, Pascal Onguéné, Lydia L Lifongo, Jean Ndom, Wolfgang Sippl, and Luc Mbaze. "The potential of anti-malarial compounds derived from African medicinal plants, part II: a pharmacological evaluation of non-alkaloids and non-terpenoids", Malaria Journal, 2014. Publication	<1%

---

29 [repositorio.unicamp.br](http://repositorio.unicamp.br)

Internet Source

<1%

---

30 Submitted to University of Sheffield

Student Paper

<1%

---

31 Sergio Luiz Dalmora, Silvia Maria Krug Masiero, Paulo Renato de Oliveira, Maximiliano da Silva Sangoi, Liberato Brum Junior. "Validation of an RP-LC Method and Assessment of rhG-CSF in Pharmaceutical Formulations by Liquid Chromatography and Biological Assay", Journal of Liquid Chromatography & Related Technologies, 2007

Publication

<1%

---

32 Submitted to Panjab University

Student Paper

<1%

---

Exclude quotes  On

Exclude matches  < 5 words

Exclude bibliography  On

# Isolation of Active Compound from Zingiber Purpureum Roxb. Using Bioassay Guided

---

## GRADEMARK REPORT

---

FINAL GRADE

**/100**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---