

Structural Conformational Study of Isoflavon Derivatives in Soybean

by Rollando S. Farm

Submission date: 31-Oct-2019 05:46PM (UTC-0700)

Submission ID: 1204629730

File name: ral_Conformational_Study_of_Isoflavon_Derivatives_in_Soybean.pdf (512.29K)

Word count: 2451

Character count: 12190

Structural Conformational Study of Isoflavon Derivatives in Soybean Using Semi empirical Methods

Yuyun Yuniati^{1*}, Rokiy Alfanaar¹, Rollando Rollando²

¹ Program of Chemistry, Faculty of Science and Technology, Ma Chung University, Malang, Villa Puncak Tidar N-01, Malang 65151.

² Program of Pharmacy, Faculty of Science and Technology, Ma Chung University, Malang Villa Puncak Tidar N-01, Malang 65151.

* Corresponding Author: Yuyun Yuniati

Abstract

Isoflavon was flavonoid group which was a polyphenolic compound. As a member of flavonoid groups, isoflavon was a bioactive compound that contains phenolic group reported to have an ability as antioxidant and prevent free radical damage through two mechanism. The mechanism are donating hydrogen ions and directly acting as a free radical scavenger. One of the seed plants which contain isoflavon was soybean. Isoflavon derivatives in soybean are biochanin a, genistein, daidzein, equol, formononetin, dan glycitein. The compounds can be applied in medical and pharmaceutical. Generally the is of lavon derivatives in soy bean have been conducted to anti-allergic, antioxidant, dan anti cholesterol. These studies based on optimization and structural conformational to support QSAR and Molecular docking studies. In this study, semiempirical AM1 method was used for optimization all the structure. The distance of two nearby atoms affect the structure stability and the strain energy because the lower the strain energy, the more stable that compound. The most stable structure for biochanin a, genwastein, daidzein, glycitein, formononetin was 40° while the most stable structure for equol was 280°.

Keywords: Conformation, Derivatives, Energy, Isoflavon, Soybean.

Introduction

Isoflavon was flavonoid group which was a polyphenolic compound. The based chemistry structure of isoflavon almost equal with flavonoid, that consist of two benzene rings (A-ring and B-ring) which was bind to heterocyclic pyran ring (C-ring). The difference was in the B-ring orientation. B-ring in flavonoid compound was bind by the number 2 carbon (C2) in C-ring to the number 1 carbon in B-ring; hence, isoflavon was bind by the number 3 carbon (C3) in C-

ring [1]. As a member of flavonoid groups, isoflavon was a bioactive compound that contains phenolic group reported to have ability as antioxidant and prevent free radical damage through two mechanism [2].

The Mechanism of Antioxidant in Flavonoid Groups Are

- By donating hydrogen ions [3]
- By directly acting as a free radical scavenger [1]




Figure 1: Structure of the flavonoids [with two aromatic benzol rings (A and B rings)] and a C3 portion cyclized with oxygen (C ring)

One of the seed plants which contain isoflavon was soybean. As we know, soy was widely used in Asia as a staple food and consumed regularly in traditional food items such as tofu, miso, natto, edamame (whole soybeans), soybean paste, and shoyu (fermented soy sauce). In soybean contain isoflavon compound that have an ability as an antioxidant that can be anti cancer. Hence, the isoflavone intake among Asians

was about a factor of 100 higher than that of people in the Western world. The daily isoflavone intake among Southeast Asians ranges between 15 and 47 mg, while European consumes only between 0.15 and 1.7 mg isoflavones per day. Isoflavon derivatives in soybean are biochanin a, genwastein, daidzein, equol, formononetin, dan glycitein [4].



Figure 2: Structure of is flavone: geniste in, daidze in, for mononeter, biochanin A, and glycitein [3].

The compounds can be applied in medical and pharmaceutical. Generally the isoflavon derivatives in soybean have been conducted to anti-allergic, antioxidant, dan anti cholesterol. Genistein has role as antioxidant, anti-angiogenik, metastasis cancer cell, protein kinase inhibitor and proliferasi inhibitor [2].

Biochanin a can prevent high cholesterol, anti-fungal, anti tumor, regulate hormonal stage in human body, anti cancer, antispasmodic, and lower lipidin blood. Daidzein can prevent and lower cancer risk, absence of tumor, cure syndrome activity, such as menopause syndrome and anti-osteoporosis. Formononetin was anti cancer (breast cancer, prostat cancer, dan colon cancer), have effect to reduce fat. Glycitein can be use for bone healthy, menopause

symptoms, to reduce the stress hormones and anti inflammation that lead by cancer [3]. One of the best way to know all the function from the isoflavones derivatives through QSAR and Molecular docking.

These studies based on optimization and structural conformational to support QSAR and Molecular Docking studies [7]. The result in study can find the possibility structure of isoflavones derivatives in soybean as a cancer drugs. In this study, semiempirical AM1 method was used for optimization all the structure. Austin Model 1 or AM 1 was first introduced by Dewar when he was in the Texas University, Austin. AM1 has a specific parameters than MNDO, 14 parameters/atom. The other difference was AM 1 uses a new tribe to describe their interaction [8].

$$E_{AB} = Z_A Z_B < S_A S_A | S_B S_B > + \frac{Z_A Z_B}{R_{AB}} \sum_{i=1}^n [\alpha_i^A \exp(-b_i^A (r_{AB} - c_i^A)^2) + \alpha_i^B \exp(-b_i^B (r_{AB} - c_i^B)^2)]$$

a,b and c are parameters from Gaussian function that was associated with atoms in the definition (superscript).

Experimental Section

Materials

Computer (ASUS Vivo book Flip TP410UA-DB71T 14-inch FHD Touchscreen, Intel Core i7).

Optimized Structural Conformational

Structural conformational of the optimized isoflavones derivatives was carried out with the use of an efficient program, AM1. Hypechem was used to draw and optimization all the structure. To get the result for structural conformation, constrain variation bond angle between C(2)-C(3), C(3)-C(1), and C(1)-C(2) each ten degree, from

zero degree to three hundred and sixty degree. So, strain energy and bond length has been obtained. Make strain energy graphics to identify the stable and unstable structure from isoflavones derivatives. Calculate the bond length based on numbering of carbon atoms (IUPAC). The methodology can be expressed as following (Figure 3).




Figure 3: Methodology of structural conformational study of isoflavon derivatives

Results and Discussion

Optimization AM 1 for Isoflavon Derivatives in Soybean

The bond length result in Table 1 show the bond length between one compound and another was similar where was the difference of one compound and the other compounds was about 0,016 Å. The differences from the

bond length results of one compound minus with the others and then sum all the result. The totals divide by the amount of all bond lengths. But there was a structure difference between the five isoflavons and equal, the difference was in the C(3)-C(2) bond, in which the orbital changes from sp^2 to sp^3 . The bond length of equal was longer than the other structures.

Table 1: Bond Length between Each Atoms of Isoflavon Derivatives in Soybean

Atom	Bond Length (Å)					
	Genistein	Daidzein	Glycitein	Formononetin	Biochanin A	Equal
C(1)-C(2)	1.375	1.3736	1.3720	1.3738	1.3752	1.4021
C(2)-C(3)	1.3538	1.3559	1.3564	1.3559	1.3538	1.5207
C(3)-C(4)	1.4784	1.4745	1.4735	1.4745	1.4784	1.5197
C(5)-C(6)	1.4007	1.3858	1.3872	1.3858	1.4087	1.3873
C(6)-C(7)	1.4032	1.4095	1.4231	1.4049	1.4032	1.4070
C(7)-C(8)	1.4006	1.3998	1.3947	1.3997	1.4005	1.401
C(8)-C(1)	1.4597	1.4595	1.4595	1.4594	1.4595	1.455
C(1)-C(2)	1.4044	1.4019	1.4044	1.4004	1.4051	1.4019
C(2)-C(3)	1.3893	1.3915	1.3892	1.3933	1.3879	1.3907
C(3)-C(4)	1.4940	1.4918	1.495	1.3995	1.4078	1.4026
C(5)-C(6)	1.4018	1.4049	1.4017	1.4078	1.3885	1.4041
C(6)-C(7)	1.3915	1.389	1.3915	1.3875	1.3932	1.3969
C(8)-C(1)	1.4925	1.495	1.4924	1.4957	1.4911	1.3999

Conformational Study

In structural conformation study, the strain energy graphic (Figure 2) indicates the extreme values of each compound in soybean isoflavones derivatives. Biochanin a,

daidzein, glycitein, formononetin, and genistein have the similar structure (Figure 2). Because of that, they have similar angle of extreme values at 40°, 80°, 140°, 180°, 220°.

dan 320°. But there was a difference extreme values in equal, the extreme values were at 80°, 140°, 280°, dan 340°. It was because there

was no double bond in equal structure, which the others have [9].




Figure 4: Strain Energy of Isoflavon Derivatives in Soybean a) Biochanin a; b) Daidzein; c) Glycitein; d) Genistein; e) Formononetin; f) Equol

Table 2: Distance of Two nearby Atoms

Angle	Atom	Distance (Å)					
		Biochanin a	Daidzein	Glycitein	Formononetin	Genistein	Equol
40°	O-H(C2)	2,281	2,292	2,281	2,281	2,281	-
	H(C2)-H(C6)	2,029	1,994	2,029	2,029	2,029	-
80°	H(C2)-H(C6)	3,278	3,280	3,278	3,278	3,278	2,558
	O-H(C2)	3,423	3,394	3,423	3,423	3,423	1,701
140°	O-H(C6)	2,064	2,168	2,064	2,064	2,064	-
	H(C2)-H(C2)	2,004	2,007	2,004	2,004	2,004	1,730
	H(C3)-H(C2)	-	-	-	-	-	2,422
180°	O-H(C6)	1,821	1,814	1,821	1,821	1,821	-
	H(C2)-H(C2)	1,706	1,675	1,706	1,706	1,706	-
190°	O-H(C6)	-	-	-	-	-	1,842
	H(C3)-H(C2)	-	-	-	-	-	2,533
220°	O-H(C6)	2,064	2,069	2,064	2,064	2,064	-
	H(C2)-H(C2)	1,959	1,953	1,959	1,959	1,959	-
360°	O-H(C6)	3,007	3,003	3,007	3,007	3,007	-
	H(C2)-H(C2)	2,780	2,770	2,780	2,780	2,780	-
280°	H(C2)-H(C6)	-	-	-	-	-	1,900
	H(C3)-H(C2)	-	-	-	-	-	2,461
320°	O-H(C2)	2,128	2,107	2,128	2,128	2,128	-
	H(C2)-H(C6)	1,889	1,874	1,889	1,888	1,889	-
340°	O-H(C2)	-	-	-	-	-	1,560
	H(C2)-H(C6)	-	-	-	-	-	2,025

Table 3: Energy from Extreme Value of Isoflavon Derivatives in Soybean

Angle	Energy (kcal/mol)					
	Biochanin a	Daidzein	Glycitein	Formononetin	Genistein	Equol
40°	-3775,037	-3406,4587	-3778,365	-3675,219	-3506,286	-
80°	-3773,662	-3408,031	-3777,047	-3673,792	-3504,914	-3532,874
140°	-3774,984	-3406,137	-3778,101	-3674,006	-3506,235	-3529,888
180°	-3776,012	-3406,902	-3773,596	-3669,719	-3501,180	-
220°	-3774,745	-3406,125	-3778,357	-3674,885	-3505,952	-
260°	-3773,551	-3404,919	-3777,070	-3673,670	-3504,804	-
280°	-	-	-	-	-	-3531,685
320°	-3774,994	-3406,192	-3778,557	-3674,953	-3506,255	-
340°	-	-	-	-	-	-3530,504

Table 2 show the distance of nearby atoms in extreme values angles each derivatives compound of isoflavon in soybean. Nearby atoms was two neighboring atoms that occur repulsion energy between that atoms. The extreme values were happened because the interaction of the two nearby atoms and make the energy higher. The lower the strain energy, the more stable that compound. From Table 3, the most minimum energy of derivatives compound of isoflavon in soybean biochanin a, daidzein, glycitein, formononetin, genistein, and equol was at 40°.

The most maximum energy from that compound at 180° because when the torsion at 180°, two nearby atoms was very close that makes the structure more unstable. Besides, equol has the minimum energy at 280° and the most maximum energy at 140°. For example in biochanin a at 40°, the nearby atoms in H (C2)-H(C2') was 2,00438 Å. The nearby atoms at 180° in O-H(C6) was 1,82086 Å and H(C2)-H(2) was 1,70648 Å. Biochanin a, daidzein, glycitein, formononetin, genistein and equal have a nearby atoms that was affect the maximum and minimum energy. Hence, this was the proof that distance of two nearby atoms affects the structure stability and strain energy.

References

- Nijveldt BJ (2001) Flavonoids: a Review of Probable mechanism of Action and Potential Applications, *Am. J. Clin. Nutr.* 74:418-425.
- Yuniati L, Rollando R (2018) Isolation of antibacterial compounds from endophyte fungal of fusarium sp. In *phyllanthus niruri* linn. Leaves, *J. Pharm. Sci. & Res.* 10(2):260-264.
- Rollando R, Hedyotis corymbosa L, *Sterculia quadrifida* R (2018) Br Ethanollic Extract Enhances Cisplatin's Cytotoxicity on T47D Breast Cancer Cells Through Cell Cycle Modulation, *The Journal of Pure and Applied Chemistry Research*, 7(2):63-75.

Conclusion

The structural conformational study of derivatives compound of isoflavon in soybean was performed based on the desktop molecular modeling program by IS 1 semi empirical and selected. The distances of two nearby atoms affect the structure stability and the strain energy because the lower the strain energy, the more stable that compound.

In this study, the most stable structure for biochanin a, genistein, daidzein, glycitein, formononetin was 40° while the most stable structure for equol was 280°. From the angles, the next study can found the possibility of derivatives compound of isoflavon in soybean as a cancer drugs by molecular docking research. From that study, researchers no need to test every derivatives compound of isoflavon in soybean in laboratory and more efficient in time, money, substances, and energy.

Acknowledgements

We gratefully acknowledge the financial support from Program of Chemistry and Program of Pharmacy, Faculty of Science and Technology, Ma Chung University.

4. Arora A, Byrem TM, Nair MG, Strasburg GM (2000) Modulation of liposomal membrane fluidity by flavonoids and isoflavonoids. Arch Biochem Biophys, 373:102-109.
5. Zubik L, M Meydani (2003) Biosvability of soybean Isoflavon from Aglycone and Glucoside form in American Women, Am. J. Clin. Nutr. 77:1459-1465.
6. Yuan D, Yingni PAN, Chen Y, Uno T, Zhang S, Kano Y (2008) An Improved Method for Basic Hydrolysis of Isoflavone Malonyl glucosides and Quality Evaluation of Chinese Soy Materials. Chem. Pharm. Bull.56 (1):1-6.
7. Mahboub R (2014) Structural Conformational Study of Eugenol Derivatives Using Semiempirical Methods. Hindawi Publishing Corporation,1-5
8. Arora A, Byrem TM, Nair MG, Strasburg GM (2000) Modulation of liposomal membrane fluidity by flavonoids and isoflavonoids. Arch Biochem Biophys, 373:102-109.
9. Medjakovic S, Mueller M, Jugbauer A (2010) Potential Health-modulating Effects of Isoflavones and Metabolites via Activation of PPAR and AhR. Nutrients, 2:241-279.

Structural Conformational Study of Isoflavon Derivatives in Soybean

ORIGINALITY REPORT

11 %

SIMILARITY INDEX

10 %

INTERNET SOURCES

7 %

PUBLICATIONS

5 %

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

5%

★ www.jgpt.co.in

Internet Source

Exclude quotes On

Exclude bibliography On

Exclude matches < 5 words

Structural Conformational Study of Isoflavon Derivatives in Soybean

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
